

~~B physics~~ Heavy Flavour Results from D0

*X(4140), B_s Lifetime, B^+ F-B Asymmetry,
 D_s CP Violation, Dimuon Asymmetry*


Iain Bertram
Beauty 2014, Edinburgh
17 July 2014



Search for X(4140)




- $B^+ \rightarrow J/\psi \phi K^+$: resonance $X(4140) \rightarrow J/\psi \phi$
 - Standard quark model does not predict a state at this mass
 - Decay suggests cc, but mass is above open charm threshold
- Reconstruct $B^+ \rightarrow J/\psi \phi K^+$ (where $J/\psi \rightarrow \mu\mu$ and $\phi \rightarrow KK$)
 - veto $\psi(2S)$ and check for $J/\psi + K$ or π structures
 - Fit for B^+ yield in bins of $M(J/\psi KK)$





Search for X(4140)



- Evidence for X(4140) at 3.1σ

$$M = 4159.0 \pm 4.3 \pm 6.6 \text{ MeV}$$

$$\Gamma = 19.9 \pm 12.6^{+3.0}_{-8.0} \text{ MeV}$$

$$\frac{\mathcal{B}(B^+ \rightarrow X(4140)K^+)}{\mathcal{B}(B^+ \rightarrow J/\Psi \phi K^+)} = (19 \pm 7 \pm 4)\%$$

- Status
 - First evidence at CDF (Y(4140)) at 3.8σ
No evidence at Belle $\gamma\gamma \rightarrow J/\psi\phi$ in but higher mass state reported
 - No evidence at LHCb
 - Evidence at CMS at $>5\sigma$



B_s Lifetime



- Test theoretical predictions
 - Heavy Quark Expansion: $\tau(B_s^0)/\tau(B_d^0) = 1.00 \pm 0.01$
[Phys. Rev. D 70, 094031](#)
 - ... most recent, Lattice inputs: $= 1.001 \pm 0.002$
[Lenz review, arXiv:1405.3601](#)
 - Need for understanding of complex B_s^0 mixed system
- Target
 - Lifetimes of B_d^0 and B^+ measured to < 1% precision at B-factories
 - Updating latest DØ measurement (precision 3.7%) with full data set ($0.4 \rightarrow 10.4 \text{ fb}^{-1}$)



B_s Lifetime



- B_s⁰ lifetime depends on final state!
 - $\Delta m_s = B_s^L - B_s^H$ Mixing, mass eigenstates...
 - $\Delta \Gamma_s = \Gamma_s^L - \Gamma_s^H$... with different lifetimes
 - $\Delta \Gamma_s = \Gamma_s^{\text{CP-even}} - \Gamma_s^{\text{CP-odd}}$ if no CP violation
- Lifetimes
 - B_s⁰ $\rightarrow J/\Psi f_0(980)$ Pure CP-odd, Only single lifetime, $\Gamma_s^{\text{CP-odd}}, \Gamma_s^H$
 - B_s⁰ $\rightarrow K^+ K^-$ Pure CP-even, $\Gamma_s^{\text{CP-even}}, \Gamma_s^L$
 - B_s⁰ $\rightarrow D_s^- \mu^+ \nu$ Flavour specific, 50% CP-even, CP-odd at t=0
 - B_s⁰ $\rightarrow J/\Psi \phi$ complicated mix of CP-even and -odd, complex analysis to extract.

$$\Gamma_s = \frac{\Gamma_s^L + \Gamma_s^H}{2}$$

define

$$\bar{\tau}(B_s^0) = 1/\Gamma_s$$



B_s Lifetime

- B_s⁰ lifetime depends on final state!
 - Δm_s = B_s^L – B_s^H Mixing, mass eigenstates...
 - ΔΓ_s = Γ_s^L – Γ_s^H ... with different lifetimes
 - ΔΓ_s = Γ_s^{CP-even} – Γ_s^{CP-odd} if no CP violation
- Lifetimes
 - B_s⁰ → J/Ψf₀(980) Pure CP-odd, Only single lifetime, Γ_s^{CP-odd}, Γ_s^H
 - B_s⁰ → K⁺ K⁻ Pure CP-even, Γ_s^{CP-even}, Γ_s^L
 - B_s⁰ → D_s⁻μ⁺ν Flavour specific, 50% CP-even, CP-odd at t=0
 - B_s⁰ → J/ΨΦ complicated mix of CP-even and -odd, complex analysis to extract.

$$\Gamma_s = \frac{\Gamma_s^L + \Gamma_s^H}{2}$$

define


$$\bar{\tau}(B_s^0) = 1/\Gamma_s$$



B_s Reconstruction



- Reconstruct B_s^0 using D_s^- with opposite signed muon.
- Single and dimuon triggers
 - No IP based triggers
- Kinematic Requirements
 - μ^\pm with $p_T > 1.5 \text{ GeV}$ and $p_{\text{tot}} > 3.0 \text{ GeV}$
 - ϕ : K^\pm with $p_T > 1.0 \text{ GeV}$ and $1.08 \leq m(KK) \leq 1.32 \text{ GeV}$
 - D_s^- : π^\pm with $p_T > 0.7 \text{ GeV}$ and $1.6 \leq m(\phi\pi) \leq 2.3 \text{ GeV}$
 - B_s : $2.5 \leq m(\mu D_s) \leq 5.5 \text{ GeV}$






B_s Reconstruction



$$B_s^0 \rightarrow D_s^- \mu^+ \nu X$$
$$\downarrow \phi \pi^-$$
$$\downarrow K^+ K^-$$

Reconstruct a D_s^- associated
with a correct-sign muon






Likelihood



$$\mathcal{L} = \prod_{i \in \text{sig.sample}} [f_{\text{sig}} \mathcal{F}_{\text{sig}}^i + (1 - f_{\text{sig}}) \mathcal{F}_{\text{bckg}}^i] \prod_{i \in \text{bckg.sample}} \mathcal{F}_{\text{bckg}}^i$$

- f_{sig} from D_s mass fit, $\mathcal{F}_{\text{bckg}}$ Mass sidebands and WS signal.
- Lifetime models: convolved exponentials,
cc : Gaussians, combinatorial: multiple exponentials







Likelihood



$$\mathcal{L} = \prod_{i \in \text{sig.sample}} [f_{\text{sig}} \mathcal{F}_{\text{sig}}^i + (1 - f_{\text{sig}}) \mathcal{F}_{\text{bckg}}^i] \prod_{i \in \text{bckg.sample}} \mathcal{F}_{\text{bckg}}^i$$

- f_{sig} from D_s mass fit, $\mathcal{F}_{\text{bckg}}$ Mass sidebands and WS signal.
- Lifetime models: convolved exponentials,
cc : Gaussians, combinatorial: multiple exponentials




- Use $B_d \rightarrow D^- \mu^+ \nu X$ peak fit for ratio
 $R = \tau(B_s^0)_{\text{fs}} / \tau(B_d^0)$



Systematics



Uncertainty source	B_s^0 (μm)	B^0 (μm)	ΔR
Resolution Model	0.7	2.1	0.003
Combinatorial Background Model	5.0	4.9	0.001
K -factor determination	1.6	1.3	0.006
Non-Combinatorial Background	2.6	2.0	0.001
Signal Fraction	1.0	1.8	0.002
Alignment of the detector	2.0	2.0	0.000
Total	6.3	6.4	0.007






Results



$$\tau(B_s^0)_{\text{fs}} = 1.479 \pm 0.010 \pm 0.021 \text{ ps}$$

Contours of $\Delta(\log L) = 0.5$



$[\tau(B_s^0)/\tau(B_d^0)]^{\text{pred}}$ and WA of $\tau(B_d^0)$

$$\tau(B_s^0)_{\text{fs}} = \frac{1}{\Gamma_s} \frac{1 + (\Delta\Gamma_s/2\Gamma_s)^2}{1 - (\Delta\Gamma_s/2\Gamma_s)^2}$$


- Measurement precision better than previous world average



B_s Lifetime



$$R = \tau(B_s^0)_{\text{fs}} / \tau(B_d^0) = 0.964 \pm 0.013 \pm 0.007$$



- DØ working on $B_s^0 \rightarrow J/\psi f_0(980)$
- Pure CP-odd
- Stay tuned!


R Van Kooten




B_s Lifetime



$$R = \tau(B_s^0)_{\text{fs}} / \tau(B_d^0) = 0.964 \pm 0.013 \pm 0.007$$



R Van Kooten



Contours of $\ln(L)$:
39% CL for the en
68% CL for the ba

- NOT INCLU
- $B_s \rightarrow D_s\pi, K\pi$
- $B_s \rightarrow D_s D_s$
- $B_s \rightarrow J/\Psi (\pi\pi)$


Gray = combin



B $^\pm$ F-B Asymmetry



- Forward-backward asymmetry may probe for new physics.
- D0 uses $B^\pm \rightarrow J/\Psi K^\pm$ to probe asymmetry of b-quarks.





$$A_{FB} = \frac{N_F - N_B}{N_F + N_B}$$

Forward: b-quark in same direction as proton
anti-b in same direction as anti-proton



B^\pm F-B Asymmetry

- An unbinned maximum likelihood fit is used to extract the number of B meson decays in each category.
- Unblinded projections: [D0 Note 6441-CONF](#)



$$A_{FB} = [-0.26 \pm 0.41 \pm 0.17] \%$$





B $^\pm$ F-B Asymmetry



$$A_{FB} = [-0.26 \pm 0.41 \pm 0.17] \%$$

- Comparison with MC@NLO

$$A_{\text{MC@NLO}} = [1.63 \pm 0.43 \pm X.XX] \%$$



- Theory systematics to come: different PDFs, renorm. scale. etc.



Direct CPV in $D_s^\pm \rightarrow \phi\pi^\pm$



- Motivation:
 - Direct CP violation can occur if tree and loop (penguin) can interfere with different strong and weak phases
 - No CP violation is expected in decay $D_s^\pm \rightarrow \phi\pi^\pm$ (all process have same weak phase)
 - Non-zero value implies new physics
 - Motivation
 - Assume zero-CPV in many other analyses:



e.g.: $B_s^0 \rightarrow \bar{B}_s^0 \rightarrow D_s \mu\nu$	$\sigma(D_s^\pm)$
CPV in mixing	Production asymmetry (LHCb)
 - Experimentally measure
- $$A_{D_s} = \frac{N_{D_s^+} - N_{D_s^-}}{N_{D_s^+} + N_{D_s^-}},$$



Direct CPV in $D_s^\pm \rightarrow \phi\pi^\pm$



- Use similar techniques for CP asymmetries as other D \emptyset analyses
- $D_s^\pm \rightarrow \phi\pi^\pm \hookrightarrow K^+K^-$ Dominant kaon charge asymmetry \sim cancels!




$$A_{CP} = (-0.38 \pm 0.26 \pm 0.08)\%$$

- Most precise measurement
- Consistent with zero



Dimuon Charge Asymmetry



- CP violation *in mixing*: $\Gamma(B_{(s)}^0 \rightarrow \bar{B}_{(s)}^0 \rightarrow \mu^- X) \neq \Gamma(\bar{B}_{(s)}^0 \rightarrow B_{(s)}^0 \rightarrow \mu^- X)$
- Measure via

$$A = \frac{N(\mu^+ \mu^+) - N(\mu^- \mu^-)}{N(\mu^+ \mu^+) + N(\mu^- \mu^-)}$$

Inclusive single muons

$a^{\text{raw}} = \frac{n(\mu^+) - n(\mu^-)}{n(\mu^+) + n(\mu^-)}$

Constrain backg.
Reduce syst.

Mostly background

$$A_{sl}^b = \frac{N_b(\mu^+ \mu^+) - N_b(\mu^- \mu^-)}{N_b(\mu^+ \mu^+) + N_b(\mu^- \mu^-)}$$

Correct for backgrounds, fraction from b 's

- Asymmetry is a linear combination semileptonic charge asymmetries of B_d^0 and B_s^0

$$A_{sl}^b = C_d a_{sl}^d + C_s a_{sl}^s ;$$

$$a_{sl}^b = \frac{\Gamma(\bar{B} \rightarrow \mu^+ X) - \Gamma(B \rightarrow \mu^- X)}{\Gamma(\bar{B} \rightarrow \mu^+ X) + \Gamma(B \rightarrow \mu^- X)}$$




Dimuon: 2011 Result



$$A_{sl}^b = (-0.787 \pm 0.172(\text{stat}) \pm 0.093(\text{syst})) \%$$

- $9.0 \text{ fb}^{-1} \rightarrow 10.4 \text{ fb}^{-1}$
- More detailed study of asymmetry dependence on impact parameter (IP) on each muon
- More detailed study of asymmetry dependence on muon (p_T, η)
- Another cross check using independent alternative way to measure background
- Additional CP-violating process included to interpret result

CPV in interference of decays w/ and w/o mixing & special decay class



PRD 84 052007 (2011)




Dimuon Charge Asymmetry



IP Sample	Muon IP (μm)
IP=1	0-50
IP=2	50-120
IP=3	120-3000



More backgrounds due to $K \rightarrow \mu, \pi \rightarrow \mu$ which result in an asymmetry since $\sigma(K^-N) > \sigma(K^+N)$



$$a_{\text{CP}} = a_{\text{raw}} - a_{\text{det}}$$

Similar result if bin in (η, p_T)
where there are 27 independent bins




Dimuon Charge Asymmetry




IP Sample	Muon IP (μm)
IP=1	0-50
IP=2	50-120
IP=3	120-3000



More backgrounds due to $K \rightarrow \mu, \pi \rightarrow \mu$ which result in an asymmetry since $\sigma(K^-N) > \sigma(K^+N)$



Similar result if bin in (η, p_T)
where there are 27 independent bins





Dimuon Charge Asymmetry



IP Sample	Muon IP (μm)
IP=1	0-50
IP=2	50-120
IP=3	120-3000

Using same sign
dimuons


$$A_{\text{CP}} = (-0.235 \pm 0.064 \pm 0.055)\%$$

3.6σ deviation from SM

$$A_{\text{CP}}^{\text{mix}}(\text{SM}) + A_{\text{CP}}^{\text{int}}(\text{SM})$$

$$(-0.8 \pm 0.1) \times 10^{-4} + (-3.5 \pm 0.8) \times 10^{-4}$$

PRD 87, 074020 (2013)






Dimuon Charge Asymmetry



- Interpretation & results
 - Fractional mix of B_s and B_d in each IP bin and A_{CP} proportional to $\Delta\Gamma_d$
- $a_{sl}^s = (-0.82 \pm 0.99) \%$,
 $a_{sl}^d = (-0.62 \pm 0.43) \%$,
 $\Delta\Gamma_d/\Gamma_d = (+0.50 \pm 1.38) \%$,
 $\rho_{s,d} = -0.61$, $\rho_{d,\Delta\Gamma} = -0.03$,
 $\rho_{s,\Delta\Gamma} = +0.66$.

3.0 σ deviation from SM of three values

Result consistent with D0
measurements of a_{sl}^s and a_{sl}^d






Dimuon Charge Asymmetry



- Interpretation & results
 - Sensitive to $\Delta\Gamma_d/\Gamma_d$
 - Fix to WA $\Delta\Gamma_d/\Gamma_d = 0.015$
 - Deviation now only 1.9σ






Dimuon Charge Asymmetry



- Average of all three D0 semi-leptonic charge asymmetries

$$\begin{aligned}
 a_{\text{sl}}^s &= (-1.33 \pm 0.58)\%, \\
 a_{\text{sl}}^d &= (-0.09 \pm 0.29)\%, \\
 \Delta\Gamma_d/\Gamma_d &= (+0.79 \pm 1.15)\%, \\
 \rho_{s,d} &= -0.34, \quad \rho_{d,\Delta\Gamma} = +0.24, \\
 \rho_{s,\Delta\Gamma} &= +0.55.
 \end{aligned}$$

3.1σ deviation from SM





Summary




- DØ has a well understood detector & dataset with well developed analysis techniques.
 - small levels of pile-up
 - p-anti-p CP symmetric initial state
 - regular flipping of magnet polarities
- Still producing results with LHC in niche areas.
 - new tests of CPV and FB asymmetries
 - Leaving Dimuon charge asymmetry puzzle
 - New Physics
 - Is $\Delta\Gamma_d/\Gamma_d$ the solution?

Backup Slides



Backup B_s Lifetime



If fit with a single exponential, measure:

$$\tau(B_s^0)_{fs} = \frac{1}{\Gamma_s} \frac{1 + (\Delta\Gamma_s / 2\Gamma_s)^2}{1 - (\Delta\Gamma_s / 2\Gamma_s)^2}$$

Difficult to distinguish two exponentials
in a stable fit...




B_s Reconstruction



$$\begin{aligned} B_s^0 &\rightarrow D_s^- \mu^+ \nu X \\ &\downarrow \phi \pi^- \\ &\downarrow K^+ K^- \end{aligned}$$

Reconstruct a D_s^- associated
with a correct-sign muon





B_s Lifetime Systematics




- Decay length resolution
Replace double-Gaussian model with single + exponential tails
- Combinatorial Background
Use single samples (each of mass side-bands, wrong-sign)
- K factors
Use different MC, vary composition and relevant lifetimes within uncertainties
- Non-Combinatorial Background
Vary composition within uncertainties
- Detector Alignment
Use different silicon microvertex detector alignment files with sensors moved within uncertainties
- Signal fraction
Varied within uncertainties from mass fit, different mass models



B $^\pm$ F-B Asymmetry



- Correlation between parent b-quark and reconstructed B meson from MC@NLO. About 80% of the time, the B meson tracks the parent b-quark.





B^\pm F-B Asymmetry




BDT Variations	0.14%
Fit Variations	0.080%
Polarity Weighting	0.0001%
Detector Asymmetries	0.058%
Systematic	0.17%
Statistical	0.41%
Total	0.44%


Determination of the raw asymmetries: $D^\pm \rightarrow \phi\pi^\pm$

$L=3\text{fb}^{-1}$

$D^+ \rightarrow \phi\pi^+$



$D^- \rightarrow \phi\pi^-$



— signal
— combinatorial bkg
bkg

Physics Letters B 713 (2012) 186–195

$$A_P = \frac{\sigma(D_s^+) - \sigma(D_s^-)}{\sigma(D_s^+) + \sigma(D_s^-)} = (-0.33 \pm 0.22 \pm 0.10)\%.$$

-018
D

Decay Mode	Yield
$D^\pm \rightarrow K_S^0\pi^\pm$	$4\,834\,440 \pm 2\,555$
$D_s^\pm \rightarrow K_S^0\pi^\pm$	$120\,976 \pm 692$
$D^\pm \rightarrow K_S^0K^\pm$	$1\,013\,516 \pm 1\,379$
$D_s^\pm \rightarrow K_S^0K^\pm$	$1\,476\,980 \pm 2\,354$
$D^\pm \rightarrow \phi\pi^\pm$	$7\,020\,160 \pm 2\,739$
$D_s^\pm \rightarrow \phi\pi^\pm$	$13\,144\,900 \pm 3\,879$


[%]	Total
$\mathcal{A}_{\text{meas}}^{D^\pm \rightarrow K_S^0\pi^\pm}$	-0.95 ± 0.05
$\mathcal{A}_{\text{meas}}^{D_s^\pm \rightarrow K_S^0\pi^\pm}$	-0.15 ± 0.46
$\mathcal{A}_{\text{meas}}^{D^\pm \rightarrow K_S^0K^\pm}$	$+0.01 \pm 0.19$
$\mathcal{A}_{\text{meas}}^{D_s^\pm \rightarrow K_S^0K^\pm}$	$+0.27 \pm 0.11$
$\mathcal{A}_{\text{meas}}^{D_s^\pm \rightarrow \phi\pi^\pm}$	-0.41 ± 0.05



B^\pm F-B Asymmetry




- Detector Asymmetry



Additional Source of CPV in Like-Sign Dimuons

Borissov, Hoeneisen, arXiv:1303.0175v1 [hep-ex],
Understanding the like-sign dimuon charge asymmetry in pp(bar) collisions



e.g.,

$\Gamma \rightarrow \mu^- X$ $B^- B^0$ $\hookrightarrow D^+ D^-$ $\hookrightarrow \mu^- X$	$\Gamma \rightarrow \mu^+ X$ $B^+ \bar{B}^0$ $\hookrightarrow D^+ D^-$ $\hookrightarrow \mu^+ X$
---	---

but due to interference between mixing and decay in B system:

$$\Gamma(B^0 \rightarrow D^+ D^-) \neq \Gamma(\bar{B}^0 \rightarrow D^+ D^-) \quad \mathcal{A} = -\sin(2\beta) \frac{x_d}{1 + x_d^2}$$

$$\mathcal{A}_{CP}^{mix}(SM) = (-0.8 \pm 0.1) \times 10^{-4}$$

$$\mathcal{A}_{CP}^{int}(SM) = (-3.5 \pm 0.8) \times 10^{-4} \quad \leftarrow \text{additional}$$




Semi-Leptonic CPV WA



- Direct Measurements of $a_{\text{SI}}^d = (+0.23 \pm 0.26)\%$
 - Previous B-factory results HFAG arXiv:1207.1158 $(-0.05 \pm 0.56)\%$
 - DØ: PRD 86, 072009 (2012) $(+0.68 \pm 0.47)\%$
 - BaBar: PRL 111, 101802 (2013) $(+0.06 \pm 0.38)\%$
- Direct Measurements of $a_{\text{SI}}^s = (+0.50 \pm 0.52)\%$
 - DØ: PRL 110, 011801 (2013) $(-1.12 \pm 0.76)\%$
 - LHCb: PLB 728C (2014) $(-0.06 \pm 0.63)\%$
- Direct Measurements of $\Delta\Gamma_d/\Gamma_d = (-0.4 \pm 2.0)\%$
 - Previous B-factory results, HFAG, arXiv:1207.1158 $(+1.15 \pm 1.80)\%$
(Belle, BaBar, [DELPHI])
 - LHCb: JHEP04(2014)114 $(-4.4 \pm 2.7)\%$
- Dimuon Charge Asymmetry (see talk)




Semi-Leptonic CPV WA



$$a_{sl}^s = (-0.92 \pm 0.43)\%,$$

$$a_{sl}^d = (-0.11 \pm 0.21)\%,$$

$$\Delta\Gamma_d/\Gamma_d = (+1.09 \pm 0.93)\%,$$



$$\rho_{s,d} = -0.24, \quad \rho_{d,\Delta\Gamma} = +0.23, \quad \rho_{s,\Delta\Gamma} = +0.48.$$

$$\chi^2(\text{comb}) = 4.98/3\text{d.o.f.}$$

2.8 σ deviation from SM