

# Rubisco activation limits photosynthesis in wheat



Samuel H. Taylor & Stephen P. Long: Lancaster Environment Centre

## Rationale:

Light available for photosynthesis fluctuates continuously in the field, as clouds cross the sun and as the movement of the sun causes shadows to move across leaves. Transgenic manipulations to allow more rapid relaxation of non-photochemical quenching during sun-shade transitions increased productivity in the field by 14-20% [1]. Rubisco activation is a key limit on photosynthesis during induction following shade-sun transitions [2]. This suggests there may also be potential to increase photosynthesis and crop productivity by speeding up Rubisco activation.

## Key findings:

- 1) Dynamic  $A/c_i$  analysis shows that  $V_{\text{cmax}}$  is the slowest relaxing biochemical limitation during photosynthetic induction in flag-leaves of wheat
- 2) Modelling diurnal  $\text{CO}_2$  assimilation using photosynthetic light responses and kinetics for  $V_{\text{cmax}}$  shows that  $V_{\text{cmax}}$  kinetics limit flag leaf photosynthesis by as much as 21%

## Dynamic $A/c_i$ analysis

- Flag leaf photosynthesis of wheat, cv. Highbury, was carboxylation-limited at steady state (Fig. 1).
- $A/c_i$  response curves at 10 s time intervals showed that photosynthesis was limited by  $V_{\text{cmax}}$  for longer than  $J$  (Fig. 2).
- Time constants were obtained for increases in  $V_{\text{cmax}}$  in the light:  $\tau_{\text{act}} \approx 180$  s; and decreases in the shade,  $\tau_{\text{de-act}} \approx 300$  s.

Fig. 1  $A/c_i$  responses illustrating biochemical limitations to photosynthesis in flag leaves of wheat, cv. Highbury. Carboxylation-limited region (shaded), and  $A/c_i$  components ( $A_c$ ,  $A_i$ ) are shown for steady state conditions (PPFD,  $1200 \mu\text{mol m}^{-2} \text{s}^{-1}$ ;  $T_{\text{leaf}}$ ,  $25^\circ\text{C}$ ; and  $\text{VPD}_{\text{leaf}}$ ,  $1 \text{ kPa}$ ) and snapshots 2, 4 and 8 min after transition from shade (30 min,  $50 \mu\text{mol m}^{-2} \text{s}^{-1}$  PPFD) to saturating light ( $1200 \mu\text{mol m}^{-2} \text{s}^{-1}$ ). Dynamic responses were obtained following Soleh et al. [3], by repeating measurements at  $[\text{CO}_2]$  of 50, 100, 200, 300, 400, 500, 600, 800 and  $1000 \mu\text{mol mol}^{-1}$ .





Fig. 2 Dynamic response to shade-sun transitions for biochemical factors limiting photosynthesis in flag leaves of wheat, cv. Highbury. Rubisco carboxylation rate,  $V_{\text{cmax}}$ , and electron transport rate,  $J$ , determined at 10 s intervals after transition from 30 mins deep shade (PPFD,  $50 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) to saturating light ( $1200 \mu\text{mol m}^{-2} \text{s}^{-1}$ ). Activation and de-activation time constants ( $\tau$ ) for  $V_{\text{cmax}}$  were determined using the exponential  $y = y_{\text{sat}} - (y_{\text{sat}} - y_0)e^{-t/\tau}$  [4].

## Impact on diurnal $\text{CO}_2$ assimilation

Impacts of  $V_{\text{cmax}}$  kinetics on gross  $\text{CO}_2$  assimilation at steady state  $c_i$  ( $A^*$ ) were estimated by comparing two scenarios:

- 1) immediate responses of  $A^*$  to PPFD; and 2)  $A^*$  responses with kinetics similar to  $V_{\text{cmax}}$ .
- $V_{\text{cmax}}$  kinetics decreased diurnal photosynthesis by 21% (Fig. 3).




Fig. 3 Impact of Rubisco activation kinetics on integrated gross  $\text{CO}_2$  assimilation ( $A^*$ ). (A) A minute-by-minute simulation of PPFD in the second layer of a crop canopy on a clear-sky day [5] and (B) photosynthetic light response curves, were used to model (C)  $A^*$  during a diurnal period, and (D) the impact on cumulative potential  $\text{CO}_2$  uptake ( $\Sigma A^*$ ).

$A^*$  was integrated following [4], as  $\bar{A}^* = A_f^* t - (A_f^* - A_i) \tau + (A_f^* - A_i) \tau e^{-t/\tau}$ .  $A_f^*$  and  $A_i$  are predicted final and initial values for  $A^*$ .  $A_i$  is initially 0, then  $A_i = A_f^* - (A_f^* - A_i) e^{-t/\tau}$  from the previous interval  $A_f^* = \phi I + A_{\text{sat}} - \sqrt{(\phi I + A_{\text{sat}})^2 - 4\phi I A_{\text{sat}}}/2\theta$  (non-rectangular hyperbola) fit to photosynthetic light response curves (B).  $\tau$  is the relevant time constant scenario 1:  $\tau = 0$  scenario 2:  $\tau = \tau_{\text{act}}$  with  $\uparrow$  PPFD, or  $\tau_{\text{de-act}}$  with  $\downarrow$  PPFD  $t$  is the time interval ( $\sim 60$  s).



## References

1. Kromdijk et al. 2016 Science 354:857
2. Kaiser et al. 2015 J Exp Bot 66:2415
3. Soleh et al. 2016 Plant Cell Environ 39:685
4. Mott & Woodrow 2000 J Exp Bot 51:399
5. Zhu et al. 2004 J Exp Bot 55:1167

## Acknowledgements

Dr. Elizabete Carmo-Silva shared the cv. Highbury plants. This research was supported by LEC, Lancaster University

Forthcoming publication Taylor SH & Long SP (in press) Philos T Roy Soc B