

MAPPING CONES IN THE BOUNDED DERIVED CATEGORY OF A GENTLE ALGEBRA

İLKE ÇANAKÇI, DAVID PAUKSZTELLO, AND SIBYLLE SCHROLL

ABSTRACT. In this article we describe the triangulated structure of the bounded derived category of a gentle algebra by describing the triangles induced by the morphisms between indecomposable objects in a basis of their Hom-space.

INTRODUCTION

Derived categories provide a common framework for homological algebra in subjects such as algebra, geometry and mathematical physics. For example, in mathematical physics, in the context of homological mirror symmetry, they are the natural setting for Bridgeland's stability conditions [11]. In algebraic geometry, they arise in the study of non-commutative crepant resolutions which are often studied via an algebra whose derived category is equivalent to the derived category of the smooth variety resolving the singularity [47]. In representation theory, derived categories are the natural setting for tilting theory of finite dimensional algebras, see for example [34]. Thus understanding the structure of derived categories and their properties is an important problem. However, owing to their complexity, in general, this is difficult to achieve. Therefore in the cases, where this is achievable, it is of great value to obtain as much detailed knowledge of the derived category as possible.

In the context of the representation theory of finite dimensional algebras, we will now describe a situation where it is possible to gain insight into the structure of derived categories. According to the tame-wild dichotomy [26], algebras are either of tame representation type, that is, all indecomposable finite-dimensional modules may be classified, or they are of wild representation type and it is considered that a complete classification is impossible. Therefore much of the work in representation theory has been focused on tame algebras and for particular classes of tame algebras we have a good understanding of their (classical) representation theory, that is of their module categories. A good example of this are special biserial algebras [48], a prominent class of tame algebras, that has been widely studied, with many exciting recent developments, see for example [1, 2, 3, 21, 23, 24, 27, 29, 46, 49]. One of the reasons that special biserial algebras are so well-understood is that their indecomposable representations are classified in terms of strings and bands [32, 43, 48].

The notion of derived-tameness was introduced in [30]. In the case of derived-tame algebras we can gain concrete insight into the structure of the derived category. Gentle algebras are derived tame [8]. They form a subclass of special biserial algebras which is closed under derived equivalence [45]. They are, therefore, a natural class of algebras whose derived category is the object of intensive study. As a result, our understanding of the structure of derived categories of gentle algebras parallels that of our understanding of their module categories.

Gentle algebras first arose in the setting of tilting theory in the classification of iterated tilted algebras of type A and type \tilde{A} in [5] and [7] respectively. They now play an important role in many areas of mathematics: in algebra, they occur in cluster theory as Jacobian algebras associated to surface triangulations [6, 28, 39], and in recent advances in invariant theory [22]. In addition to their widespread appearance in representation theory and algebra, gentle algebras

2010 *Mathematics Subject Classification.* 18E30, 16G10, 05E10.

Key words and phrases. bounded derived category, gentle algebra, homotopy string and band, string combinatorics, mapping cone

This work was supported through EPSRC grants EP/K026364/1, EP/K022490/1, EP/N005457/1 and EP/P016294/1.

also are increasingly ubiquitous in geometric contexts. For example, their singularity category [14] – which measures how far an algebra or variety is away from being nonsingular – was described in [37]. They feature prominently in the programme to understand singularities of nodal curves [15, 16, 17] and in an algebraic approach to mirror symmetry [10, 33, 38, 41].

From now on let Λ be a gentle algebra and $D^b(\Lambda)$ be its bounded derived category with shift functor Σ . The indecomposable objects in $D^b(\Lambda)$ have been classified [8] in terms of string combinatorics: namely they are given in terms of *homotopy strings* and *homotopy bands*, the terminology originating in [9]; see also [15] for a similar approach in the context of nodal algebras. The corresponding indecomposable complexes are called *string complexes* and *band complexes*, respectively.


Using the Happel functor [34], the Auslander–Reiten (AR) structure of the perfect category $K^b(\text{proj}(\Lambda))$ was determined in [9]. See also [4, §6] for similar results without the use of the Happel functor. A canonical basis for the morphisms between string complexes and (one-dimensional) band complexes in $D^b(\Lambda)$ was given in [4] in terms of three types of morphism: *graph maps*, *single maps* and *double maps*. We will call this the *standard basis*.

A general description of the middle terms of triangles of the form $Q^\bullet \rightarrow E^\bullet \rightarrow P^\bullet \rightarrow \Sigma Q^\bullet$, where P^\bullet and Q^\bullet are indecomposable objects, is not known. In this article, using (homotopy) string combinatorics, we describe the mapping cones of elements of the standard basis for the morphisms between string complexes and (one-dimensional) band complexes. As such, we explicitly describe the middle terms of extensions for a standard basis of $\text{Ext}_{D^b(\Lambda)}^1(P^\bullet, Q^\bullet) := \text{Hom}_{D^b(\Lambda)}^1(P^\bullet, \Sigma Q^\bullet)$ where P^\bullet and Q^\bullet are string or band complexes. Understanding middle terms of extensions is important for many applications, for example: in cluster theory, these form the basis of the so-called exchange triangles and exchange relations [13, 18, 20, 35, 36]; understanding middle terms of extensions is a key ingredient in classifying torsion pairs in triangulated categories [25, 31, 40, 42, 50]; the description of middle terms of extensions in the derived category of derived-discrete algebras is instrumental in the classification of thick subcategories of discrete derived categories [12] and they have been used in [19] to show that the short exact sequences given in [44] form a basis of the Ext^1 -space between any two indecomposable Λ -modules, thus answering this longstanding open question. Moreover, the results of this paper have been applied in [41] to give a geometric interpretation of mapping cones in terms of resolutions of crossing curves in the context of a geometric model of the derived category of a gentle algebra.


In $D^b(\Lambda)$ the middle term E^\bullet of an extension $Q^\bullet \rightarrow E^\bullet \rightarrow P^\bullet \xrightarrow{f^\bullet} \Sigma Q^\bullet$ is given by the inverse shift of the mapping cone $M_{f^\bullet}^\bullet$ of the map f^\bullet . In this paper, we describe the indecomposable summands of the mapping cones of the standard basis elements, that is of maps f^\bullet in the canonical basis of $\text{Hom}_{D^b(\Lambda)}(P^\bullet, Q^\bullet)$, where P^\bullet and Q^\bullet are indecomposable complexes. Our description is general in that our results cover both string and band complexes. Our main results can be summarised in the following graphical description of the mapping cones of standard basis elements:

Theorem. *Let Λ be a gentle algebra. Suppose Q_σ^\bullet and Q_τ^\bullet are indecomposable objects in $D^b(\Lambda)$, where σ and τ are the corresponding homotopy strings or bands. Let $f^\bullet \in \text{Hom}_{D^b(\Lambda)}(Q_\sigma^\bullet, Q_\tau^\bullet)$ be a standard basis element. Then the indecomposable summands of the mapping cone $M_{f^\bullet}^\bullet$ are given by the homotopy strings and bands occurring in the green and red boxes resulting from the following graphical calculus.*


- (1) Let $\sigma = \beta\sigma_L\rho\sigma_R\alpha$ and $\tau = \delta\tau_L\rho\tau_R\gamma$ and suppose f^\bullet is a graph map corresponding to the overlap ρ . Then $M_{f^\bullet}^\bullet$ is given by:



(2) Let $\sigma = \beta\sigma_L\sigma_R\alpha$ and $\tau = \delta\tau_L\tau_R\gamma$ and suppose f^\bullet is a single map. Then M_{f^\bullet} is given by:



(3) Let $\sigma = \beta\sigma_L\sigma_C\sigma_R\alpha$ and $\tau = \delta\tau_L\tau_C\tau_R\gamma$ and suppose f^\bullet is a double map. Then M_{f^\bullet} is given by:



We refer the reader to Sections 2, 3 and 4 for precise statements and details.

Convention. Throughout this article, all modules will be left modules and all maps will be composed from left to right.

1. BACKGROUND

1.1. The homotopy category and mapping cones. The required background on derived and triangulated categories in the setting of representation theory can be found in [34].

Let Λ be a finite-dimensional algebra over an algebraically closed field \mathbf{k} . The category of interest in this article will be $\mathcal{K}^{b,-}(\text{proj}(\Lambda))$, the homotopy category of right bounded complexes of projective left Λ -modules with bounded cohomology. The morphisms of $\mathcal{K}^{b,-}(\text{proj}(\Lambda))$ are cochain maps of complexes up to homotopy. It is well known that $\mathcal{K}^{b,-}(\text{proj}(\Lambda)) \simeq \mathcal{D}^b(\Lambda)$, the bounded derived category of finitely generated left Λ -modules and for convenience we shall always identify $\mathcal{D}^b(\Lambda)$ with $\mathcal{K}^{b,-}(\text{proj}(\Lambda))$.

Definition 1.1. Let (P^\bullet, d_P) and (Q^\bullet, d_Q) be complexes in $\mathcal{K}^{b,-}(\text{proj}(\Lambda))$ and suppose $f^\bullet: P^\bullet \rightarrow Q^\bullet$ is a cochain map. The *mapping cone of f^\bullet* is the complex $(M_{f^\bullet}^\bullet, d_{M_{f^\bullet}^\bullet})$ given by

$$M_{f^\bullet}^n = P^{n+1} \oplus Q^n \quad \text{and} \quad d_{M_{f^\bullet}^\bullet}^n = \begin{bmatrix} -d_P^{n+1} & f^{n+1} \\ 0 & d_Q^n \end{bmatrix}$$

1.2. Gentle algebras. The definition of a gentle algebra goes back to [7], where they first occurred as iterated tilted algebras of type A .

Definition 1.2. A finite-dimensional \mathbf{k} -algebra is called *gentle* if it is Morita equivalent to an algebra $\mathbf{k}Q/I$ where Q is a quiver and I an admissible ideal in $\mathbf{k}Q$ such that

- (1) For each vertex $i \in Q_0$ there are at most two arrows starting at i and at most two arrows ending at i ;
- (2) For each arrow $a \in Q_1$ there is at most one arrow b such that $ba \notin I$ and at most one arrow c such that $ac \notin I$;
- (3) For each arrow $a \in Q_1$ there is at most one arrow b with $s(b) = e(a)$ and $ba \in I$ and at most one arrow c with $e(c) = s(a)$ and $ac \in I$;
- (4) The ideal I is generated by length two monomial relations.

Note that if one removes condition (3) and relaxes condition (4) so that I is generated simply by monomial relations one obtains a so-called *string algebra*.

From now on, Λ will be a gentle algebra.

1.3. String and band complexes. The indecomposable complexes in $D^b(\Lambda)$ are parametrised by homotopy strings and bands. The original reference is [8] but we use the notation and terminology of [4] dropping some of the formality regarding cohomological degrees. All modules will be finitely generated left modules and therefore paths in the quiver will be read from right to left. For each $a \in Q_1$ we define a formal inverse arrow $\bar{a} = a^{-1}$ with $s(\bar{a}) = e(a)$ and $e(\bar{a}) = s(a)$. For a path $p = a_n \cdots a_1$ in (Q, I) the inverse path is $\bar{p} = \bar{a}_1 \cdots \bar{a}_n$.

Definitions 1.3. We recall the following definitions.

- (1) A *walk* in (Q, I) is a sequence $w = w_l \cdots w_1$ satisfying $s(w_{i+1}) = e(w_i)$, where each w_i is either an arrow or an inverse arrow, and where the sequence does not contain any subsequence of the form $a\bar{a}$ or $\bar{a}a$ for some $a \in Q_1$.
- (2) A (*finite*) *homotopy string* is a walk of finite length in (Q, I) . In addition there are *trivial homotopy strings* for each vertex $x \in Q_0$.
- (3) A subwalk $p = w_j \cdots w_i$ of a homotopy string $\sigma = w_l \cdots w_1$ is a *homotopy letter* if
 - (a) p or \bar{p} is a path in (Q, I) ; and,
 - (b) $w_i, \bar{w}_{i-1} \in Q_1$, or $\bar{w}_i, w_{i-1} \in Q_1$, or $w_i w_{i-1} \in I$, or $\bar{w}_{i-1} \bar{w}_i \in I$; and,
 - (c) $w_j, \bar{w}_{j+1} \in Q_1$, or $\bar{w}_j, w_{j+1} \in Q_1$, or $w_{j+1} w_j \in I$, or $\bar{w}_j \bar{w}_{j+1} \in I$.

The subwalk p is a *direct homotopy letter* if it is a path in (Q, I) and an *inverse homotopy letter* if \bar{p} is a path in (Q, I) . This partitions a homotopy string σ into homotopy letters and we write $\sigma = \sigma_n \cdots \sigma_1$ for this decomposition. A *homotopy subletter* of p is a subwalk of p .

- (4) A *trivial homotopy letter* is one of the form $p = 1_x$ for some $x \in Q_0$. Sometimes we shall assert the nonexistence of homotopy letters, and in this case we write $p = \emptyset$.
- (5) Let $\sigma = \sigma_n \cdots \sigma_1$ be a homotopy string decomposed into its homotopy letters. A subwalk $\tau = \sigma_j \cdots \sigma_i$ with $1 \leq i \leq j \leq n$ is called a *homotopy substring* of σ .
- (6) A *homotopy band* is a homotopy string $\sigma = \sigma_n \cdots \sigma_1$ with $s(\sigma) = e(\sigma)$, $\sigma_1 \neq \bar{\sigma}_n$, $\sigma \neq \tau^m$ for some homotopy substring τ and $m > 1$, and σ has equal numbers of direct and inverse homotopy letters.
- (7) A (possibly infinite) walk $w = w_n \cdots w_1$ is called a *direct antipath* if each $w_i \in Q_1$ is a direct homotopy letter; it is called an *inverse antipath* if each $\bar{w}_i \in Q_1$ and each w_i is an inverse homotopy letter.
- (8) A left infinite walk $w = \cdots w_n \cdots w_2 w_1$ is a *left infinite homotopy string* if there exists $m \geq 1$ such that $v = \cdots w_n \cdots w_{m+1} w_m$ is a direct antipath.
- (9) A right infinite walk $w = w_{-1} w_{-2} \cdots w_{-n} \cdots$ is a *right infinite homotopy string* if there exists $m \geq 1$ such that $v = w_{-m} w_{-m-1} \cdots w_{-n} \cdots$ is an inverse antipath.
- (10) A two-sided infinite walk $w = \cdots w_2 w_1 w_0 w_{-1} \cdots$ is called a *two-sided infinite homotopy string* if there exist integers $n > m$ such that $\cdots v_{n+1} v_n$ is a direct antipath and $v_m v_{m-1} \cdots$ is an inverse antipath.
- (11) A two-sided, left or right infinite homotopy string is called an *infinite homotopy string*.

Let σ be a (possibly infinite) homotopy string. Modulo the equivalence relation $\sigma \sim \bar{\sigma}$, the set of homotopy strings determine the *string complexes* P_σ^\bullet . Up to the equivalence relation given by inversion and cyclic permutation, the homotopy bands together with scalars $\lambda \in \mathbf{k}^*$ and strictly positive integers n determine the *band complexes* $B_{\sigma, \lambda, n}^\bullet$. By [8, Thm. 3], the string and band complexes form a complete set of indecomposable complexes in $K^{b,-}(\text{proj}(\Lambda))$ up to isomorphism. We refer to [8] for precise details of the construction of the complexes P_σ^\bullet and $B_{\sigma, \lambda, n}^\bullet$ from a homotopy string or band σ . The indecomposable perfect complexes are precisely the band complexes and the string complexes arising from finite homotopy strings. In what follows, we will only consider band complexes $B_{\sigma, \lambda, 1}^\bullet$ which we will write as $B_{\sigma, \lambda}^\bullet$. When we do not wish to explicitly identify whether an indecomposable complex is a string or band complex we use the notation

$$Q_\sigma^\bullet = \begin{cases} P_\sigma^\bullet & \text{if } \sigma \text{ is a (possibly infinite) homotopy string;} \\ B_{\sigma, \lambda}^\bullet & \text{if } \sigma \text{ is a homotopy band.} \end{cases}$$

Let $\sigma = \sigma_n \cdots \sigma_1$ be a homotopy string or band. We shall make frequent use of the following *unfolded diagram* notation from [4]:

$$\bullet \xrightarrow{\sigma_n} \bullet \cdots \bullet \xrightarrow{\sigma_2} \bullet \xrightarrow{\sigma_1} \bullet .$$

1.4. The standard basis. Let σ and τ be homotopy strings or bands. The main theorem of [4] establishes a basis for $\text{Hom}_{\mathbb{D}^b(\Lambda)}(Q_\sigma^\bullet, Q_\tau^\bullet)$ which we recall here.

Theorem 1.4 ([4, Theorem 3.15]). *Let σ and τ be homotopy strings or bands. Then there is a basis of $\text{Hom}_{\mathbb{D}^b(\Lambda)}(Q_\sigma^\bullet, Q_\tau^\bullet)$ given by:*

- graph maps $f^\bullet: Q_\sigma^\bullet \rightarrow Q_\tau^\bullet$;
- singleton single maps $f^\bullet: Q_\sigma^\bullet \rightarrow Q_\tau^\bullet$;
- singleton double maps $f^\bullet: Q_\sigma^\bullet \rightarrow Q_\tau^\bullet$;
- quasi-graph maps $\varphi: Q_\sigma^\bullet \rightsquigarrow \Sigma^{-1}Q_\tau^\bullet$.

A quasi-graph map is not a map but in fact determines a homotopy class of single and double maps, hence we use the notation \rightsquigarrow . We now briefly recap the types of maps occurring in Theorem 1.4.

1.4.1. Graph maps. Suppose σ and τ are homotopy strings or bands of the form,

$$\sigma = \beta\sigma_L\rho\sigma_R\alpha \text{ and } \tau = \delta\tau_L\rho\tau_R\gamma, \text{ or } \sigma = \beta\sigma_L\rho \text{ and } \tau = \delta\tau_L\rho,$$

where α, β, γ and δ are homotopy substrings, $\sigma_L, \sigma_R, \tau_L$ and τ_R are homotopy letters (possibly empty, in which case the corresponding homotopy substring α, β, γ or δ would be empty as well), and ρ is a (possibly trivial) maximal common homotopy substring, and in the second case an infinite homotopy substring of σ and τ . Moreover, we assume that ρ occurs in the same cohomological degrees in both homotopy strings. These setups are indicated in the following unfolded diagrams of Q_σ^\bullet and Q_τ^\bullet :

$$(1) \quad \begin{array}{c} Q_\sigma^\bullet: \\ \quad \beta \quad \sigma_L \quad \rho_k \quad \rho_{k-1} \quad \dots \quad \rho_2 \quad \rho_1 \quad \sigma_R \quad \alpha \\ \quad \downarrow f_L \quad (*) \quad \parallel \quad \parallel \quad \dots \quad \parallel \quad \parallel \quad (**) \quad \downarrow f_R \\ Q_\tau^\bullet: \\ \quad \delta \quad \tau_L \quad \rho_k \quad \rho_{k-1} \quad \dots \quad \rho_2 \quad \rho_1 \quad \tau_R \quad \gamma \end{array}$$

$$(2) \quad \begin{array}{c} P_\sigma^\bullet: \\ \quad \beta \quad \sigma_L \quad \rho_{-1} \quad \rho_{-2} \quad \rho_{-3} \quad \dots \\ \quad \downarrow f_L \quad (*) \quad \parallel \quad \parallel \quad \parallel \\ P_\tau^\bullet: \\ \quad \delta \quad \tau_L \quad \rho_{-1} \quad \rho_{-2} \quad \rho_{-3} \quad \dots \end{array}$$

where we require $(*)$ and $(**)$ to commute. This data defines a map of complexes $f^\bullet: Q_\sigma^\bullet \rightarrow Q_\tau^\bullet$ called a *graph map*. The isomorphisms indicated in diagrams (1) and (2) together with f_L and f_R , should they exist, are the *components* of f^\bullet .

Below we explicitly write down the *right endpoint conditions* making $(**)$ commute, where we have separated out the situation in (2) as a kind of ‘infinite right endpoint condition’:

$$(RG1) \quad \begin{array}{ccc} \bullet \xrightarrow{\sigma_R} \bullet & \xrightarrow{\quad} & \bullet \xrightarrow{\quad} \alpha \\ \parallel & \downarrow f_R & \parallel \\ \bullet \xrightarrow{\tau_R} \bullet & \xrightarrow{\tau_R = \sigma_R f_R} & \bullet \xrightarrow{\quad} \gamma \end{array} \quad (RG2) \quad \begin{array}{ccc} \bullet \xrightarrow{\sigma_R = \bar{\omega} \bar{f}_R} \bullet & \xrightarrow{\quad} & \bullet \xrightarrow{\quad} \alpha \\ \parallel & \downarrow f_R & \parallel \\ \bullet \xrightarrow{\tau_R = \bar{\omega}} \bullet & \xrightarrow{\quad} & \bullet \xrightarrow{\quad} \gamma \end{array}$$

$$(RG3) \quad \begin{array}{ccc} \bullet \xrightarrow{\sigma_R} \bullet & \xrightarrow{\quad} & \bullet \xrightarrow{\quad} \alpha \\ \parallel & & \parallel \\ \bullet \xleftarrow{\tau_R} \bullet & \xleftarrow{\quad} & \bullet \xleftarrow{\quad} \gamma \end{array} \quad (RG\infty) \quad \begin{array}{c} \bullet \xrightarrow{\sigma_L} \bullet \xrightarrow{\rho_{-1}} \bullet \xrightarrow{\rho_{-2}} \bullet \xrightarrow{\rho_{-3}} \bullet \cdots \\ \downarrow f_L \quad (*) \quad \parallel \quad \parallel \quad \parallel \\ \bullet \xrightarrow{\tau_L} \bullet \xrightarrow{\rho_{-1}} \bullet \xrightarrow{\rho_{-2}} \bullet \xrightarrow{\rho_{-3}} \bullet \cdots \end{array}$$

The *left endpoint conditions* making $(*)$ commute are defined dually.

conditions,

$$\begin{array}{ccc}
\text{(RQ1)} & \begin{array}{c} \bullet \xrightarrow{\sigma_R = \tau_R \tau'_R} \bullet \text{---} \alpha \\ \parallel \\ \bullet \xrightarrow{\tau_R} \bullet \text{---} \gamma \end{array} & \text{(RQ2)} & \begin{array}{c} \bullet \xleftarrow{\sigma_R = \bar{\omega}} \bullet \text{---} \alpha \\ \parallel \\ \bullet \xleftarrow{\tau_R = \bar{\omega} \omega'} \bullet \text{---} \gamma \end{array} & \text{(RQ3)} & \begin{array}{c} \bullet \xleftarrow{\sigma_R} \bullet \text{---} \alpha \\ \parallel \\ \bullet \xrightarrow{\tau_R} \bullet \text{---} \gamma \end{array}
\end{array}$$

where ω and ω' are direct homotopy letters, and all primed homotopy (sub)letters are nontrivial. In (RQ3) we also permit σ_R or τ_R (but not both) to be zero. The *quasi-graph map left endpoint conditions* are defined dually.

A quasi-graph map $\varphi: Q_\sigma^\bullet \rightsquigarrow \Sigma^{-1}Q_\tau^\bullet$ determines a family of homotopy equivalent single and/or double maps. All nonzero single or double maps that are not singleton occur in this way. We refer to [4, Prop. 4.8] for explicit details on the single or double maps lying in a homotopy class determined by a given quasi-graph map.

Finally, we remark ‘quasi-graph maps’ satisfying the graph map endpoint conditions give rise to a family of null-homotopic single and/or double maps. As such quasi-graph maps never arise from diagrams of the form (2) because such diagrams always satisfy a graph map endpoint conditions, namely (RG ∞) or its dual (LG ∞).

2. MAPPING CONES OF GRAPH MAPS

In this section we compute the mapping cones of graph maps. We first start by describing the unfolded diagram of the mapping cone of a graph map.

2.1. The unfolded diagram of the mapping cone of a graph map. Let $f^\bullet: Q_\sigma^\bullet \rightarrow Q_\tau^\bullet$ be a graph map. The unfolded diagram of $M_{f^\bullet}^\bullet$ has one of the following forms.

$$\begin{array}{cccccccccccc}
\text{---} \beta & \xrightarrow{-\sigma_L} & \bullet & \xrightarrow{-\rho_k} & \bullet & \xrightarrow{-\rho_{k-1}} & \dots & \xrightarrow{-\rho_2} & \bullet & \xrightarrow{-\rho_1} & \bullet & \xrightarrow{-\sigma_R} & \bullet & \text{---} \alpha \\
& \searrow f_L & \parallel & \searrow f_R & & \\
\text{---} \delta & \xrightarrow{\tau_L} & \bullet & \xrightarrow{\rho_k} & \bullet & \xrightarrow{\rho_{k-1}} & \dots & \xrightarrow{\rho_2} & \bullet & \xrightarrow{\rho_1} & \bullet & \xrightarrow{\tau_R} & \bullet & \text{---} \gamma \\
& & \parallel & & & \\
& & \text{---} \rho_3 & \xrightarrow{-\rho_2} & \bullet & \xrightarrow{-\rho_1} & \bullet & \xrightarrow{-\sigma_R} & \bullet & \xrightarrow{-\alpha} & \bullet & \searrow f_R & & \\
& & \parallel & & & \\
& & \text{---} \rho_3 & \xrightarrow{\rho_3} & \bullet & \xrightarrow{\rho_2} & \bullet & \xrightarrow{\rho_1} & \bullet & \xrightarrow{\tau_R} & \bullet & \text{---} \gamma & &
\end{array}$$

2.2. Mapping cones of graph maps between string complexes. Let σ and τ be homotopy strings and $f^\bullet: P_\sigma^\bullet \rightarrow P_\tau^\bullet$ be a graph map between the corresponding string complexes. In this case the generic situation is for $M_{f^\bullet}^\bullet$ to have two indecomposable summands, which are again string complexes and whose strings can be read off from σ , τ and f^\bullet in a natural way. We start with a technical definition which ensures that the unfolded diagram of the graph map is properly oriented for the mapping cone calculus in the case that f^\bullet is a graph map supported in precisely one degree.

Definition 2.1. Let σ and τ be homotopy strings and suppose $f^\bullet: P_\sigma^\bullet \rightarrow P_\tau^\bullet$ is a graph map concentrated in precisely one degree, i.e. the maximal common homotopy substring ρ is trivial and satisfies the graph map endpoint conditions (RG3) and (LG3):

$$\begin{array}{ccc}
\text{---} \beta & \xleftarrow{\sigma_L} & \bullet & \xrightarrow{\sigma_R} & \bullet & \text{---} \alpha \\
& & \parallel & & & \\
\text{---} \delta & \xrightarrow{\tau_L} & \bullet & \xleftarrow{\tau_R} & \bullet & \text{---} \gamma
\end{array}$$

We say that the homotopy strings σ and τ are *compatibly oriented* if $\sigma_L \tau_L \neq 0$ and $\sigma_R \tau_R \neq 0$.

We observe that if a graph map is supported in more than one degree, either by having precisely one isomorphism and satisfying endpoint conditions so that one of f_L or f_R is nonempty, or by having isomorphisms in at least two cohomological degrees (making ρ of length at least 1), then the homotopy strings and graph map between them is forced to be ‘compatibly oriented’.

Note that in the following if a homotopy string $\sigma = \emptyset$ then the corresponding string complex $P_\sigma^\bullet \cong 0^\bullet$.

Theorem 2.2. *Let σ and τ be (possibly infinite) homotopy strings and suppose $f^\bullet: P_\sigma^\bullet \rightarrow P_\tau^\bullet$ is a graph map. Suppose that σ and τ are compatibly oriented. Then we have the following cases.*

(1) *If $\sigma = \beta\sigma_L\rho\sigma_R\alpha$ and $\tau = \delta\tau_L\rho\tau_R\gamma$ then the mapping cone $M_{f^\bullet}^\bullet$ is isomorphic in $\mathbf{K}^{b,-}(\text{proj}(\Lambda))$ to the direct sum of $P_{c_1}^\bullet \oplus P_{c_2}^\bullet$, where*

$$c_1 = \begin{cases} \bar{\gamma}\bar{\tau}_R\sigma_R\alpha & \text{if } \sigma_R \neq \emptyset \text{ and } \tau_R \neq \emptyset; \\ \bar{\gamma} & \text{if } \sigma_R = \emptyset; \\ \alpha & \text{if } \tau_R = \emptyset; \\ \emptyset & \text{if } \sigma_R = \emptyset \text{ and } \tau_R = \emptyset, \end{cases} \quad \text{and } c_2 = \begin{cases} \beta\sigma_L\bar{\tau}_L\bar{\delta} & \text{if } \sigma_L \neq \emptyset \text{ and } \tau_L \neq \emptyset; \\ \bar{\delta} & \text{if } \sigma_L = \emptyset; \\ \beta & \text{if } \tau_L = \emptyset; \\ \emptyset & \text{if } \sigma_L = \emptyset \text{ and } \tau_L = \emptyset, \end{cases}$$

(2) *If $\sigma = \rho\sigma_R\alpha$ and $\tau = \rho\tau_R\gamma$ then the mapping cone $M_{f^\bullet}^\bullet$ is isomorphic in $\mathbf{K}^{b,-}(\text{proj}(\Lambda))$ to P_c^\bullet , where*

$$c = \begin{cases} \bar{\gamma}\bar{\tau}_R\sigma_R\alpha & \text{if } \sigma_R \neq \emptyset \text{ and } \tau_R \neq \emptyset; \\ \bar{\gamma} & \text{if } \sigma_R = \emptyset; \\ \alpha & \text{if } \tau_R = \emptyset; \\ \emptyset & \text{if } \sigma_R = \emptyset \text{ and } \tau_R = \emptyset. \end{cases}$$

Remark 2.3. If in Theorem 2.2(1) the graph map right endpoint conditions (RG1) or (RG2) occur then in either case $\bar{\gamma}\bar{\tau}_R\sigma_R\alpha = \bar{\gamma}\bar{f}_R\alpha$. Similarly, if the left endpoint conditions (LG1) or (LG2) occur then $\beta\sigma_L\bar{\tau}_L\bar{\delta} = \beta f_L\bar{\delta}$.

In order to prove Theorem 2.2 we first need the following well-known result. For the convenience of the reader we provide a short proof.

Lemma 2.4. *Let P^\bullet be a complex of the form*

$$P^\bullet: \quad \dots \longrightarrow P^{n-1} \xrightarrow{[a_1 \ a_2]} P^n \oplus Q \xrightarrow{\begin{bmatrix} b_1 & b_2 \\ b_3 & 1 \end{bmatrix}} P^{n+1} \oplus Q \xrightarrow{\begin{bmatrix} c_1 \\ c_2 \end{bmatrix}} P^{n+2} \longrightarrow \dots$$

Then $P^\bullet \cong (P')^\bullet \oplus Q^\bullet$, where $(P')^\bullet$ and Q^\bullet are the following complexes,

$$(P')^\bullet: \quad \dots \longrightarrow P^{n-1} \xrightarrow{a_1} P^n \xrightarrow{b_1 - b_2 b_3} P^{n+1} \xrightarrow{c_1} P^{n+2} \longrightarrow \dots;$$

$$Q^\bullet: \quad \dots \longrightarrow 0 \longrightarrow Q \xrightarrow{1} Q \longrightarrow 0 \longrightarrow \dots,$$


where $(P')^i = P^i$ and $Q^i = 0$ whenever $i \notin \{n, n+1\}$. In particular, in the homotopy category $P^\bullet \cong (P')^\bullet$.

Proof. The required isomorphism (and its inverse) are recorded in the following commutative diagram of complexes.


$$\begin{array}{ccccccc} P^\bullet: & & P^{n-1} & \xrightarrow{[a_1 \ a_2]} & P^n \oplus Q & \xrightarrow{\begin{bmatrix} b_1 & b_2 \\ b_3 & 1 \end{bmatrix}} & P^{n+1} \oplus Q & \xrightarrow{\begin{bmatrix} c_1 \\ c_2 \end{bmatrix}} & P^{n+2} \\ \sim \downarrow & & \parallel & & \begin{bmatrix} 1 & b_2 \\ 0 & 1 \end{bmatrix} \downarrow & & \downarrow \begin{bmatrix} 1 & 0 \\ -b_3 & 1 \end{bmatrix} & & \parallel \\ (P')^\bullet \oplus Q^\bullet: & & P^{n-1} & \xrightarrow{[a_1 \ 0]} & P^n \oplus Q & \xrightarrow{\begin{bmatrix} b_1 - b_2 b_3 & 0 \\ 0 & 1 \end{bmatrix}} & P^{n+1} \oplus Q & \xrightarrow{\begin{bmatrix} c_1 \\ 0 \end{bmatrix}} & P^{n+2} \\ \sim \downarrow & & \parallel & & \begin{bmatrix} 1 & -b_2 \\ 0 & 1 \end{bmatrix} \downarrow & & \downarrow \begin{bmatrix} 1 & 0 \\ b_3 & 1 \end{bmatrix} & & \parallel \\ P^\bullet: & & P^{n-1} & \xrightarrow{[a_1 \ a_2]} & P^n \oplus Q & \xrightarrow{\begin{bmatrix} b_1 & b_2 \\ b_3 & 1 \end{bmatrix}} & P^{n+1} \oplus Q & \xrightarrow{\begin{bmatrix} c_1 \\ c_2 \end{bmatrix}} & P^{n+2} \end{array}$$

Note that the commutativity of each square follows from observing that $a_2 = -a_1 b_2$ and $c_2 = -b_3 c_1$, which follows from the fact that P^\bullet is a complex so that the differential squares to zero. The final statement follows by observing that Q^\bullet is a contractible complex. \square

Remark 2.5 (Unfolded diagram interpretation of Lemma 2.4). We can reformulate the situation of Lemma 2.4 in terms of unfolded diagrams as follows. The initial complex P^\bullet has the following ‘unfolded diagram’:




The complex $(P')^\bullet$, to which P^\bullet is homotopic, has the following ‘unfolded diagram’:



Let σ and τ be (possibly infinite) homotopy strings and suppose $f^\bullet: P_\sigma^\bullet \rightarrow P_\tau^\bullet$ is a graph map. Since f^\bullet is a graph map, at least one nonzero component of f^\bullet is an isomorphism. The next lemma summarises a straightforward analysis of the form of an unfolded diagram of f^\bullet at one such component.

Lemma 2.6. *Let σ and τ be (possibly infinite) homotopy strings and suppose $f^\bullet: P_\sigma^\bullet \rightarrow P_\tau^\bullet$ is a graph map. At any given component that is an isomorphism, the unfolded diagram of f^\bullet locally has the one of the following six forms, up to inversion of σ , τ or both:*




Here σ_L , σ_R , τ_L and τ_R need not be located at the endpoints of overlaps and may be empty paths. The homotopy letters f_L , f_R may be paths, trivial paths or empty paths.

Recall the form of the unfolded diagram of $M_{f^\bullet}^\bullet$ from Section 2.1. The following corollary applies the unfolded diagram interpretation of Lemma 2.4 to each of the local situations occurring in Lemma 2.6. The resulting unfolded diagram gives rise to a complex that is homotopic to $M_{f^\bullet}^\bullet$.

Corollary 2.7. *Let σ and τ be (possibly infinite) homotopy strings and suppose $f^\bullet: P_\sigma^\bullet \rightarrow P_\tau^\bullet$ is a graph map. For each of the cases in Lemma 2.6 above, the following (compatibly oriented) unfolded diagrams correspond to complexes that are homotopic to $M_{f^\bullet}^\bullet$, where we use the notation*

of Lemma 2.6.



Here the dashed arrows indicate components of the map $b_1 - b_2 b_3$ in Lemma 2.4. Note that in each case $b_1 = 0$.


Remark 2.8. Note that in cases (iii) and (iv) of Corollary 2.7 the components b_2 and b_3 are also zero, hence why no dashed arrows appear in those diagrams.

We are now ready for the proof of Theorem 2.2.

Proof of Theorem 2.2. For convenience we distinguish between three cases: $\text{len}(\rho) = 1$, $\text{len}(\rho) = k > 1$ and $\text{len}(\rho) = \infty$. The first two cases correspond to statement (1) of the theorem.

Lemma 2.6 shows all possible forms, up to inversion and inclusion of empty homotopy letters, in the case $\text{len}(\rho) = 1$. Observe that the right and left graph map endpoint conditions together with compatible orientation in case (v) (see Definition 2.1) force, in each case, $\tau_L \sigma_R = 0$ and $\sigma_L \bar{\tau}_L = 0$. In particular, the diagrams (i) – (vi) in Corollary 2.7 give precisely the unfolded diagrams of the string complexes whose homotopy strings are c_1 and c_2 . It follows that $M_{f_\bullet}^\bullet \simeq P_{c_1}^\bullet \oplus P_{c_2}^\bullet$ in this case.


Now suppose $\text{len}(\rho) = k > 1$. Firstly consider the case ρ_1 is direct. Two applications of Corollary 2.7 show that the mapping cone $M_{f_\bullet}^\bullet$ has an unfolded diagram of the following form. On the left hand side we are in one of cases (i), (iii), (iv) or (vi) of Corollary 2.7, and on the right hand side we are in cases (i), (ii) or (iii).



Here, $\mu_1 = f_L$ is a downward pointing direct homotopy letter whenever σ_L and τ_L have the same orientation (cases (i), (iii) and (vi)) and $\mu_1 = \sigma_L \bar{\tau}_L$ is an upward pointing inverse homotopy letter whenever σ_L and τ_L have opposite orientation (case (iv)). Note that, in this case $\sigma_L \bar{\tau}_L \neq 0$. In cases (iii) and (vi) μ_2 does not exist for degree reasons and in cases (i) and (iv), $\mu_2 = \tau_L \rho_k$, which as above is zero. Now we turn to the right hand side of the diagram. In the case that σ_R is direct, whence we fall into (i) or (ii), $\nu_1 = \rho_1 \sigma_R = 0$. In the case that σ_R is inverse, putting us in case (iii), then ν_1 does not exist for degree reasons. Finally, if σ_R and τ_R have the same orientation (cases (i) and (iii)), we have $\nu_2 = f_R$ is a downward pointing direct homotopy letter. If σ_R and τ_R have opposite orientations (case (ii)) then $\nu_2 = \bar{\sigma}_R \tau_R$ is an upward pointing inverse homotopy letter, which, as before, is nonzero.

Now we consider the case that ρ_1 is inverse. We have already dealt with the left hand side of the diagram above, so we only consider the right hand side of the unfolded diagram. Here we


fall into case (vi) or the duals to cases (i) and (iv).



Here, if both σ_R and τ_R are direct (putting us in case (vi)), ν_1 does not exist for degree reasons and $\nu_2 = f_R$ is a downward pointing direct homotopy letter. If τ_R is inverse, we fall into the dual of case (i) or case (iv) and $\nu_1 = \rho_1 \tau_R = 0$. If both σ_R and τ_R are inverse (putting us in the dual of case (i)), then $\nu_2 = f_R$ is a downward pointing direct homotopy letter. If σ_R is direct and τ_R is inverse (dual to case (iv)), then $\nu_2 = \overline{\sigma_R} \tau_R$ is an upwards pointing inverse homotopy letter.

It follows that $M_{f_\bullet}^\bullet$ is homotopic to the direct sum of the string complexes $P_{c_1}^\bullet$ and $P_{c_2}^\bullet$ and the mapping cone of $\text{id}^\bullet: P_{\rho'}^\bullet \rightarrow P_{\rho'}^\bullet$ where ρ' is the homotopy string $\rho' = \rho_{k-1} \cdots \rho_2$. In particular, $M_{\text{id}^\bullet}^\bullet$ is contractible. Note that in the case $n = 1$, $\rho' = \emptyset$ is the empty homotopy string and therefore $P_{\rho'}^\bullet = 0^\bullet$ and when $n = 2$, $\rho' = 1_{e(\rho_1)}$ is a trivial homotopy string. It follows that $M_{f_\bullet}^\bullet \simeq P_{c_1}^\bullet \oplus P_{c_2}^\bullet$.

Finally, suppose $\text{len}(\rho) = \infty$, which puts us into statement (2) of the theorem, or its dual. Let's consider the setup in statement (2), which gives us the following unfolded diagram after one application of Corollary 2.7, where μ_1 and μ_2 are defined as above.



It follows that $M_{f_\bullet}^\bullet$ is homotopic to the direct sum of the string complex P_c^\bullet and the mapping cone of $\text{id}^\bullet: P_{\rho'}^\bullet \rightarrow P_{\rho'}^\bullet$, where ρ' is the infinite homotopy string $\rho' = \rho_{-2} \rho_{-3} \rho_{-4} \cdots$. Hence, $M_{f_\bullet}^\bullet \simeq P_c^\bullet$. \square

2.3. Mapping cones of graph maps involving a band complex. Suppose σ and τ are homotopy strings or bands, with at least one being a homotopy band. In this section we consider the mapping cones of graph maps $f^\bullet: Q_\sigma^\bullet \rightarrow Q_\tau^\bullet$. The difference with Theorem 2.2 is that now $M_{f_\bullet}^\bullet$ has only one indecomposable summand. Moreover, this summand is a band complex precisely when both σ and τ are homotopy bands, and is a string complex otherwise.

We start with the situation that both σ and τ are homotopy bands. We impose the convention that the scalars λ and μ are placed on direct arrows of σ and τ , respectively.

Proposition 2.9. *Let $\sigma = \beta \sigma_L \rho \sigma_R \alpha$ and $\tau = \delta \tau_L \rho \tau_R \gamma$ be homotopy bands. Suppose $f^\bullet: B_{\sigma, \lambda}^\bullet \rightarrow B_{\tau, \mu}^\bullet$ is a compatibly oriented graph map which is not the identity. Then*

$$M_{f_\bullet}^\bullet \cong \begin{cases} B_{c, \lambda \mu^{-1}}^\bullet & \text{if } \text{len}(\rho) \text{ is even;} \\ B_{c, -\lambda \mu^{-1}}^\bullet & \text{if } \text{len}(\rho) \text{ is odd,} \end{cases}$$

where $c = \beta \sigma_L \bar{\tau}_L \bar{\delta} \bar{\gamma} \bar{\tau}_R \sigma_R \alpha$ and where the scalar $\lambda \mu^{-1}$ is placed on a direct homotopy letter of α , β , $\bar{\gamma}$ or $\bar{\delta}$.

Note that here we make use Remark 2.3 when describing the homotopy band c .

Proof. We first check that $c = \beta \sigma_L \bar{\tau}_L \bar{\delta} \bar{\gamma} \bar{\tau}_R \sigma_R \alpha$ is a homotopy band. To do this, we introduce some notation. Let

$$\partial(\sigma) = \# \text{ direct homotopy letters of } \sigma, \text{ and } \iota(\sigma) = \# \text{ inverse homotopy letters of } \sigma.$$

We must show that $\partial(c) = \iota(c)$. Suppose, as usual, that $\sigma = \sigma_m \cdots \sigma_1$ and $\tau = \tau_n \cdots \tau_1$. Then we have $m = 2m'$ and $n = 2n'$, where $m' = \partial(\sigma) = \iota(\sigma)$ and $n' = \partial(\tau) = \iota(\tau)$.

We start by dealing with the case that $f^\bullet: B_{\sigma,\lambda}^\bullet \rightarrow B_{\tau,\mu}^\bullet$ does not satisfy the graph map endpoint conditions (RG3) or (LG3); note that f^\bullet cannot satisfy (RG ∞) or (LG ∞) because σ and τ are homotopy bands. In this case $c = \beta f_L \bar{\delta} \bar{\gamma} \bar{f}_R \alpha$.

Suppose $\partial(\sigma_L \rho \sigma_R) = l = \partial(\tau_L \rho \tau_R)$ and $\iota(\sigma_L \rho \sigma_R) = l' = \iota(\tau_L \rho \tau_R)$. Then $\partial(\alpha) + \partial(\beta) = m' - l$ and $\iota(\alpha) + \iota(\beta) = m' - l'$. Similarly, $\partial(\gamma) + \partial(\delta) = n' - l$ and $\iota(\gamma) + \iota(\delta) = n' - l'$. It follows that

$$\begin{aligned} \partial(c) &= \partial(\alpha) + \partial(\beta) + \iota(\gamma) + \iota(\delta) + 1 = m' - l + n' - l' + 1, \text{ and,} \\ \iota(c) &= \iota(\alpha) + \iota(\beta) + \partial(\gamma) + \partial(\delta) + 1 = m' - l' + n' - l + 1. \end{aligned}$$

The remaining cases are:

- f^\bullet satisfies (RG3) but not (LG3) so that $c = \beta f_L \bar{\delta} \bar{\gamma} (\bar{\tau}_R \sigma_R) \alpha$, where since τ_R is an inverse homotopy letter then $(\bar{\tau}_R \sigma_R)$ is a direct homotopy letter;
- f^\bullet satisfies (LG3) but not (RG3) so that $c = \beta (\sigma_L \bar{\tau}_L) \bar{\delta} \bar{\gamma} \bar{f}_R \alpha$, where since σ_L is an inverse homotopy letter then $(\sigma_L \bar{\tau}_L)$ is also an inverse homotopy letter;
- f^\bullet satisfies (RG3) and (LG3) so that $c = \beta (\sigma_L \bar{\tau}_L) \bar{\delta} \bar{\gamma} (\bar{\tau}_R \sigma_R) \alpha$, with the same observations as above.

In each case, we obtain $\partial(c) = m' + n' - l - l' + 1 = \iota(c)$. Hence c is a homotopy band, as claimed.

The remainder of the proof is the same as the proof of Theorem 2.2 except that we have to take into account the scalar $\pm \lambda \mu^{-1}$. First let us deal with the sign. Consider a homotopy band $\sigma = \sigma_m \cdots \sigma_1$ and place a minus sign on any two homotopy letters, say σ_j and σ_i with $m \geq j > i \geq 1$. Then the following diagram induces an isomorphism of band complexes.

$$\begin{array}{cccccccccccccccc} \cdots & \xrightarrow{\sigma_1} & \bullet & \xrightarrow{\sigma_m} & \bullet & \cdots & \bullet & \xrightarrow{-\sigma_j} & \bullet & \xrightarrow{\sigma_{j-1}} & \bullet & \cdots & \bullet & \xrightarrow{\sigma_{i+1}} & \bullet & \xrightarrow{-\sigma_i} & \bullet & \cdots & \bullet & \xrightarrow{\sigma_1} & \bullet & \xrightarrow{\sigma_m} & \cdots \\ & & \parallel & & \parallel & & \parallel & \downarrow -1 & \parallel & \downarrow -1 & \parallel & & \parallel & \downarrow -1 & \parallel & \downarrow -1 & \parallel & & \parallel & & \parallel & & \parallel & \\ \cdots & \xrightarrow{\sigma_1} & \bullet & \xrightarrow{\sigma_m} & \bullet & \cdots & \bullet & \xrightarrow{\sigma_j} & \bullet & \xrightarrow{\sigma_{j-1}} & \bullet & \cdots & \bullet & \xrightarrow{\sigma_{i+1}} & \bullet & \xrightarrow{\sigma_i} & \bullet & \cdots & \bullet & \xrightarrow{\sigma_1} & \bullet & \xrightarrow{\sigma_m} & \cdots \end{array}$$

Now the unfolded diagram of the complex homotopic to M_f^\bullet , given by the arguments of Theorem 2.2 using Lemma 2.4 and Corollary 2.7 corresponds precisely to the homotopy band c . In this unfolded diagram, the homotopy letters carrying a minus sign are precisely those of the homotopy substrings α and β . The number of homotopy letters in α and β is equal to $m - k + 2$, which is even if and only if $k = \text{len}(\rho)$ is even. In the case that $\text{len}(\rho)$ is even, all the minus signs can be eliminated by an isomorphism as above. In the case that $\text{len}(\rho)$ is odd, all but one minus sign can be removed by such an isomorphism. This deals with the signs.

To deal with the scalar $\lambda \mu^{-1}$ note that a scalar can be moved along to an adjacent homotopy letter as indicated in the following diagram.

$$\begin{array}{ccc} \cdots \bullet \xrightarrow{\lambda \sigma_i} \bullet \xrightarrow{\sigma_{i-1}} \bullet \cdots & & \cdots \bullet \xrightarrow{\lambda \sigma_i} \bullet \xleftarrow{\sigma_{i-1}} \bullet \cdots \\ \parallel & \downarrow \lambda^{-1} & \parallel \\ \cdots \bullet \xrightarrow{\sigma_i} \bullet \xrightarrow{\lambda \sigma_{i-1}} \bullet \cdots & & \cdots \bullet \xrightarrow{\sigma_i} \bullet \xleftarrow{\lambda^{-1} \sigma_{i-1}} \bullet \cdots \end{array}$$

This completes the proof. □

It is useful to illustrate the proof of Proposition 2.9 in an example.

Example 2.10. Let Λ be the algebra given by the following quiver with relations.

$$\begin{array}{ccccc} 3 & \xrightarrow{c} & 4 & \xrightarrow{f} & 6 \\ \uparrow b & & \uparrow d & & \uparrow g \\ 2 & \xrightarrow{e} & 5 & \xrightarrow{h} & 7 \\ \uparrow a & & & & \uparrow i \\ 1 & \xrightarrow{j} & & & 8 \end{array}$$

Consider the following homotopy bands: $\sigma = \bar{e}\bar{d}cb$, $\tau = \bar{j}\bar{i}\bar{g}fcb$. We consider graph maps $f^\bullet: B_{\sigma,\lambda}^\bullet \rightarrow B_{\tau,\mu}^\bullet$ defined by the following unfolded diagram.

$$\begin{array}{c}
B_{\sigma,\lambda}^\bullet: \\
f^\bullet \downarrow \\
B_{\tau,\mu}^\bullet:
\end{array}
\begin{array}{ccccccccccc}
\bullet & \xleftarrow{\lambda^{-1}\bar{e}} & \bullet & \xleftarrow{\bar{d}} & \bullet & \xrightarrow{c} & \bullet & \xrightarrow{b} & \bullet & \xleftarrow{\lambda^{-1}\bar{e}} & \bullet \\
& & & & \parallel & & \parallel & & \downarrow a & & \\
\bullet & \xleftarrow{\bar{j}} & \bullet & \xleftarrow{\bar{i}} & \bullet & \xleftarrow{\mu^{-1}\bar{g}} & \bullet & \xrightarrow{f} & \bullet & \xrightarrow{c} & \bullet & \xrightarrow{ba} & \bullet & \xleftarrow{\bar{j}} & \bullet
\end{array}$$

Let $c_\tau = \bar{e}\bar{f}\bar{d}gij\bar{a}$. Let us work through the calculation of $M_{f^\bullet}^\bullet$, using the techniques of Lemma 2.4 and Corollary 2.7.

$$\begin{array}{ccccccccccc}
\bullet & \xleftarrow{-\lambda^{-1}\bar{e}} & \bullet & \xleftarrow{-\bar{d}} & \bullet & \xrightarrow{-c} & \bullet & \xrightarrow{b} & \bullet & \xleftarrow{-\lambda^{-1}\bar{e}} & \bullet \\
& & & & \parallel & & \parallel & & \searrow a & & \\
\bullet & \xleftarrow{\bar{j}} & \bullet & \xleftarrow{\bar{i}} & \bullet & \xleftarrow{\mu^{-1}\bar{g}} & \bullet & \xrightarrow{f} & \bullet & \xrightarrow{c} & \bullet & \xrightarrow{ba} & \bullet & \xleftarrow{\bar{j}} & \bullet
\end{array}$$

Removing the isomorphisms using Corollary 2.7 we get the following unfolded diagram of a homotopic complex, where \circ denotes the removed endpoints of the isomorphisms.

$$\begin{array}{ccccccc}
\bullet & \xleftarrow{-\lambda^{-1}\bar{e}} & \bullet & \xleftarrow{\bar{d}\bar{f}} & \circ & \circ & \bullet & \xleftarrow{-\lambda^{-1}\bar{e}} & \bullet \\
& & & \searrow & & & \searrow a & & \\
\bullet & \xleftarrow{\bar{j}} & \bullet & \xleftarrow{\bar{i}} & \bullet & \xleftarrow{\mu^{-1}\bar{g}} & \bullet & \xleftarrow{\bar{j}} & \bullet
\end{array}$$

Straightening this out, we get the following unfolded diagram,

$$\cdots \xleftarrow{-\lambda^{-1}\bar{e}} \bullet \xleftarrow{\bar{d}\bar{f}} \bullet \xrightarrow{\mu^{-1}\bar{g}} \bullet \xrightarrow{i} \bullet \xrightarrow{j} \bullet \xleftarrow{\bar{a}} \bullet \xleftarrow{-\lambda^{-1}\bar{e}} \bullet \xleftarrow{\bar{d}\bar{f}} \cdots,$$

which gives rise to a band complex isomorphic to a band complex with the unfolded diagram,

$$\cdots \xleftarrow{\bar{e}} \bullet \xleftarrow{\bar{d}\bar{f}} \bullet \xrightarrow{-\lambda\mu^{-1}\bar{g}} \bullet \xrightarrow{i} \bullet \xrightarrow{j} \bullet \xleftarrow{\bar{a}} \bullet \xleftarrow{\bar{e}} \bullet \xleftarrow{\bar{d}\bar{f}} \cdots.$$

We now state the corresponding statements when only one of σ or τ is a homotopy band. In these cases $M_{f^\bullet}^\bullet$ is a string complex and there are no signs or scalars to take care of.

Proposition 2.11. *Let $\sigma = \beta\sigma_L\rho\sigma_R\alpha$ be a homotopy band and $\tau = \delta\tau_L\rho\tau_R\gamma$ be a homotopy string. Suppose $f^\bullet: B_{\sigma,\lambda}^\bullet \rightarrow P_\tau^\bullet$ is a compatibly oriented graph map. Then $M_{f^\bullet}^\bullet$ is isomorphic to the string complex P_c^\bullet , where $c = \delta\tau_L\bar{\sigma}_L\bar{\beta}\bar{\alpha}\bar{\sigma}_R\tau_R\gamma$.*

Proposition 2.12. *Let $\sigma = \beta\sigma_L\rho\sigma_R\alpha$ be a homotopy string and $\tau = \delta\tau_L\rho\tau_R\gamma$ be a homotopy band. Suppose $f^\bullet: P_\sigma^\bullet \rightarrow B_{\tau,\mu}^\bullet$ is a compatibly oriented graph map. Then $M_{f^\bullet}^\bullet$ is isomorphic to the string complex P_c^\bullet , where $c = \beta\sigma_L\bar{\tau}_L\bar{\delta}\bar{\gamma}\bar{\tau}_R\sigma_R\alpha$.*

3. MAPPING CONES OF SINGLE MAPS


In this section, we describe the mapping cone calculus for single maps. As in Section 2, we first describe the unfolded diagram of the mapping cone and then describe the mapping cone calculus in the case of a single map between string complexes. We then deal with the cases in which (at least) one of the indecomposable complexes is a band complex.

3.1. The unfolded diagram of the mapping cone of a single map. Let $f^\bullet: Q_\sigma^\bullet \rightarrow Q_\tau^\bullet$ be a single map with single component f . We illustrate the unfolded diagram of $M_{f^\bullet}^\bullet$ in the generic situation.

$$\begin{array}{ccccccc}
\text{---} \bullet & \xrightarrow{-\sigma_L} & \bullet & \xrightarrow{-\sigma_R} & \bullet & \text{---} \\
& & \searrow f & & & \\
\text{---} \bullet & \xrightarrow{\tau_L} & \bullet & \xleftarrow{\tau_R} & \bullet & \text{---} \\
& & \delta & & \gamma &
\end{array}$$

3.2. Mapping cones of single maps between string complexes. We start with a technical definition analogous to Definition 2.1.

Definition 3.1. Let σ and τ be homotopy strings and suppose $f^\bullet: P_\sigma^\bullet \rightarrow P_\tau^\bullet$ is a (not necessarily singleton) single map sitting in the following orientation.



We shall say that the homotopy strings σ and τ are *compatibly oriented* (for f^\bullet) if

- (i) the homotopy letters σ_L and τ_L do not contain f as a subletter;
- (ii) if σ_L is direct then $\sigma_L f = 0$ and if τ_L is inverse $f \tau_L = 0$;
- (iii) $\sigma_R = f f_R$ and $\tau_R = f_L f$ for some (possibly trivial) paths f_R and f_L in (Q, I) .


We note that the presentation of a singleton single map given in Section 1.4.2 is already compatibly oriented. For a general single map we simply allow f_L or f_R to be trivial.

Theorem 3.2. Let $f^\bullet: P_\sigma^\bullet \rightarrow P_\tau^\bullet$ be a singleton single map with single component f . Suppose that $\sigma = \beta \sigma_L \sigma_R \alpha$ and $\tau = \delta \tau_L \tau_R \gamma$ are compatibly oriented for f . Then $M_f^\bullet \simeq P_{c_1}^\bullet \oplus P_{c_2}^\bullet$, where


$$c_1 = \begin{cases} \beta \sigma_L f \tau_L \bar{\delta} & \text{if } \sigma_L \neq \emptyset \text{ and } \tau_L \neq \emptyset; \\ \beta \sigma_L f & \text{if } \sigma_L \neq \emptyset \text{ and } \tau_L = \emptyset; \\ f \tau_L \bar{\delta} & \text{if } \sigma_L = \emptyset \text{ and } \tau_L \neq \emptyset; \\ f & \text{if } \sigma_L = \emptyset \text{ and } \tau_L = \emptyset, \end{cases} \quad \text{and} \quad c_2 = \begin{cases} \bar{\gamma} f_L f f_R \alpha & \text{if } \sigma_R \neq \emptyset \text{ and } \tau_R \neq \emptyset; \\ \alpha & \text{if } \sigma_R \neq \emptyset \text{ and } \tau_R = \emptyset; \\ \bar{\gamma} & \text{if } \sigma_R = \emptyset \text{ and } \tau_R \neq \emptyset; \\ \emptyset & \text{if } \sigma_R = \emptyset \text{ and } \tau_R = \emptyset. \end{cases}$$

Remark 3.3. Observe that the form of c_2 in the case that $\sigma_R = \emptyset$ or $\tau_R = \emptyset$ is not what one would expect from naive word combinatorics.

Proof. The strategy of the proof is to define an isomorphism $i^\bullet: P_{c_1}^\bullet \oplus P_{c_2}^\bullet \rightarrow M_f^\bullet$ directly. This isomorphism is easier to see at the level of unfolded diagrams. For word combinatorial reasons it is useful to state Theorem 3.2 using the compatible orientation of Definition 3.1. However, the unfolded diagrams of the mapping cones look ‘more like complexes’ and are thus easier to work with if the opposite orientation is taken for τ . For clarity, the diagram below indicates how the map looks using this choice of orientation.



For convenience, assume $\beta = \sigma_m \cdots \sigma_{i+2}$, $\sigma_L = \sigma_{i+1}$, $\sigma_R = \sigma_i$, $\alpha = \sigma_{i-1} \cdots \sigma_1$ and $\gamma = \tau_1 \cdots \tau_{j-1}$, $\tau_R = \tau_j$, $\tau_L = \tau_{j+1}$, $\delta = \tau_n \cdots \tau_{j+2}$. Then the unfolded diagram of the mapping cone M_f^\bullet is



We define the maps $i_1^\bullet: P_{c_1}^\bullet \rightarrow M_f^\bullet$ and $i_2^\bullet: P_{c_2}^\bullet \rightarrow M_f^\bullet$, respectively, in the figures below. In the figure, we write $x_i = e(\sigma_i)$, $x_o = s(\sigma_1)$, $y_j = e(\tau_j)$ and $y_0 = s(\tau_1)$.




FIGURE 1. Unfolded diagram for $i_1^\bullet: P_{c_1}^\bullet \rightarrow M_{f^\bullet}^\bullet$.




FIGURE 2. Unfolded diagram for $i_2^\bullet: P_{c_2}^\bullet \rightarrow M_{f^\bullet}^\bullet$.

It is straightforward to check that these induce well-defined morphisms of complexes: only the commutativity of the central part of the diagram in each case is not immediately clear. The central part of the diagram defining $i_1^\bullet: P_{c_1}^\bullet \rightarrow M_{f^\bullet}^\bullet$ is,

$$\begin{array}{ccc} P(x_i) & \xrightarrow{f} & P(y_j) \\ \begin{bmatrix} 1 & 0 \end{bmatrix} \downarrow & & \downarrow \begin{bmatrix} -f_R & 1 \end{bmatrix} \\ P(x_i) \oplus P(y_{j-1}) & \xrightarrow{\begin{bmatrix} -ff_R & f \\ 0 & f_Lf \end{bmatrix}} & P(x_{i-1}) \oplus P(y_j), \end{array}$$

which is clearly commutative. We need to check that $f_R\sigma_{i-1} = 0$ in the case that σ_{i-1} is direct. But this is clear since in that case $\sigma_i\sigma_{i-1} = 0$ and $\sigma_i = ff_R$ forcing $f_R\sigma_{i-1} = 0$. For the central part of the diagram defining $i_2^\bullet: P_{c_2}^\bullet \rightarrow M_{f^\bullet}^\bullet$ we have,

$$\begin{array}{ccc} P(y_{j-1}) & \xrightarrow{f_Lf_R} & P(x_{i-1}) \\ \begin{bmatrix} -f_L & 1 \end{bmatrix} \downarrow & & \downarrow \begin{bmatrix} 1 & 0 \end{bmatrix} \\ P(x_i) \oplus P(y_{j-1}) & \xrightarrow{\begin{bmatrix} -ff_R & f \\ 0 & f_Lf \end{bmatrix}} & P(x_{i-1}) \oplus P(y_j), \end{array}$$

which is clearly commutative. We need to check that $f_L\bar{\sigma}_{i+1} = 0$ in the case that $\sigma_{i+1} = \sigma_L$ is inverse. Again, this is clear since $\sigma_i = \sigma_R = ff_R$ together with $f_Lf \neq 0$ and gentleness imply that $f_L\bar{\sigma}_{i+1} = 0$. Putting together the two diagrams, we get the following,

$$\begin{array}{ccc} P(x_i) \oplus P(y_{j-1}) & \xrightarrow{\begin{bmatrix} f & 0 \\ 0 & f_Lf_R \end{bmatrix}} & P(y_j) \oplus P(x_{i-1}) \\ \begin{bmatrix} 1 & 0 \\ -f_L & 1 \end{bmatrix} \downarrow & & \downarrow \begin{bmatrix} -f_R & 1 \\ 1 & 0 \end{bmatrix} \\ P(x_i) \oplus P(y_{j-1}) & \xrightarrow{\begin{bmatrix} -ff_R & f \\ 0 & f_Lf \end{bmatrix}} & P(x_{i-1}) \oplus P(y_j), \end{array}$$

where the downward maps are clearly full rank matrices. In particular, $i^\bullet = \begin{bmatrix} i_1^\bullet \\ i_2^\bullet \end{bmatrix} : P_{c_1}^\bullet \oplus P_{c_2}^\bullet \rightarrow M_{f^\bullet}^\bullet$ consists of full rank matrices in each degree, and is therefore an isomorphism of complexes.

We note that in the case that $\sigma_R = \emptyset$, the right hand side of the second diagram above does not exist and $\overline{\tau_1} \cdots \overline{\tau_{j-1}} = \overline{\gamma}$ and $c_2 = \gamma$. Similarly, if $\tau_R = \emptyset$ then the left hand side of the same diagram does not exist and $\alpha = \sigma_{i-1} \cdots \sigma_1$ and $c_2 = \alpha$. If both $\sigma_R = \emptyset$ and $\tau_R = \emptyset$ then the top line of this diagram does not exist and $c_2 = \emptyset$. \square

3.3. Mapping cones of single maps involving a band complex. Suppose σ and τ are homotopy strings or bands, with at least one being a homotopy band. We now consider the mapping cones of single maps $f^\bullet : Q_\sigma^\bullet \rightarrow Q_\tau^\bullet$. As was the case in Section 2.3, $M_{f^\bullet}^\bullet$ is now indecomposable. Moreover, $M_{f^\bullet}^\bullet$ is a band complex if and only if σ and τ are homotopy bands. This is the situation with which we start.

If both σ and τ are homotopy bands, then any single map $f^\bullet : B_{\sigma,\lambda}^\bullet \rightarrow B_{\tau,\mu}^\bullet$ is necessarily of type (iv). We again impose the convention that the scalars λ and μ are placed on direct arrows of σ and τ , respectively.

Proposition 3.4. *Suppose σ and τ are homotopy bands and $f^\bullet : B_{\sigma,\lambda}^\bullet \rightarrow B_{\tau,\mu}^\bullet$ with single component f . Suppose that $\sigma = \beta\sigma_L\sigma_R\alpha$ and $\tau = \delta\tau_L\tau_R\gamma$ are compatibly oriented for f^\bullet . Then $M_{f^\bullet}^\bullet$ is isomorphic to a band complex $B_{c,-\lambda\mu^{-1}}^\bullet$, where $c = \beta\sigma_L f \overline{\tau_L} \overline{\delta} \overline{\gamma} \overline{\tau_R} \overline{f} \sigma_R \alpha$ and where the scalar $-\lambda\mu^{-1}$ is placed on a direct homotopy letter of α , β , $\overline{\gamma}$ or $\overline{\delta}$.*

Note that in the statement above $\overline{\tau_R} \overline{f} \sigma_R = f_L f \overline{f} f f_R = f_L f f_R$.

Proof. The verification that c is indeed a homotopy band is similar to that in the proof of Proposition 2.9. The construction of an isomorphism $i^\bullet : B_{c,-\lambda\mu^{-1}}^\bullet \rightarrow M_{f^\bullet}^\bullet$ proceeds exactly as in the proof of Theorem 3.2. The sign occurs by observing that one of the homotopy substrings $\sigma_m \cdots \sigma_{i+1}$ and $\sigma_{i-1} \cdots \sigma_1$ has an even number of homotopy letters and the other an odd number of homotopy letters since σ has in total an even number of homotopy letters and only σ_i has been removed. Therefore the signs on the identity morphisms with domain $P(x_m) = P(x_0)$ in the unfolded diagrams in Figures 1 and 2 are different. However, i_1^\bullet and i_2^\bullet are ‘glued together’ to form i^\bullet by identifying these two maps with opposite signs. This is only possible if a minus sign is introduced to an odd number of homotopy letters of $\beta\sigma_L$ and α to make it that these maps have the same sign. By the argument of Theorem 3.2 this odd number of minus signs can then be reduced to one, which we position on the direct homotopy letter bearing the scalar. \square

The following two propositions deal with the cases where precisely one of σ or τ is a homotopy band and the other is a homotopy string. In these cases $M_{f^\bullet}^\bullet$ is a string complex. The proofs proceed exactly as in Theorem 3.2. Note that when σ is a homotopy band we have $\sigma_L \neq \emptyset$ and $\sigma_R \neq \emptyset$.

Proposition 3.5. *Suppose σ is a homotopy band and τ is a homotopy string. Suppose $f^\bullet : B_{\sigma,\lambda}^\bullet \rightarrow P_\tau^\bullet$ is a single map with single component f . Suppose that $\sigma = \beta\sigma_L\sigma_R\alpha$ and $\tau = \delta\tau_L\tau_R\gamma$ is compatibly oriented for f^\bullet . Then $M_{f^\bullet}^\bullet \simeq P_c^\bullet$, where*

$$c = \begin{cases} \alpha\beta\sigma_L f \overline{\tau_L} \overline{\delta} & \text{if } \tau_R = \emptyset; \\ \overline{\gamma} \overline{\tau_R} \overline{f} \sigma_R \alpha \beta \sigma_L f \overline{\tau_L} \overline{\delta} & \text{if } \tau_R \neq \emptyset. \end{cases}$$

Proposition 3.6. *Suppose σ is a homotopy string and τ is a homotopy band. Suppose $f^\bullet : P_\sigma^\bullet \rightarrow B_{\tau,\mu}^\bullet$ is a single map with single component f . Suppose that $\sigma = \beta\sigma_L\sigma_R\alpha$ and $\tau = \delta\tau_L\tau_R\gamma$ is compatibly oriented for f^\bullet . Then $M_{f^\bullet}^\bullet \simeq P_c^\bullet$, where*

$$c = \begin{cases} \beta\sigma_L f \overline{\tau_L} \overline{\delta} \overline{\gamma} & \text{if } \sigma_R = \emptyset; \\ \beta\sigma_L f \overline{\tau_L} \overline{\delta} \overline{\gamma} \overline{\tau_R} \overline{f} \sigma_R \alpha & \text{if } \sigma_R \neq \emptyset. \end{cases}$$

4. MAPPING CONES OF DOUBLE MAPS

We now turn to the statement for double maps. Note that double maps are automatically ‘compatibly oriented’ and therefore we do not require such a definition in this case.

Theorem 4.1. *Let $f^\bullet: P_\sigma^\bullet \rightarrow P_\tau^\bullet$ be a double map with components (f_L, f_R) . Suppose that $\sigma = \beta\sigma_L\sigma_C\sigma_R\alpha$ and $\tau = \delta\tau_L\tau_C\tau_R\gamma$. Then $M_{f^\bullet} \simeq P_{c_1}^\bullet \oplus P_{c_2}^\bullet$, where $c_1 = \overline{\gamma\tau_R f_R \sigma_R \alpha}$ and $c_2 = \beta\sigma_L f_L \overline{\tau_L \delta}$.*

Proof. In terms of unfolded diagrams, the map f^\bullet is of the following form, where f is permitted to be a trivial path. The case when f is not trivial corresponds to a singleton double map.

$$\begin{array}{ccccccc}
 \beta & \cdot & \xrightarrow{\sigma_L} & \cdot & \xrightarrow{\sigma_C=f_L f} & \cdot & \xrightarrow{\sigma_R} & \cdot & \xrightarrow{\alpha} \\
 & & & \downarrow f_L & & \downarrow f_R & & & \\
 \delta & \cdot & \xrightarrow{\tau_L} & \cdot & \xrightarrow{\tau_C=f f_R} & \cdot & \xrightarrow{\tau_R} & \cdot & \xrightarrow{\gamma}
 \end{array}$$

For convenience, assume $\sigma_C = \sigma_i$ and $\tau_C = \tau_j$ etc. Then the unfolded diagram of the mapping cone M_{f^\bullet} is

$$\begin{array}{ccccccccccc}
 \cdot & \xrightarrow{-\sigma_m} & \cdot & \xrightarrow{-\sigma_{i+1}} & \cdot & \xrightarrow{-\sigma_i=-f_L f} & \cdot & \xrightarrow{\sigma_{i-1}} & \cdot & \xrightarrow{\sigma_1} & \cdot \\
 & & & & & \searrow f_L & & \searrow f_R & & & \\
 \cdot & \xrightarrow{\tau_n} & \cdot & \xrightarrow{\tau_{j+1}} & \cdot & \xrightarrow{\tau_j=f f_R} & \cdot & \xrightarrow{\tau_{j-1}} & \cdot & \xrightarrow{\tau_1} & \cdot
 \end{array}$$

The proof is the same as that of Theorem 3.2, therefore, we just write down the maps $i_1^\bullet: P_{c_1}^\bullet \rightarrow M_{f^\bullet}$ and $i_2^\bullet: P_{c_2}^\bullet \rightarrow M_{f^\bullet}$ at the level of unfolded diagrams in the figures below. It is then straightforward to check these define full rank matrices in each cohomological degree, so that $i^\bullet = \begin{bmatrix} i_1^\bullet \\ i_2^\bullet \end{bmatrix}: P_{c_1}^\bullet \oplus P_{c_2}^\bullet \rightarrow M_{f^\bullet}$ is an isomorphism.

$$\begin{array}{cccccccccccc}
 P(x_{i-1}) & \xrightarrow{\sigma_{i-1}} & P(x_{i-2}) & \xrightarrow{\sigma_{i-2}} & P(x_{i-3}) & \cdots & P(x_2) & \xrightarrow{\sigma_2} & P(x_1) & \xrightarrow{\sigma_1} & P(x_0) \\
 \downarrow f_R & & \downarrow f_R \\
 P(y_{j-1}) & \xrightarrow{\tau_{j-1}} & P(y_{j-2}) & \xrightarrow{\tau_{j-2}} & P(y_{j-3}) & \cdots & P(y_2) & \xrightarrow{\tau_2} & P(y_1) & \xrightarrow{\tau_1} & P(y_0) \\
 \downarrow f_L & & \downarrow f_L \\
 P(x_i) & \xrightarrow{-\sigma_i=-f_L f} & P(x_{i-1}) & \xrightarrow{-\sigma_{i-1}} & P(x_{i-2}) & \xrightarrow{-\sigma_{i-2}} & P(x_{i-3}) & \cdots & P(x_2) & \xrightarrow{-\sigma_2} & P(x_1) & \xrightarrow{-\sigma_1} & P(x_0) \\
 \downarrow f_L & & \downarrow f_L \\
 P(y_j) & \xrightarrow{f f_R} & P(y_{j-1}) & \xrightarrow{f f_R} & P(y_{j-2}) & \xrightarrow{f f_R} & P(y_{j-3}) & \cdots & P(y_2) & \xrightarrow{f f_R} & P(y_1) & \xrightarrow{f f_R} & P(y_0)
 \end{array}$$

$$\begin{array}{cccccccccccc}
 P(x_m) & \xrightarrow{\sigma_m} & P(x_{m-1}) & \cdots & P(x_{i+2}) & \xrightarrow{\sigma_{i+2}} & P(x_{i+1}) & \xrightarrow{\sigma_{i+1}} & P(x_i) & \xrightarrow{f_L} & P(y_j) \\
 \downarrow f_L & & \downarrow f_L \\
 P(y_n) & \xrightarrow{\tau_n} & P(y_{n-1}) & \cdots & P(y_{j+2}) & \xrightarrow{\tau_{j+2}} & P(y_{j+1}) & \xrightarrow{\tau_{j+1}} & P(y_j) & \xrightarrow{-f} & P(x_{i-1}) \\
 \downarrow f_L & & \downarrow f_L \\
 P(x_m) & \xrightarrow{-\sigma_m} & P(x_{m-1}) & \cdots & P(x_{i+2}) & \xrightarrow{-\sigma_{i+2}} & P(x_{i+1}) & \xrightarrow{-\sigma_{i+1}} & P(x_i) & \xrightarrow{f_L f} & P(x_{i-1}) & \xrightarrow{f_R} & P(y_{j-1}) \\
 \downarrow f_L & & \downarrow f_L \\
 P(y_n) & \xrightarrow{\tau_n} & P(y_{n-1}) & \cdots & P(y_{j+2}) & \xrightarrow{\tau_{j+2}} & P(y_{j+1}) & \xrightarrow{\tau_{j+1}} & P(y_j) & \xrightarrow{f f_R} & P(y_{j-1})
 \end{array}$$

We note that the signs on the isomorphisms $P(x_0) \rightarrow P(x_0)$ and $P(x_m) \rightarrow P(x_m)$ defined in the above diagram are necessarily different because one of $\sigma_m \cdots \sigma_{i+1}$ and $\sigma_{i-1} \cdots \sigma_1$ has an even number of homotopy letters and the other an odd number of homotopy letters. \square

We now give the analogous statements for double maps $f^\bullet: Q_\sigma^\bullet \rightarrow Q_\tau^\bullet$ where at least one of σ or τ is a homotopy band. As was the case for graph maps and single maps before, M_{f^\bullet} is indecomposable and is a band complex if and only if both σ and τ are homotopy bands. In

this case we get an extra sign on the scalar because of the different parities in the lengths of $\sigma_m \cdots \sigma_{i+1}$ and $\sigma_{i-1} \cdots \sigma_1$ noted above.

Proposition 4.2. *Let $\sigma = \beta\sigma_L\sigma_C\sigma_R\alpha$ and $\tau = \delta\tau_L\tau_C\tau_R\gamma$ be homotopy bands. Suppose $f^\bullet: B_{\sigma,\lambda}^\bullet \rightarrow B_{\tau,\mu}^\bullet$ is a double map with components (f_L, f_R) . Then $M_{f^\bullet} \simeq B_{c, -\lambda\mu^{-1}}^\bullet$, where $c = \beta\sigma_L f_L \bar{\tau}_L \bar{\delta} \bar{\gamma} \bar{\tau}_R \bar{f}_R \sigma_R \alpha$ and where the scalar $-\lambda\mu^{-1}$ is placed on a direct homotopy letter of $\alpha, \beta, \bar{\gamma}$ or $\bar{\delta}$.*

Proposition 4.3. *Let $\sigma = \beta\sigma_L\sigma_C\sigma_R\alpha$ be a homotopy band and $\tau = \delta\tau_L\tau_C\tau_R\gamma$ be a homotopy string. Suppose $f^\bullet: B_{\sigma,\lambda}^\bullet \rightarrow P_\tau^\bullet$ is a double map with components (f_L, f_R) . Then $M_{f^\bullet} \simeq P_c^\bullet$, where $c = \delta\tau_L f_L \bar{\sigma}_L \bar{\beta} \bar{\alpha} \bar{\sigma}_R \bar{f}_R \tau_R \gamma$.*

Proposition 4.4. *Let $\sigma = \beta\sigma_L\sigma_C\sigma_R\alpha$ be a homotopy string and $\tau = \delta\tau_L\tau_C\tau_R\gamma$ be a homotopy band. Suppose $f^\bullet: P_\sigma^\bullet \rightarrow B_{\tau,\mu}^\bullet$ is a double map with components (f_L, f_R) . Then $M_{f^\bullet} \simeq P_c^\bullet$, where $c = \beta\sigma_L f_L \bar{\tau}_L \bar{\delta} \bar{\gamma} \bar{\tau}_R \bar{f}_R \sigma_R \alpha$.*

5. MAPPING CONES AND QUASI-GRAPH MAPS

Single maps and double maps that are not singleton occur in a homotopy class that is determined by a quasi-graph map $P_\sigma^\bullet \rightsquigarrow \Sigma^{-1}P_\tau^\bullet$. In this case, it is possible to read off the mapping cone of any representative of the homotopy class from the quasi-graph map. Before we state how this is done, we need the following bookkeeping definition to cover the case when a quasi-graph map is supported in precisely one cohomological degree.

Definition 5.1. Let σ and τ be homotopy strings or bands and suppose $\varphi: Q_\sigma^\bullet \rightsquigarrow \Sigma^{-1}Q_\tau^\bullet$ is a quasi-graph map supported in exactly one degree, i.e. corresponds to the following diagram

$$\begin{array}{ccccccc} \beta & & \sigma_L & & \sigma_R & & \alpha \\ \sim & \bullet & \xrightarrow{\quad} & \bullet & \xrightarrow{\quad} & \bullet & \sim \\ & & & \parallel & & & \\ \delta & \bullet & \xrightarrow{\quad} & \bullet & \xrightarrow{\quad} & \bullet & \sim \\ & & \tau_L & & \tau_R & & \gamma \end{array}$$

We say that the homotopy strings or bands σ and τ are *compatibly oriented on the left* if

- (1) If σ_L is direct then τ_L is either empty, inverse with $\sigma_L\tau_L \neq 0$ or direct with $\tau_L = \sigma'_L\sigma_L$ for some nontrivial σ'_L .
- (2) If σ_L is inverse then τ_L is inverse and $\sigma_L = \tau_L\tau'_L$ for some nontrivial τ'_L .
- (3) If $\sigma_L = \emptyset$ then τ_L is inverse.

Similarly, we say that the homotopy strings or bands σ and τ are *compatibly oriented on the right* if the following dual conditions hold:

- (1) If σ_R is inverse then τ_R is either empty, direct with $\sigma_R\tau_R \neq 0$ or inverse with $\tau_R = \sigma'_R\sigma_R$ for some nontrivial σ'_R .
- (2) If σ_R is direct then τ_R is direct and $\sigma_R = \tau_R\tau'_R$ for some nontrivial τ'_R .
- (3) If $\sigma_R = \emptyset$ then τ_R is direct.

We say that the homotopy strings or bands σ and τ are *compatibly oriented for φ* if they are compatibly oriented on the left and on the right.

Note that when the maximal common homotopy substring ρ determining a quasi-graph map $\varphi: Q_\sigma^\bullet \rightsquigarrow \Sigma^{-1}Q_\tau^\bullet$ is of length at least one, the homotopy strings or bands σ and τ are automatically compatibly oriented for φ in an unfolded diagram of φ .


Proposition 5.2. *Let σ and τ be homotopy strings or bands. Suppose $\varphi: Q_\sigma^\bullet \rightsquigarrow \Sigma^{-1}Q_\tau^\bullet$ is a quasi-graph map determined by a maximal common homotopy substring ρ , i.e. $\sigma = \beta\sigma_L\rho\sigma_R\alpha$ and $\tau = \delta\tau_L\rho\tau_R\gamma$. Assume further that σ and τ are compatibly oriented for φ . Suppose $f^\bullet: Q_\sigma^\bullet \rightarrow Q_\tau^\bullet$ is a representative of the homotopy set determined by φ .*

- (1) *If σ and τ are homotopy strings then M_{f^\bullet} is isomorphic to $P_{c_1}^\bullet \oplus P_{c_2}^\bullet$, where $c_1 = \beta\sigma_L\rho\tau_R\gamma$ and $c_2 = \delta\tau_L\rho\sigma_R\alpha$.*


- (2) If (σ, λ) and (τ, μ) are homotopy bands then $M_{f_\bullet}^\bullet$ is isomorphic to $B_{c, -\lambda\mu^{-1}}^\bullet$, where $c = \beta\sigma_L\rho\tau_R\gamma\delta\tau_L\rho\sigma_R\alpha$ and where the scalar $-\lambda\mu^{-1}$ is placed on a direct homotopy letter of α or β or an inverse homotopy letter of γ or δ .
- (3) If (σ, λ) is a homotopy band and τ is a homotopy string then $M_{f_\bullet}^\bullet$ is isomorphic to P_c^\bullet , where $c = \delta\tau_L\rho\sigma_R\alpha\beta\sigma_L\rho\tau_R\gamma$.
- (4) If σ is a homotopy string and (τ, μ) is a homotopy band then $M_{f_\bullet}^\bullet$ is isomorphic to P_c^\bullet , where $c = \beta\sigma_L\rho\tau_R\gamma\delta\tau_L\rho\sigma_R\alpha$.

Proof. For ease of notation, we assume σ and τ are homotopy strings. Let $\varphi: P_\sigma^\bullet \rightsquigarrow \Sigma^{-1}P_\tau^\bullet$ be a quasi-graph map determined by a maximal common homotopy string ρ . We note that by orienting the unfolded diagrams so that they are compatibly oriented for φ we have interchanged the roles of γ and δ and τ_L and τ_R vis-à-vis the statements in Sections 3 and 4. Firstly, if the homotopy class determined by φ contains a double map, then we simply compute the mapping cone of such a double map as in Theorem 4.1 and Propositions 4.2, 4.3 and 4.4. So we may assume that the homotopy class of φ does not contain a double map. By [4, Prop. 4.8] this occurs if and only if quasi-graph map endpoint conditions (RQ3) and (LQ3) do not occur.

We first assume that ρ is a trivial homotopy string and choose a single map representative of the homotopy class determined by φ . Since ρ is trivial, this corresponds to the situation in Theorem 3.2 where precisely one of f_L or f_R is trivial. Assume f_R is nontrivial and f_L is trivial. The left-hand diagram below indicates this situation, where the notation has been updated to reflect the set up of the quasi-graph map in Definition 5.1, indicated on the right-hand side.



Here σ_L corresponds to the homotopy letter σ_{i+1} and τ_L corresponds to $\overline{\tau_{j-1}}$ in the proof of Theorem 3.2. Since f_R is nontrivial, the map $i_1^\bullet: P_{c_1}^\bullet \rightarrow M_{f_\bullet}^\bullet$ is defined as in Figure 1 in the proof of Theorem 3.2. However, the map $i_2^\bullet: P_{c_2}^\bullet \rightarrow M_{f_\bullet}^\bullet$ defined in Figure 2 is only well defined in the case that σ_L is direct. In the case that σ_L is inverse, we need to modify the definition of $i_2^\bullet: P_{c_2}^\bullet \rightarrow M_{f_\bullet}^\bullet$ as follows. First note that this means that φ satisfies the left endpoint condition (LQ1), meaning that $\sigma_L = \overline{g'g}$ and $\tau_L = \overline{g}$ for some nontrivial paths g and g' with $gg' \neq 0$. Then $i_2^\bullet: P_{c_2}^\bullet \rightarrow M_{f_\bullet}^\bullet$ can be modified as in the diagram below, with the remaining components defined as in Figure 2.




The argument when f_R is trivial and f_L is nontrivial is dual to the argument above.

Finally, in the case that both f_L and f_R are trivial, then $\text{len}(\rho) > 0$. In this case, one modifies i_1^\bullet above using identity morphisms with alternating signs along the top string until one reaches a direct homotopy letter or an endpoint as indicated in the second diagram above. Similarly for i_2^\bullet , where one proceeds until one reaches an inverse homotopy letter or an endpoint as the first diagram above. When one of σ or τ is a homotopy band, one deals with the signs exactly as in Propositions 3.4, 3.5 and 3.6. \square

6. EXAMPLES

In this section we will illustrate the graphical mapping cone calculus in $D^b(\Lambda)$ developed in Sections 2-5 on some concrete examples. In the first example, we consider maps involving only string complexes and in the second example we consider band complexes.


Example 6.1. Let Λ be the gentle algebra given by the following quiver and relations:



(1) Consider the homotopy strings $\sigma = edcbad\bar{}$ and $\tau = \bar{e}fcbaf\bar{e}$, and the graph map $f^\bullet: P_\sigma^\bullet \rightarrow P_\tau^\bullet$ given by

$$\begin{array}{cccccccccccc}
 P_\sigma & & P(3) & \xrightarrow{e} & P(4) & \xrightarrow{dc} & P(2) & \xrightarrow{b} & P(1) & \xrightarrow{a} & P(0) & \xleftarrow{d} & P(4) \\
 \downarrow & & & & & & \parallel & & \parallel & & \downarrow f & & \\
 P_\tau & & P(4) & \xleftarrow{e} & P(3) & \xleftarrow{f} & P(0) & \xrightarrow{c} & P(2) & \xrightarrow{b} & P(1) & \xrightarrow{af} & P(3) & \xrightarrow{e} & P(4)
 \end{array}$$


By Theorem 2.2, the mapping cone M_{f^\bullet} is isomorphic to $P_{c_1}^\bullet \oplus P_{c_2}^\bullet$ where $c_1 = d\bar{a}afe = dfe$ and $c_2 = edc\bar{c}fe = edfe$ (cf. green and red boxes in the figure below).



(2) Non-singleton double maps and single maps arise in the context of quasi-graph maps. As described in Section 1.4.4, a given quasi-graph map gives rise to a class of (single and double) maps, which are all homotopy equivalent to each other. In particular, they all have the same mapping cone, which is the ‘mapping cone of the quasi-graph map’. We will now illustrate this with an example. Consider a quasi-graph map $\varphi: P_\sigma^\bullet \rightsquigarrow \Sigma^{-1}P_\tau^\bullet$, for homotopy strings $\sigma = bacb$ and $\tau = \bar{f}cba$, given by

$$\begin{array}{cccccccccccc}
 P_\sigma & & P(2) & \xrightarrow{b} & P(1) & \xrightarrow{a} & P(0) & \xrightarrow{c} & P(2) & \xrightarrow{b} & P(1) \\
 \downarrow \text{zigzag} & & & & & & \parallel & & \parallel & & \parallel \\
 \Sigma^{-1}P_\tau & & P(3) & \xleftarrow{f} & P(0) & \xrightarrow{c} & P(2) & \xrightarrow{b} & P(1) & \xrightarrow{a} & P(0)
 \end{array}$$

By Proposition 5.2(1), the mapping cone of any single or double map $f^\bullet: P_\sigma^\bullet \rightarrow P_\tau^\bullet$ in the homotopy set determined by φ is isomorphic to $P_{c_1}^\bullet \oplus P_{c_2}^\bullet$, where $c_1 = bacba$ and $c_2 = \bar{f}cb$ (cf. green and red boxes in the figure below).



We now consider a single map $f^\bullet: P_\sigma^\bullet \rightarrow P_\tau^\bullet$ and a double map $g^\bullet: P_\sigma^\bullet \rightarrow P_\tau^\bullet$ in the homotopy set determined by the quasi-graph map φ .

(i) Let $f^\bullet : P_\sigma^\bullet \rightarrow P_\tau^\bullet$ be a single map in the homotopy set determined by φ given by

$$\begin{array}{ccccccc}
 P_\sigma & & P(2) & \xrightarrow{b} & P(1) & \xrightarrow{a} & P(0) & \xrightarrow{c} & P(2) & \xrightarrow{b} & P(1) \\
 \downarrow & & & & & & \downarrow c & & & & \\
 P_\tau & & P(0) & \xleftarrow{a} & P(1) & \xleftarrow{b} & P(2) & \xleftarrow{c} & P(0) & \xrightarrow{f} & P(3)
 \end{array}$$

where we have drawn the unfolded diagram so that it is compatibly oriented (see Definition 3.1). By Theorem 3.2, the mapping cone is $M_{f^\bullet}^\bullet \cong P_{c_1}^\bullet \oplus P_{c_2}^\bullet$, where $c_1 = bacba$ and $c_2 = \bar{f}cb$ (cf. green and red boxes in the figure below).

$$\begin{array}{ccccccc}
 \boxed{P(2) \xrightarrow{b} P(1) \xrightarrow{a} P(0)} & \xrightarrow{c} & \boxed{P(2) \xrightarrow{b} P(1)} \\
 \downarrow c & & \uparrow c \\
 \boxed{P(0) \xleftarrow{a} P(1) \xleftarrow{b} P(2)} & \xleftarrow{c} & \boxed{P(0) \xrightarrow{f} P(3)}
 \end{array}$$

(ii) Let $g^\bullet : P_\sigma^\bullet \rightarrow P_\tau^\bullet$ be a double map in the homotopy set determined by φ given by

$$\begin{array}{ccccccc}
 P_\sigma & & P(2) & \xrightarrow{b} & P(1) & \xrightarrow{a} & P(0) & \xrightarrow{c} & P(2) & \xrightarrow{b} & P(1) \\
 \downarrow & & & & \downarrow a & & \downarrow f & & & & \\
 P_\tau & & P(0) & \xleftarrow{a} & P(1) & \xleftarrow{b} & P(2) & \xleftarrow{c} & P(0) & \xrightarrow{f} & P(3)
 \end{array}$$

By Theorem 4.1, its mapping cone is $M_{g^\bullet}^\bullet \cong P_{c_1}^\bullet \oplus P_{c_2}^\bullet$, where $c_1 = \bar{f}cb$ and $c_2 = bacba$ (cf. red and green boxes in the figure below).

$$\begin{array}{ccccccc}
 \boxed{P(2) \xrightarrow{b} P(1)} & \xrightarrow{a} & \boxed{P(0) \xrightarrow{c} P(2) \xrightarrow{b} P(1)} \\
 \downarrow a & & \downarrow f \\
 \boxed{P(0) \xleftarrow{a} P(1) \xleftarrow{b} P(2)} & \xrightarrow{f} & \boxed{P(3)}
 \end{array}$$

We finally give an example involving band complexes.

Example 6.2. Let Λ be the algebra given by the quiver with relations in Example 2.10. Let $\sigma = gh\bar{d}\bar{f}$ and $\tau = b\bar{e}\bar{d}c$ be homotopy bands with corresponding scalars λ and μ respectively. By Proposition 2.9, the mapping cone for the graph map $f^\bullet : B_{\sigma,\lambda}^\bullet \rightarrow B_{\tau,\mu}^\bullet$ given below is $M_{f^\bullet}^\bullet \cong B_{c,\lambda\mu^{-1}}^\bullet$, where $c = ghe\bar{b}\bar{c}d\bar{d}\bar{f} = ghe\bar{b}\bar{c}\bar{f}$. Graphically this corresponds to the following diagrams of unfolded complexes.

$$\begin{array}{ccccccc}
 \dots\dots\dots P(6) & \xrightarrow{\lambda gh} & P(5) & \xleftarrow{df} & P(6) & \dots\dots\dots \\
 \downarrow \lambda ghe & & \parallel & & \downarrow f & \\
 \dots\dots\dots P(3) & \xrightarrow{b} & P(2) & \xleftarrow{e} & P(5) & \xleftarrow{d} & P(4) & \xrightarrow{\mu c} & P(3) & \dots\dots\dots
 \end{array}$$

REFERENCES

- [1] C. Amiot, Y. Grimeland, *Derived invariants for surface algebras*, J. Pure Appl. Algebra **220** (2016), 3133–3155.
- [2] C. Amiot, D. Labardini-Fragoso, P.-G. Plamondon, *Derived invariants for surface cut algebras II: the punctured case*, arXiv:1606.07364.
- [3] S. Ariki, K. Iijima, E. Park, *Representation type of finite quiver Hecke algebras of type $A_\ell^{(1)}$ for arbitrary parameters*, Int. Math. Res. Not. IMRN 2015, no. 15, 6070–6135.
- [4] K. K. Arnesen, R. Laking, D Pauksztello, *Morphisms between indecomposable complexes in the bounded derived category of a gentle algebra*, J. Algebra **467** (2016), 1–46.
- [5] I. Assem, *Tilted algebras of type A_n* , Comm. Algebra **9** (1982), 2101–2125.

- [6] I. Assem, T. Brüstle, G. Charbonneau-Jodoin, P-G. Plamondon, *Gentle algebras arising from surface triangulations*, Algebra Number Theory **4** (2010), 201–229.
- [7] I. Assem, A. Skowroński, *Iterated tilted algebras of type \tilde{A}* , Math. Z. **195** (1987), 269–290.
- [8] V. Bekkert, H. Merklen, *Indecomposables in derived categories of gentle algebras*, Algebr. Represent. Theory **6** (2003), 285–302.
- [9] G. Bobiński, *The almost split triangles for perfect complexes over gentle algebras*, J. Pure Appl. Algebra **215** (2011), 642–654.
- [10] R. Bocklandt, *A dimer ABC*, Bull. Lond. Math. Soc. **48** (2016), no. 3, 387–451.
- [11] T. Bridgeland, *Stability conditions on triangulated categories*, Ann. of Math. **166** (2007), 317–345.
- [12] N. Broomhead, *Thick subcategories of discrete derived categories*, arXiv:1608.06904.
- [13] A. B. Buan, R. Marsh, M. Reineke, I. Reiten, G. Todorov, *Tilting theory and cluster combinatorics*, Adv. Math. **204** (2006), 572–618.
- [14] R-O. Buchweitz, *Maximal Cohen-Macaulay modules and Tate cohomology over Gorenstein rings*, unpublished manuscript.
- [15] I. Burban, Y. Drozd, *Derived categories of nodal algebras*, J. Algebra **272** (2004), 46–94.
- [16] I. Burban, Y. Drozd, *Maximal Cohen-Macaulay modules over non-isolated surface singularities and matrix problems*, Mem. Amer. Math. Soc. 248 (2017), no. 1178.
- [17] I. Burban, Y. Drozd, *Tilting on non-commutative rational projective curves*, Math. Ann. **351** (2011), 655–709.
- [18] P. Caldero, F. Chapoton, R. Schiffler, *Quivers with relations arising from clusters (A_n case)*, Trans. Amer. Math. Soc. **358** (2006), 1347–1364.
- [19] İ. Çanakçı, D. Pauksztello, S. Schroll, *On extensions for gentle algebras*, arXiv:1707.06934.
- [20] İ. Çanakçı, S. Schroll, *Extensions in Jacobian algebras and cluster categories of marked surfaces*, Adv. Math. **313** (2017), 1–49.
- [21] A. T. Carroll, *Generic modules for gentle algebras*, J. Algebra **437** (2015), 177–201.
- [22] A. T. Carroll, C. Chindris, *On the invariant theory for acyclic gentle algebras*, Trans. Amer. Math. Soc. **367** (2015), 3481–3508.
- [23] C. Cibils, *A finite dimensional algebra of the diagram of a knot*, Proc. Amer. Math. Soc. 142 (2014), no. 11, 3741–3746.
- [24] R. Coelho Simões, M. Parsons, *Endomorphism algebras for a class of negative Calabi-Yau categories*, J. Algebra **491** (2017), 32–57.
- [25] R. Coelho Simões, D. Pauksztello, *Torsion pairs in a triangulated category generated by a spherical object*, J. Algebra **448** (2016), 1–47.
- [26] Y. A. Drozd, *Tame and wild matrix problems* (in Russian), in “Representations and quadratic forms”, 39–74, Institute of Mathematics, Academy of Sciences, Ukrainian SSR, Kiev, 1977.
- [27] K. Erdmann, *On Hochschild cohomology for selfinjective special biserial algebras*, Algebras, quivers and representations, 79–94, Abel Symp., 8, Springer, Heidelberg, 2013.
- [28] S. Fomin, M. Shapiro, D. Thurston, *Cluster algebras and triangulated surfaces. Part I: Cluster complexes*, Acta Mathematica **201** (2008), 83–146.
- [29] A. Garver, T. McConville, *Oriented Flip Graphs and Noncrossing Tree Partitions*, arXiv:1604.06009.
- [30] C. Geiß, H. Krause, *On the notion of derived tameness*, J. Algebra Appl. **1** (2002), 133–157.
- [31] S. Gratz, T. Holm, P. Jørgensen, *Cluster tilting subcategories and torsion pairs in Igusa–Todorov cluster categories of Dynkin type A_∞* , arXiv:1711.07528.
- [32] I. M. Gelfand, V. A. Ponomarev, *Indecomposable representations of the Lorentz group*, Russian Math. Surveys **23** (1968), 1–58.
- [33] F. Haiden, L. Katzarkov, M. Kontsevich, *Flat surfaces and stability structures*, Publ. Math. Inst. Hautes Études Sci. 126 (2017), 247–318.
- [34] D. Happel, *Triangulated Categories in the Representation Theory of Finite Dimensional Algebras*, London Math. Soc. Lecture Notes Series **119**, Cambridge University Press (1988).
- [35] K. Igusa, G. Todorov, *Cluster categories coming from cyclic posets*, Comm. Alg. **43** (2015), 4367–4402.
- [36] O. Iyama, Y. Yoshino, *Mutation in triangulated categories and rigid Cohen-Macaulay modules*, Invent. Math. **172** (2008), 117–168.
- [37] M. Kalck, *Singularity categories of gentle algebras*, Bull. London Math. Soc. **47** (2015), 65–74.
- [38] Y. Lekili, A. Polishchuk, *Derived equivalences of gentle algebras via Fukaya categories*, arXiv:1801.06370.
- [39] D. Labardini-Fragoso, *Quivers with potentials associated to triangulated surfaces*, Proc. London Math. Soc. **98** (2009), 797–839.
- [40] P. Ng, *A characterization of torsion theories in the cluster category of Dynkin type A_∞* , arXiv:1005.4364.
- [41] S. Oppen, P-G. Plamondon, S. Schroll, *A geometric model for the derived category of gentle algebras*, arXiv:1801.09659.
- [42] Y. Qiu, Y. Zhou, *Cluster categories for marked surfaces: punctured case*, Compos. Math. 153 (2017), no. 9, 1779–1819.
- [43] C. M. Ringel, *The indecomposable representations of the dihedral 2-groups*, Math. Ann. **214** (1975), 19–34.

- [44] J. Schröer, *Modules without self-extensions over gentle algebras*, J. Algebra **216** (1999), 178–189.
- [45] J. Schröer, A. Zimmermann, *Stable endomorphism algebras of modules over special biserial algebras*, Math. Z. **244** (2003), 515–530.
- [46] S. Schroll, *Trivial extensions of gentle algebras and Brauer graph algebras*, J. Algebra **444** (2015), 183–200.
- [47] M. van den Bergh, *Non-commutative crepant resolutions*, in “The Legacy of Niels Henrik Abel”, 749–770, Springer, Berlin 2004.
- [48] B. Wald, J. Waschbüsch, *Tame biserial algebras*, J. Algebra **95** (1985), 480–500.
- [49] D. Xu, *Auslander-Reiten conjecture and special biserial algebras*, Arch. Math. (Basel) **105** (2015), 13–22.
- [50] J. Zhang, Y. Zhou, B. Zhu, *Cotorsion pairs in the cluster category of a marked surface*, J. Algebra **391** (2013), 209–226.

SCHOOL OF MATHEMATICS, STATISTICS AND PHYSICS, NEWCASTLE UNIVERSITY, NEWCASTLE UPON TYNE, NE1 7RU, UNITED KINGDOM.

E-mail address: `ilke.canakci@ncl.ac.uk`

DEPARTMENT OF MATHEMATICS AND STATISTICS, LANCASTER UNIVERSITY, LANCASTER, LA1 4YF, UNITED KINGDOM

E-mail address: `d.pauksztello@lancaster.ac.uk`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF LEICESTER, UNIVERSITY ROAD, LEICESTER, LE1 7RH, UNITED KINGDOM.

E-mail address: `schroll@leicester.ac.uk`