

This is the Accepted Manuscript version of “Re-evaluating the DSM-I” by Rachel Cooper and Roger Blashfield which is forthcoming in *Psychological Medicine*. This version has been accepted for publication and will appear in a revised form post copyediting. Please see <http://journals.cambridge.org/action/displayJournal?jid=PSM> for the final version.

The *Diagnostic and Statistical Manual: Mental Disorders* (1952), now generally known as the DSM-I, attracts little attention in the current literature. The received view is that this system was a relatively unimportant, psychoanalytically-oriented classification. For example, Allan Horwitz (2015) claimed that

All of the approximately 100 diagnostic definitions in the *DSM-I* were short, cursory, and infused with psychodynamic assumptions. (Horwitz, 2015, p.2)

In her history of the DSM-III Hannah Decker stated

Little attention was paid to the first two DSMs, which were published mainly for psychiatrists in state mental hospitals who were interested in compiling a variety of statistical information on their patients' lives and deaths. (Decker, 2013, p.xvii)

Gerald Grob, whose (1991) “Origins of DSM-I: A Study in Appearance and Reality” remains the seminal discussion of this classification, noted

The publication of DSM-I marked an internal transformation that mirrored the growing dominance of psychodynamic and psychoanalytic psychiatry and the relative weakness of the biological tradition. (Grob, 1991, p.421)

The goal of this paper is to challenge this received history. We will argue that the DSM-I was not psychoanalytic; rather it was eclectic, owing as much to Kraepelin as Freud. In addition, the DSM-I was used frequently in its time. Copies circulated widely, and the manual went through twenty printings (A.P.A., 1968, p.ix). In short, the DSM-I was much more like recent editions of the DSM than generally thought. This matters because modern writers have often severed later editions of the DSM from their historical context. Conceiving of the DSM-I as psychoanalytic and professionally unimportant has led to it being dismissed as a mere artefact of a long gone era. DSM-I and II are presented as having only their name in common with later editions, and the current epoch in psychiatry seems to start in 1980. For some this has resulted in an assumption that the classificatory strategies of the more recent DSMs delineate the only ways in which “right thinking” psychiatrists have ever divided up the domain of psychopathology. For others (critics of the DSM) the idea that the current DSM categories appeared from nowhere around a committee table in the late 1970s has facilitated their too-easy dismissal (compare, for example, Maxmen, 1985, p.35 and Whitaker, 2010, pp. 269-271; Shorter, 2015, p2). In our history, the continuities between the first DSMs (and earlier classifications) and the more recent editions DSMs become much more visible.

1. The DSM-I was not a psychoanalytic classification

We start by raising four initial points that should make readers doubt that the DSM-I can be characterized as psychoanalytic. With a prima facie case established, we then

turn to consider the way in which the DSM-I was developed. Demonstrating the influence that various prior classifications had on the DSM-I will further support our argument.

1.1. The DSM-I was not a psychoanalytic classification – the prima facie case.

1.1.1. Psychoanalysts did not write the DSM-I

The DSM-I included a list of twenty-eight individuals who served for some portion of time on the APA Committee of Nomenclature and Statistics between 1946 and 1951 (A.P.A., 1952, p.xii). These are the individuals who were credited with producing the DSM-I. The *Biographical Directory of the American Psychiatric Association* for 1950 contained details regarding twenty-five of these individuals (the other three had died prior to 1950). Of these 25, only seven (28%) had either trained as psychoanalysts, or were members of psychoanalytic societies (specifically, George N. Raines, Moses M. Frohlich, Franz Alexander, Norman Q. Brill, Jacob H. Friedman, Nolan D. C. Lewis and George S. Sprague). Of the three committee members whose details cannot be found in the 1950 *Biographical Directory*, the obituary of Jacob Kasanin listed him as a member of the American Psychoanalytic Association (Anonymous, 1946). Clarence O. Cheney and James V. May were one-time Presidents of the APA and the *American Journal of Psychiatry* contains extensive obituaries for them (APA, 1949a). These obituaries make no mention of any psychoanalytic training, or membership of psychoanalytic societies, and these two were thus almost certainly not psychoanalysts. This means that only 29% of the 28 individuals responsible for the DSM-I can be characterized as psychoanalysts.

The 1950 *Biographical Directory* also provides details of place of work and publications. The twenty-eight committee members had diverse interests (as catalogued in Anonymous, 1946; APA, 1949a; APA 1950). Many had military experience (notably Norman Brill, John Caldwell, J.P.S. Cathcart, Jacob Friedman, Moses Frohlich, Baldwin Keyes, George Raines). Others had worked for the Veteran's Administration (notably John Baird, George Brewster, Moses Frohlich, Harvey Tompkins). At least three had worked in State Mental Hospitals (Clarence O'Cheney, Jacob Kasanin, James May). There were notable neurologists (John Baird, Walter Breutsch, Norman Brill, Houston Merritt, Robert Schwab). Neil Dayton and James May had long standing interests in classification and epidemiology. Baldwin Keyes and Mabel Ross were child psychiatrists; Lawrence Kolb was a specialist on addictions; Abram Bennett and Walter Breutsch worked on somatic therapies.

Members of the committee were assigned to small groups, each of which was responsible for some section of the manual, for example, one group was responsible for the nomenclature of psychoneuroses, another for the psychoses of unknown origin, and so on (Menninger, 1963, p.475). Committee members would most likely have been allocated to groups where their expertise was greatest, making it probable that psychoanalysts would have greater input into certain sections of the manual. Still, given that overall less than one third of the committee members were psychoanalysts, and that the committee members had a wide range of interests, the DSM-I cannot be considered a psychoanalytic classification in the sense of having been written by analysts.

1.1.2. Psychoanalysts did not use the DSM-I

The DSM-I cannot be considered a psychoanalytic classification in the sense of being a classification used by psychoanalysts, as psychoanalysts in the 1950s tended not to employ the DSM-I. Between 1954 and 1958 a committee of the American Psychoanalytic Association tried to collect statistical data with the aim of measuring the effectiveness of psychoanalysis in treating particular conditions (Weinstock, 1965). They decided to use DSM-I to code the diagnosis of the patients. Psychoanalysts participating in the study were each sent a free copy, suggesting that psychoanalysts were unaccustomed to using DSM diagnoses. Early in the course of the study it became apparent that the psychoanalysts were unwilling or unable to use the DSM diagnoses. Fewer and fewer psychoanalysts returned their forms. Many of those who did continue to participate either left the diagnosis unspecified or used the same diagnosis, most often passive-aggressive personality, compulsive personality or pseudoneurotic schizophrenia (which was not a DSM-I diagnosis) for all their patients. Others abandoned the diagnostic section of the questionnaires altogether. In addition to finding the DSM unusable, the psychoanalysts were uninterested in developing their own diagnostic system. The committee invited the membership to form groups to study the problem of diagnosis, “..but as far as could be determined, only one group met and that for only one discussion.”(Weinstock, 1965,p. 68).

1.1.3. The DSM-I was not “infused” with psychoanalytic theory

Horwitz claimed that the DSM-I was “infused with psychoanalytic” theory, and provided examples of the influence of psychoanalytic thinking in the DSM-I to prove his point (2015). One of the major families of disorders in the DSM-I were the “Psychoneurotic Disorders”. The first sentences from the DSM-I description for these disorders stated:

The chief characteristic of these disorders is “anxiety” which may be directly felt and expressed or which may be unconsciously and automatically controlled by the utilization of various psychological defense mechanisms (depression, conversion, displacement, etc.) . . .

‘Anxiety’ in psychoneurotic disorders is a danger signal felt and perceived by the conscious portion of the personality. It is produced by a threat from within the personality (e.g., by supercharged repressed emotions, including such aggressive impulses as hostility and resentment), with or without stimulation from such external situations as loss of love, loss of prestige, or threat of injury. (APA, 1952, pp. 31-32)

Here the psychoanalytic influence is plain to see. If the definitions of all or even most of the mental disorders read this way, Horwitz's claim would be correct.

The hierarchical organization of the DSM-I was a top-down system that was designed to reflect clinical thinking about psychopathology at the time. The DSM-I subdivided mental disorders into (I) those with an organic basis (“Disorders caused by or associated with impairment of brain tissue function”), (II) those for which an organic basis was unknown, (“Disorders of psychogenic origin or without clearly defined physical cause or structural change in the brain”) and (III) the “mental deficiency” disorders. The organic disorders were split into (I.A) acute (13 diagnoses) vs. (I.B.) chronic forms (26 diagnoses). The (II) non-organic disorders were split into (II.A) psychotic disorders (20 diagnoses), (II.B) psychophysiological disorders (10 diagnoses), (II.C) psychoneurotic disorders (8 diagnoses), (II.D) personality disorders (22 diagnoses) and (II.E) transient situational personality disorders (10 diagnoses). Notably, the DSM-I class of “personality disorders” was wide and heterogeneous, including not only what would now be considered personality disorders, but

also sexual deviations, addictions, and “special symptom reactions”, such as speech disturbances and enuresis.

Of the 127 diagnoses in the DSM-I (including non-diagnostic terms), there were 44 categories that could be said to have partly psychoanalytic definitions: the psychophysiological disorders, the psychoneurotic disorders and the personality disorders (minus 6 diagnoses under Sociopathic Personality Disturbance). Thus, 35% of the diagnoses in the DSM-I could be said to be psychoanalytic. In contrast, 46 of the DSM-I diagnoses (36%) descended directly from Kraepelin's classification (e.g., general paralysis, schizophrenic reaction (dementia praecox), manic-depressive psychosis, etc., see Aragona, 2015, for a useful chart showing similarities between the DSM-I and Kraepelin's classification). As a result, categories with psychoanalytic terminology in their definitions did not represent the majority of diagnoses. The DSM-I drew as much on Kraepelinian concepts as it did psychoanalytic concepts.

Of course, not all DSM-I diagnoses were of equal importance; some diagnoses were used far more frequently than others. If the psychoanalytically-influenced DSM-I diagnoses were those most used in practice then perhaps the DSM-I-as-used might be considered psychoanalytic. In 1955, a year when the DSM-I had been widely adopted and statistics are available, many psychiatric patients in the U.S. continued to be treated as inpatients. That year, public mental hospitals housed 559,000 patients (excluding those in VA facilities), while an estimated 379,000 were seen as outpatients (Kramer and Pollock 1958, Bahn and Norman, 1959). Amongst inpatients, the most common diagnoses were the schizophrenic reactions (51%), cerebral arteriosclerosis (9%), and mental deficiency (8%) (National Institute of Mental Health, 1964). Outpatient clinics at the time saw more children than adults (Bahn and Norman, 1959), and the most common diagnoses given to children were Transient

Situational Personality Disorders (36%), Personality Disorders (which in the DSM-I included diagnoses for learning disturbance, speech disturbance and enuresis) (21%), and Mental Deficiency (18%). The most common adult diagnoses in outpatient clinics were Personality Disorders (which included sexual deviations and addictions) (32%), Psychoneurotic Disorders (31%), and Psychotic Disorders (20%) (Bahn and Norman, 1959). Thus, both in terms of the numbers of categories included in the DSM-I, and in terms of the numbers of patients diagnosed, the psychoanalytically-influenced diagnoses, though significant, were not dominant.

1.1.4. Contemporaries of the DSM-I did not consider it psychoanalytic

Contemporary characterizations of the DSM-I support our claim. George N. Raines was chairman of the committee that produced the DSM-I. In his view,

The present new nomenclature of The American Psychiatric Association follows...the general nomenclatural scheme of Adolf Meyer, utilizes the names originated by Kraepelin and Bleuler, and incorporates the dynamics developed by Freud and later analysts, *wherever these are applicable*. (emphasis in original, Raines, 1953a, p.425)

Commentators from the time of the DSM-I saw both a Kraepelinian and psychoanalytic influence,

It seems to be inconceivable that the Diagnostic and Statistical Manual: Mental Diseases could have been written without the preceding work of Kraepelin. (Kahn, 1955, p.395)

The modern taxonomy adopted by the American Psychiatric Association is a direct descendant of the old Kraepelinian classification of mental disorders as definite disease entities. (Blinder, 1966, p.259)

The definition of terms in the new nomenclature are largely descriptive.... In addition to the classical descriptive material, however, the revised nomenclature includes in its definitions a certain amount of psychodynamic theory as to the nature and origin of the conditions described (Bowman & Rose, 1951, p.165)

With a prima facie case established that the DSM-I was not a purely psychoanalytic classification, we turn now to consider the way in which the DSM-I was developed. Demonstrating the influence that various prior classifications had on the DSM-I will further support our argument.

1.2 Development of the DSM-I

In producing the new manual, the American Psychiatric Association's Committee on Nomenclature and Statistics, drew on two series of earlier classifications. The first of these systems was the *Statistical Manual for the Use of Institutions for the Insane* (1918) and its successor volumes (called *Statistical Manual for the Use of Hospitals for Mental Diseases*), which was used to collect statistics on mental hospital populations (American Medico-Psychological Association, 1918; APA, 1942). The second influence was classifications designed for use with military personnel during World War II (Army, Navy, and VA).

1.2.1. The DSM-I as descended from the *Statistical Manual for the Use of Institutions for the Insane* (1918)

Gerald Grob (1991) reported that in 1908 the Bureau of the Census asked the American Medico-Psychological Association (later the APA) to appoint a Committee on Nomenclature of Diseases to assist with the collection of data. The Federal Census took a particular interest in those citizens who were dependent upon governmental care for their well-being, as it was hoped that statistical knowledge could inform the development of appropriate public policy. The American Medico-Psychological Association created the requested committee in 1913 and the first classification system, produced in collaboration with the National Committee for Mental Hygiene, was finally published in 1918, as *Statistical Manual for the Use of Institutions for the Insane* (American Medico-Psychological Association, 1918).

The 1918 *Statistical Manual* organized mental diseases into 21 groups (families of disorders). Included in these 21 were groups such as traumatic psychoses (with 3 subdivisions), general paralysis (no subdivisions), psychoses due to drugs and other exogenous toxins (4 subdivisions), and dementia praecox (4 subdivisions). Thirteen of these 21 groups were “associated with organic brain disturbance” (in DSM-I terms). Of the remaining 8 groups, 4 were non-organic psychotic disorders.

James V. May sat on the committee that produced the 1918 classification. His textbook *Mental Diseases* (1922) discussed the origins of the classification. May noted that Kraepelin’s classification had been introduced to the United States by Adolph Meyer and August Hoch and became widely accepted (May, 1922, p244). The 1918 classification sought to reflect the contemporary consensus within US systems (May 1922 p.247) and was thus a modified Kraepelinian classification.

The 1918 manual went through a number of different editions. By 1929 the APA Committee of Statistics celebrated the success of the manual (now renamed *Statistical Manual for the Use of Institutions for Mental Diseases*).

The classification has been officially adopted by the Federal Census Bureau, The United States Public Health Service, the Surgeon-General of the Army, the United States Veterans Bureau, and by practically all the state hospitals for mental diseases in the United States. It is being taught in courses in psychiatry in many medical schools and is given with approval in the newer American textbooks in psychiatry (1929 report of APA Committee on Statistics, cited in Pollock, 1945,p. 10)

The *Statistical Manual* sold in fair quantities. The eighth edition (1934), for instance, sold 2700 copies in the first sixth-months of availability (APA, 1935, p.472), at a time when there were only 1510 APA members (APA, 1935, p.461). Compared to the 1918 edition of the manual, in the eighth edition the section for psychoneuroses was greatly expanded (going from 4 to 17 sub-divisions) (Dayton, 1935). The tenth edition, published in 1942, remained very similar to the eighth, although the ordering of some classes was revised (APA, 1942).

Published in 1952, the DSM-I was a continuation of this series of manuals for compiling hospital statistics. In comparing itself to the 10th edition of *The Statistical Manual*, the DSM-I described itself as a continuation of an existing series rather than as a paradigm-breaking document. The introduction to the DSM-I suggests that “Perhaps the greatest change in this revision from previous listings lies in the handling of the disorders with known organic etiologic factors” (APA, 1952, p.9); the DSM-I introduced a distinction between acute and chronic brain disorders. Other key changes were said to be as follows: the schizophrenic

reactions increased in numbers and type (including a schizo-affective type, and childhood type, not included in earlier classifications (p.27)); the manic depressive reactions were reduced in number, but a new code “000-x14 Psychotic depressive reaction” was included (p.25); and new, or greatly expanded, sections of “Psychophysiologic Autonomic and Visceral Disorders” (psychosomatic disorders), Personality Disorders and Transient Situational Personality Disorders (reactions to stress) were added. Morton Kramer, Chief of the Biometrics Branch of the National Institute of Mental Health (NIMH), prepared comparison sheets for contemporaries charting the main changes between the 10th edition of the *Statistical Manual for the Use of Institutions for Mental Diseases* and DSM-I. In Kramer's view “there were not...many basic differences” (Gottlieb, 1952, p.7).

1.2.2 The DSM-I as descended from Armed Forces Classifications

During WWII each of the U.S. armed forces, and the Veterans' Administration, developed its own classification for mental disorders. William C. Menninger was chiefly responsible for the Army classification, published as Technical Bulletin Medical 203 in 1943 (Houts, 2000). The classification caused some controversy. Menninger claims that he was initially discouraged from producing Medical 203 by the then chairman of the APA Committee for Nomenclature and Statistics (who would have been either Neil Dayton, chair 1942-1946, or Nolan Lewis, chair 1946-1948), and that the *American Journal of Psychiatry* refused to publish the classification (Menninger, 1947, p.582 ftnt 5).

Medical 203 was heavily influenced by psychoanalytic theory. The introduction explained:

The term “disorder” is used for the designation of the generic group of the specific reactions, while the specific reaction types have been termed “reactions”.

The classifications of the psychoneuroses are based on the dynamics of the

psychopathology. Of necessity, a few terms remained descriptive (symptomatic).

(Office of the Surgeon General, 2000 [orig. 1946], p.925)

Compared to the classifications intended for hospital use, the emphasis of Medical 203 was reversed. The classification started with “Simple personality reactions” (which encompassed transient personality reactions to acute or special stress, combat exhaustion, and acute situational maladjustment). Neurotic reactions, “Character and behavior reactions” and “Immaturity reactions” also received prominent billing. In contrast, the sections on psychotic disorders, especially affective disorders, were fairly brief. Very little was said about organic conditions. While organic disorders made up at least half of the *Statistical Manual for the Use of Hospitals for Mental Diseases*, the organic disorders in Medical 203 were dealt with in a single paragraph, relegated to the end of the classification.

In 1947 the Veteran’s Administration adopted a classification closely following Medical 203 (Veterans Administration, 1947). In turn, a pamphlet titled “Joint Armed Forces Nomenclature and Method of Recording Psychiatric Conditions” was released in 1949 (Army, U.S. et al., 1949). This sought to provide a common system for use in the Army, Navy and Air Force. Much of the wording remained the same as in Medical 203 and the VA classification - although the ordering of disorders differed. Given the similarities between Medical 203, the VA classification, and the Joint Armed Forces classification in what follows we will talk of ‘Medical 203/VA/Joint Forces’ when all three classifications are being discussed.

In places, especially in the schizophrenic reactions, psychophysiologic autonomic and visceral disorders, psychoneurotic disorders, and personality disorders, the text of DSM-I borrowed heavily from Medical 203/VA/Joint Forces. However, the influence of Medical 203/VA/Joint Forces was limited. In 1949, the APA Committee on Nomenclature and

Statistics reviewed the VA classification but was not overly impressed: it worried that the VA classification reflected the requirements of “expediency” and “enthusiasms of a temporary nature” (APA, 1949b, p.930). Where DSM-I borrowed phrasing from Medical 203/VA/Joint Forces, often the most “psychoanalytic” wording was deleted. In practice, it looks as if the DSM-I committee wrote sections of the DSM-I by taking Medical 203/VA/Joint Forces text and crossing out what they considered to be controversial psychoanalytic claims. Compare, for example, Medical 203/VA/Joint Forces and the DSM-I on Obsessive Compulsive Reaction. The former described this reaction as follows:

In this reaction the anxiety may be observable in connection with obsessional fear of uncontrollable impulses. On the other hand, the anxiety may be under apparent control, through a mental mechanism (isolation), by which the emotional charge becomes automatically separated from the main stream of consciousness and manifests itself in a displaced form through useless or excessive, and often repetitive activity. In the latter instance, the patient is utilizing the mental mechanisms of “undoing” – a symbolic act which temporarily protects the patient against a threat – and “displacement”. The patient himself may regard his ideas and behavior as unreasonable and even silly, but nevertheless is compelled to carry out his rituals. The diagnosis should specify the symptomatic expressions of such reactions, including touching, counting, ceremonials, handwashing, recurring thoughts, accompanied often by compulsion to repetitive action. This category includes many cases formerly classified as “psychasthenia”. (Office of the Surgeon General. 2000 [1946], p.928; Veterans Administration, 1947 p.5; Army et al., 1949,p7; italics and underlining added)

In the DSM-I, the last three sentences were almost identical (shown in underline). However, the initial part of the definition (shown in italics) was distinctly altered. The DSM-I version was more descriptive of the signs/symptoms of the disorder and was less about the theorized mental mechanisms that allegedly explained those signs/symptoms.

In this reaction the anxiety is associated with the persistence of unwanted ideas and of repetitive impulses to perform acts which may be considered morbid by the patient. The patient himself may regard his ideas and behavior as unreasonable, but nevertheless is compelled to carry out his rituals. The diagnosis will specify the symptomatic expression of such reactions, as touching, counting, ceremonials, hand-washing, or recurring thoughts (accompanied often by a compulsion to repetitive action). This category includes many cases formerly classified as “psychasthenia” (APA, 1952, p.33)

In contrast to the avowedly psychoanalytic orientation of Medical 203/VA/Joint Forces, the DSM-I presented itself a classification that “recognizes the present day descriptive nature of all psychiatric diagnoses” (APA, 1952, p.9). Still, not all psychoanalytic phrasing was deleted before text was inserted into DSM-I. For example, both Medical 203/VA/Joint Forces and the DSM-I described the paranoid personality as being characterized by a “tendency to utilize a projection mechanism” (APA, 1952, p.36; Office of the Surgeon General, 2000 [1946], p.929). In addition to borrowing wording from Medical 203/VA/Joint Forces in describing disorders, the DSM-I also adhered to the terminology of the earlier classifications by often describing disorders as “reactions” (APA, 1952, p.9).

In summary, the influence of Medical 203/VA/Joint Forces on DSM-I was limited. Most obviously, about half of the DSM-I concerned organic disorders; Medical 203/VA/Joint Forces had no influence on these disorders. Among the non-organic conditions there were also a few notable differences. Hypochondriacal reaction was included in Medical 203/VA/Joint Forces but excluded from the DSM-I. Schizophrenic reaction, schizo-affective type and schizophrenic reaction, childhood type were in the DSM-I but did not appear in Medical 203/VA/Joint Forces.

Brief note should also be made of the ICD-6, published in 1948 (World Health Organization, 1948). U.S. psychiatrists played a key role in shaping proposals for the ICD-6, thus accounting for the similarities between the ICD-6 and the DSM-I (Frohlich 1961, p.87; Kendell, 1975, p.92). Although there are some differences between DSM-I and the ICD-6, attempts were made to ensure that DSM-I codes could be converted into ICD equivalents (as outlined in Appendix A of the DSM-I). The direct influence of the ICD-6 on the DSM-I appears to have been minimal. Schizophrenic reaction, schizo-affective type is the only ICD-6 category included in the DSM-I that cannot be found in earlier U.S. classifications (either the military classifications or the *Statistical Manual for the use of Hospitals for Mental Diseases*).

2. The DSM-I was influential

As noted in the foreword of the DSM-I, one of the innovative steps in creating this classification was that the DSM-I committee distributed advance copies to 10% of the members of the APA and solicited feedback in the form of questionnaires. Of the returns, 93% expressed general approval of the new classification. Raines noted that a number of the state hospitals and mental health clinics held staff meetings to discuss the new system and its changes. The efforts by Raines and his committee to pre-test the DSM-I probably helped this

classification in achieving its goal of becoming accepted by the clinicians who were its audience.

During the 1950s, the *American Journal of Psychiatry* printed an annual note on the accounts of the APA. These recorded sales of the DSM-I as being \$1,465 to March 1952, \$10,305 from April 1952 to March 1953, and \$9,825 from April 1953-March 1954. From 1955 and 1965 no sales figures were recorded (APA 1952b, 1953, 1954). The DSM-I thus made at least \$21,595. The DSM-I sold for \$1.50 (Felix, 1956, p.405). This implies that the DSM-I sold over 14,000 copies in its first three years of publication. This equate to roughly two copies for each member of the APA (in 1952 the APA had 7105 members (APA, 1952, p.210)). The DSM-I classification was also reprinted in a number of textbooks (Masserman, 1955; McCartney, 1956; Menninger, 1952) and was used as the organizational basis for popular undergraduate textbooks, on abnormal psychology (Coleman, 1956), and psychiatry (Freedman and Kaplan, 1967).

The DSM-I was used for a variety of purposes. A year after publication, one half of public mental hospitals, one third of private mental hospitals, and three-fourths of general hospitals were using the new nomenclature for patient records (Raines, 1953b). The annual statistical surveys, *Patients in mental institutions*, employed DSM-I categories (Public Health Service, 1958). Outpatient clinics likewise came to use the DSM-I; in 1956, 39% of State and VA outpatient clinics were reporting statistical data that included a DSM-I diagnosis (Bahn and Norman, 1959). Researchers also used the nomenclature of DSM-I categories (influential examples include Pasamanick et al., 1957; O'Neal & Robins, 1958; Greenblatt et al., 1964). Stengel wrote an influential review of international psychiatric classifications for the World Health Organisation, who were concerned by the relative lack of impact of the ICD-6 on the international

psychiatric community (Stengel, 1959, p.605). In this review Stengel notes that “some other countries of the Western Hemisphere” had been considering adopting the DSM-I, and dedicates a special chapter to the classification “in view of its special importance” (Stengel, 1959, p.605). Gruenberg's summary comment in his “Foreward” to the DSM-II appears to have been an accurate assessment, “The first edition of this Manual (1952) made an important contribution to U.S. and, indeed, world psychiatry. It was reprinted twenty times through 1967 and distributed widely in the U.S. and other countries.” (APA, 1968, p.ix).

In summary, the DSM-I sold in large numbers and circulated widely. This being said, its use was by no means universal. When researchers found the DSM-I unsuitable for their purposes, they could ignore it and define the categories they studied descriptively in their papers. Some textbooks employed non-DSM diagnoses (for example, Arieti, 1959). State hospitals in New York State never moved to the DSM-I system, but moved straight from using the 1942 edition of the Statistical Manual to the DSM-II (Spitzer and Wilson, 1968). Although the DSM-I was available to psychiatrists to use as a common reference point, it could be ignored whenever a different approach seemed preferable.

3. Conclusion

The generally accepted view of the DSM-I is that this system was psychoanalytic. This view is incorrect. The roots of the DSM-I came from (1) World War II military classifications that did contain psychoanalytic concepts, and from (2) earlier classifications by the APA that were Kraepelinian in focus and intended primarily for state hospital inpatient settings.

Although U.S. psychiatry in the 1950s is often characterized as psychoanalytic in orientation

it is better described as eclectic. Articles in the *American Journal of Psychiatry* during this period were as likely to discuss somatic therapies, such as psychosurgery and ECT, as they were to discuss psychotherapy. Individual psychiatrists often combined ideas from a variety of schools in their own thinking. The DSM-I reflected this eclecticism and was a common-sense compromise among the schools of thought in American psychiatry at the end of World War II. The DSM-I sold well and circulated widely. The current tendency to dismiss the DSM-I as unimportant is an error.

Acknowledgements

We are grateful to the following individuals, who read drafts and made many helpful comments: Sam Fellowes, A. Kenneth Fuller, John Gunderson, Allan Horwitz, Arthur Houts, Jared Keeley, Martyn Pickersgill and Geoffrey Reed.

References

American Medico-Psychological Association (1918). *Statistical Manual for the Use of Institutions for the Insane*. National Committee for Mental Hygiene: New York.

American Psychiatric Association (1935). Proceedings ninety-first annual meeting. *American Journal of Psychiatry* **92**,413-486.

American Psychiatric Association (1942). *Statistical Manual for the Use of Hospitals for Mental Diseases*. State Hospitals Press: Utica.

American Psychiatric Association (1946). Reports of committees. *American Journal of Psychiatry* **103**, 249 – 269.

American Psychiatric Association (1949a). Proceedings of the one hundred and fourth annual meeting. *American Journal of Psychiatry* **105**, 851-868.

American Psychiatric Association (1949b). Committee and other reports. *American Journal of Psychiatry* **105**, 920 – 935.

American Psychiatric Association (1950). *Biographical Directory of Fellows and Members of the American Psychiatric Association*. American Psychiatric Association: New York.

American Psychiatric Association (1952). *Diagnostic and Statistical Manual: Mental Disorders*. American Psychiatric Association: Washington, DC.

American Psychiatric Association (1952b). Proceedings of the American Psychiatric Association: The one hundred and eighth annual meeting. *American Journal of Psychiatry* **109**, 210-226.

American Psychiatric Association (1953). Proceedings of the American Psychiatric Association: The one hundred and ninth annual meeting. *American Journal of Psychiatry* **110**, 209-229.

American Psychiatric Association (1954). Proceedings of the American Psychiatric Association: The one hundred and tenth annual meeting. *American Journal of Psychiatry* **111**, 207-228.

American Psychiatric Association (1968). *Diagnostic and Statistical Manual of Mental Disorders. Second Edition*. American Psychiatric Association: Washington.

Anonymous (1946). Jacob Sergei Kasanin. *Journal of Nervous and Mental Disease* **104**, 112-113.

Aragona M (2015). Rethinking received views on the history of psychiatric nosology: minor shifts, major continuities. In *Alternative Perspectives on Psychiatric Validation*. (ed. P. Zachar, D. St. Stoyanov, M. Aragona, and A. Jablensky), pp27-46. Oxford University Press: Oxford.

Arieti S (1959) *American Handbook of Psychiatry*. Basic Books: Oxford.

Army, U. S., Navy, U. S., & Air Force, U. S. (1949) *Joint Armed Forces Nomenclature and Method of Recording Psychiatric Conditions*. NavMed P-1303. US Government Printing Office: Washington, DC.

Bahn AK and Norman VB (1959). First national report on patients of mental health clinics. *Public Health Reports* **74**, 943-956.

Blinder MG (1966). The pragmatic classification of depression. *American Journal of Psychiatry* **123**, 259-269.

Bowman KM and Rose M (1951). A criticism of the terms “psychosis”, “psychoneurosis” and “neurosis”. *American Journal of Psychiatry* **108**,161-166

Coleman JC (1956). *Abnormal Psychology and Modern Life*. 2nd edition Scott, Foresman & Co.: Chicago.

Dayton NA (1935). The first year of the new standard nomenclature of diseases in Massachusetts mental hospitals. *American Journal of Psychiatry* **92**, 589-613.

Decker HS (2013). *The Making of DSM-III: A Diagnostic Manual's Conquest of American Psychiatry*. Oxford University Press: Oxford.

Felix R (1956). A bookshelf on mental health. *American Journal of Public Health and the Nation's Health* **46**, 397-407.

Freedman A and Kaplan H (1967). *Comprehensive Textbook of Psychiatry*.

The Williams & Wilkins Company: Baltimore.

Frohlich M (1961). Discussion. In *Field Studies in the Mental Disorders* (ed J. Zubin), pp.

85-88. Grune & Stratton: New York.

Gottlieb JS (1952). The responsibility of public mental hospitals in psychiatric research.

Psychiatric Services, **3(9)**, 6-10.

Greenblatt M, Grosser GH, Wechsler H (1964). Differential response of hospitalized

depressed patients to somatic therapy. *American Journal of Psychiatry* **120**, 935-943.

Grob GN (1991) Origins in DSM-I: A study in appearance and reality.

American Journal of Psychiatry **148**, 421-431.

Horwitz A (2015). DSM-I and DSM-II. In *The International Encyclopedia of*

Clinical Psychology (ed. RL Cautin and SO Lilienfeld), pp.1-6. John Wiley: New

York.

Houts AC (2000). Fifty years of psychiatric nomenclature: reflections on the 1943 war

department technical bulletin, Medical 203. *Journal of Clinical Psychology* **56**, 935-967.

Kahn E (1955). A marginal note on interpretation. *American Journal of Psychiatry* **112**, 393-395.

Kendell R (1975). *The Role of Diagnosis in Psychiatry*. Blackwell: Oxford.

Kramer M and Pollack E S (1958). Problems in the interpretation of trends in the population movement of the public mental hospitals. *American Journal of Public Health and the Nation's Health*, **48**, 1003-1019.

Masserman J (1955). *The Practice of Dynamic Psychiatry*. Saunders: Philadelphia.

Maxmen JS (1985). *The New Psychiatry*. William Morrow & Company: New York.

May JV (1922). *Mental Diseases: A Public Health Problem*. Richard G. Badger: Boston.

McCartney J (1956). *Understanding Human Behaviour*. Vantage: New York.

Menninger WC (1947). Psychiatric experience in the war, 1941-1946. *American Journal of Psychiatry* **103**, 577-586.

Menninger KA (1952). *A Manual for Case Study*. Grune and Stratton: New York.

Menninger KA (1963). *The Vital Balance*. Viking Press: New York.

National Institute of Mental Health (1964). *Patients in Mental Institutions, 1963, Part III State and County Mental Hospitals*. Public Health Services Publication: Washington DC.

Office of the Surgeon General, Armed Service Forces (2000 [1946]). Nomenclature of psychiatric disorders and reactions. *Journal of Clinical Psychology* **56**, 925-934. Originally published as War department technical bulletin, Medical 203. *Journal of Clinical Psychology* **2**, 289-296.

O'Neal P, Robins, LN (1958). The relation of childhood behavior problems to adult psychiatric status: A 30-year follow-up study of 150 subjects. *American Journal of Psychiatry* **114**, 961-969.

Pasamanick B, Roberts DW, Lemkau PV, Krueger, DE (1957). A survey of mental disease in an urban population. I. Prevalence by age, sex, and severity of impairment. *American Journal of Public Health and the Nation's Health* **47**, 923-929.

Pollock HM (1945). Development of statistics of mental disease in the United States during the past century. *American Journal of Psychiatry* **102**, 1-17.

Public Health Service (1958) *Patients in Mental Institutions 1955*. US Public Health Service,

Raines GN (1953a). Discussion of Dr Sandor Rado's academic lecture. *American Journal of Psychiatry* **110**, 425-426.

Raines GN (1953b) Comment: the new nomenclature. *American Journal of Psychiatry* **109**, 548-9.

Shorter E (2015). *What Psychiatry Left Out of the DSM-5: Historical Mental Disorders Today*. Routledge: New York.

Spitzer RL, Wilson PT (1968). An introduction to the American Psychiatric Association's new diagnostic nomenclature for New York State Department of Mental Hygiene Personnel. *Psychiatric Quarterly* **42**, 487-503.

Stengel E (1959). Classification of mental disorders. *Bulletin of the World Health Organization* **21**, 603-663.

Veterans Administration (1947). *Nomenclature of Psychiatric Disorders and Reactions* *Veterans Administration Technical Bulletin, TB 10A-78*. Oct 1 1947. Washington 25, DC.

Weinstock HI (1965). The role of classification in psychoanalytic practice. In *The Role and Methodology of Classification in Psychiatry and Psychopathology*. (ed. MM Katz, JO Cole, WE Barton), pp.62-72. U.S Dept of Health, Education and Welfare.

Whitaker R (2011). *Anatomy of an Epidemic*. Broadway: New York.

