1	End of the Kiaman Superchron in the Permian of SW England:
2	Magnetostratigraphy of the Aylesbeare Mudstone and Exeter
3	groups.
4	Mark W. Hounslow* <sup>1</sup> , Gregg McIntosh <sup>2</sup> , Richard A. Edwards <sup>3</sup> , Deryck J. C. Laming <sup>4</sup> ,
5	Vassil Karloukovski <sup>1</sup>
6 7 8 9 10	<ul> <li><sup>1</sup>Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YW, U.K.</li> <li><sup>2</sup> School of Human and Life Science, Canterbury Christchurch University, Canterbury, UK.</li> <li><sup>3</sup>Hawkridge, Thorverton, Devon, EX5 5JL, U.K (formerly British Geological Survey, Exeter, UK)</li> <li><sup>4</sup>Herrington Geoscience and David Roche GeoConsulting, Renslade House, Bonhay Road, Exeter EX4 3AY, Devon, UK.</li> </ul>
12	*Corresponding author (e-mail: <u>m.hounslow@lancaster.ac.uk</u> :Tel:+ 44 (0) 1524 510238
13	Number of: words of text (=8223), words in captions (=1140), references (=103), Tables (=1)
14	and figures (=11)
15	Abbreviated title (< 40 characters) : Kiaman Superchron, SW England
16	Keywords: Magnetostratigraphy, Permian, Triassic, Kiaman Superchron, Rotliegend
17	
18	Abstract: The chronology of Permian strata in SW England is fragmentary and largely based on
19	radiometric dating of associated volcanic units. Magnetostratigraphy from the ~2 km of
20	sediments in the Exeter and Aylesbeare Mudstone groups was undertaken to define a detailed
21	chronology, using the end of the Kiaman superchron, and the overlying reverse and normal
22	polarity in the Middle and Upper Permian as age constraints. The palaeomagnetic directions are
23	consistent with other European Permian palaeopoles, with data passing fold and reversal tests.
24	The end of the Kiaman superchron (in the Wordian) occurs in the uppermost part of the Exeter
25	Group. The overlying Aylesbeare Mudstone Group is early Capitanian to latest Wuchiapingian in
26	age. The Changhsingian and most of the Lower Triassic is absent. Magnetostratigraphic
27	comparison with the Southern Permian Basin shows that the Exeter and Aylesbeare Mudstone
28	groups are closely comparable in age to the Havel and Elbe Subgroups of the Rotleigend II
29	succession. The Altmark unconformities in these successions appear similar in age as the
30	sequence boundaries in SW England, indicating both may be climate controlled. Clasts in the
31	Exeter Group, from unroofing of the Dartmoor granite, first occurred at a minimum of ~8 Ma
32	after formation of the granite.

- 34 **Supplementary material:** Additional magnetic fabric and palaeomagnetic data is available at:
- 35 hhtp://www.geolsoc.org.uk/SUP000

37 Permian and Triassic successions in southern and SW England were produced following the 38 Variscan orogeny and occur in a number of interconnected, sag and fault-bounded basins, the 39 largest being the Wessex Basin, and various sub-basins that form the Channel Approaches Basin. 40 Some basins contain up to 8 km of post-Variscan red-bed fill (Harvey et al. 1994; Hamblin et al. 41 1992; Butler 1998; McKie & Williams 2009; Fig. 1). The Wessex Basin formed on Rheno-42 Hercynian basement (Variscan), between the Northern Variscan Front and the Lizard-Rhenish 43 Suture. The sub-basins of the Western Approaches Basin formed on Saxo-Thuringian (Variscan) 44 and Rheno-Hercynian basement (McCann et al. 2006; Strachan et al. 2014). As such, these 45 basins may share similar tectonic and stratigraphic histories with similarly situated basins in 46 France and Germany such the Saar-Nahe and Saale basins in Germany (Roscher & Schneider 47 2006; McCann et al. 2006.). However, our tectono-stratigraphic understanding of the UK basins 48 is poorly integrated into the framework of European Permian basin evolution. These 49 intramontane basins often lack the distinctive late Permian carbonate-evaporite, Zechstein 50 successions, common in basins (e.g. Southern Permian Basin) north of the Variscan front, and 51 lack the early Permian faunas of the southern Variscan basins (Roscher & Schneider 2006; 52 McCann et al. 2006).

53

54 The onshore Permian-Triassic successions in the western parts of the Wessex Basin and the 55 Credition Trough, outcrop as the Exeter, Aylesbeare Mudstone and Sherwood Sandstone groups 56 (Figs. 1 & 2). The coastal outcrops form part of the Jurassic Coast World Heritage Site (Barton et 57 al. 2011). The work of the British Geological Survey, related to the re-mapping of the Exeter 58 area (Edwards et al. 1997), generated a better regional understanding of the Permian Exeter 59 Group (Gp). The oldest successions outcropping in the Crediton Trough (and Torbay area) may 60 extend into the latest Carboniferous (Edwards et al. 1997; Leveridge et al. 2003). The units 61 below the base of the Whipton Formation (Fm) in the Exeter and Crediton Trough area contain a 62 variety of basaltic and lamprophyric lavas and intrusions, whose Ar-Ar and K-Ar ages (291-282 Ma) are older than the more tightly constrained Rb-Sr, U-Pb and Ar-Ar ages (at 280 Ma) for the 63 64 formation of the Dartmoor Granite (Scrivener 2006). These volcanic and igneous units are coeval 65 with widespread volcanic activity throughout Europe during the latest Carboniferous to early 66 Permian (Timmerman 2004). The isostatic uplift and regional denudation coeval with, and 67 following, Dartmoor Granite emplacement, was probably responsible for a major unconformity 68 (Edwards et al. 1997) separating the Whipton Fm from the older units (Fig. 2).

70 Miospores from the Whipton Fm around Exeter, and younger units equivalent to the Alphington 71 and Heavitree Breccia formations, demonstrate similarities to assemblages from the Russian 72 Kazanian and Tatarian regional stages (Warrington & Scrivener 1990; Edwards et al. 1999). 73 Consequently, the barren overlying Aylesbeare Mudstone Gp has been assigned to the Lower 74 Triassic in some subsequent studies (Newell 2001; Benton et al. 2002). Since the Aylesbeare 75 Mudstone Gp is widespread in the Wessex Basin and the western approaches (Hamblin et al. 76 1992; Butler 1998; Evans 1990; Barton et al. 2011), a Lower Triassic mudstone-dominated 77 lacustrine unit creates a major palaeogeographic problem. That is, southerly-derived clasts in the 78 Lower Triassic units, in central and Northern Britain, could not have been sourced through the 79 Wessex Basin, from the Armorican supply areas to the south, as has been widely concluded for 80 over 100 years (Ussher 1876; Thomas 1909; Wills 1970; McKie & Williams 2009; Morton et al. 81 2013).

82

83 To resolve this problem, and constrain in detail the age of the Permian successions, we use 84 magnetostratigraphy as a dating tool. The Kiaman (reverse polarity) superchron (KRPS) extends 85 from the mid Carboniferous to the mid Permian, but had ended by the early Wordian (mid 86 Guadalupian), after which reverse and normal polarity intervals (here called the Illawarra 87 Superchron) occur during the remainder of the mid and late Permian, extending into the Triassic 88 (Steiner 2006; Hounslow submitted). We demonstrate for the first time in the UK, the 89 stratigraphic position of the end of the KRPS, and also the polarity pattern through the upper part 90 of these successions (below the Budleigh Salterton Pebble Beds Fm). Our data allows a new 91 understanding of the precise age of these units, which suggests a new relationship to the better-92 studied successions in the Southern Permian Basin.

#### 93 Geology and Lithostratigraphy

Excellent exposures of the Exeter and Aylesbeare Mudstone groups occur in a series of cliff and foreshore exposures between Torbay and Budleigh Salterton (Fig. 1). These sediments have maximum burial temperatures of ca. 80±5°C attained during the early Cretaceous, with estimated maximum burial depths of 2 to 2.5 km (Carter *et al.* 1995). Faulting and folding structures in the Wessex Basin are a response to basin inversion, produced by N-S compression, along mostly E-W trending faults, many of which were former extensional structures. The basin inversion took place during the late Cretaceous and early Tertiary (Underhill & Stoneley 1998).

#### 102 Exeter Group

103 The Exeter Group is predominantly the deposits of a number of alluvial fans, with aeolian dune 104 sandstones dominating in the Dawlish Sandstone Fm, and in some units in the Torbay Breccia 105 Fm (Fig. 2). The coastal successions in Torbay are separated from those north of Oddicombe 106 (Fig. 1) by the Torquay-Babbacombe promontory which was a palaeogeographic feature in the 107 Permian (Laming 1966). Mapping work indicates the Torbay Breccia Fm (Leveridge et al. 2003), 108 can be divided into a number of separate breccias units with differing clast contents (unpublished 109 work of DJCL). The Watcombe Fm which is an on-lapping mudstone-rich breccia unit, which 110 north of the Torquay-Babbacombe promontory is unconformably overlain by the Oddicombe 111 Breccia Fm (and the equivalent Paignton breccias south of this). On the coastal outcrops the 112 Watcombe Fm has a  $9-20^{\circ}$  dip-discordance with the overlying Oddicombe Breccia (9° at 113 Whitsand Bay and 20° at Oddicombe Cove; Figs. 1, 3d; Laming 1982). The lower parts of the 114 Torbay Breccia Fm (Roundham Head breccias, with clasts derived from SW) are generally poor 115 in volcanic clasts (Laming 1982) like the oldest unit (the Cadbury Breccia Fm; Edwards & Scrivener 1999) in the Crediton Trough, and by inference may have similar ages, prior to the 116 117 early Permian basaltic volcanism.

118

119 The various breccia units below the Dawlish Sandstone Fm are largely distinguished by their 120 clast contents. These contain a variety of lithologies (limestone, sandstone, vein quartz, quartzite 121 and slate) from various Variscan basement units, together with a variety of volcanic rock 122 fragments associated with the granite and its former or earlier extrusives (Laming 1982; Selwood 123 et al. 1984; Edwards & Scrivener 1999). The Watcombe and Whipton formations consist of fine-124 grained sandy or muddy breccia with clasts of slate and sandstone with occasional porphyry. 125 They contain irregularly interbedded sandstone and mudstone units (Ussher 1913), which 126 dominate the Whipton Fm around Exeter (Edwards & Scrivener 1999). The Oddicombe Breccia 127 Fm (Fig. 2) is rich in locally derived limestone fragments, which typically displays fining-up 128 sequences (into poorly sorted sandstones or fine-breccias; Benton et al. 2002) several metres 129 thick, well displayed at Maidencombe Cove and Bundle Head (Figs. 1, 4). The Alphington 130 Breccia Fm is likewise rich in locally derived shale and sandstone fragments, and hornfelised 131 shale from the underlying Variscan basement (Edwards et al. 1997). The Teignmouth and 132 Heavitree formations are distinctive in containing common clasts of pink and white perthitic 133 feldspar (murchisonite), which Dangerfield & Hawkes (1969) interpreted as feldspar megacrysts 134 from the roof zone of the Dartmoor granite; the supply of which probably indicates synchronous

- unroofing into adjacent alluvial fan successions. The older Alphington and Oddicombe Breccia
  formations lack the murchisonite clasts (Selwood *et al.* 1984; Edwards & Scrivener 1999).
- 137

138 All the breccia units tend to be poorly sorted, and may locally contain a high proportion of mud 139 or sand. The fining-up successions in the Teignmouth Breccia Fm tend to be smaller scale (< 1 m 140 thickness) and typically display poor lateral organisation. Breccias in the upper-part of this 141 formation have interbedded aeolian sandstone units, well displayed in the Coryton Cove area (8 142 on Fig. 1; Fig. 4); which is a transitional part of this formation into the overlying Dawlish 143 Sandstone Fm. The estimated thicknesses of the Oddicombe and Teignmouth Breccia formations 144 vary widely between different authors, because of faulting, variable bedding dips and probably 145 significant palaeotopography on the Variscan basement. The thicknesses of Selwood et al. (1984) 146 are minimum thickness estimates, whereas Laming (1969; 1982) and this work suggest greater thicknesses at the upper limits indicated in Fig. 2. 147

148

149 The aeolian dune systems that dominate deposition in the Dawlish Sandstone Fm (Newell 2001), 150 also display interbedded fluvial sandstone and breccia units. Around Exeter and further north in 151 the Crediton Trough, the Dawlish Sandstone Fm onlaps onto older units, to rest on Variscan 152 basement. The overlying Exe Breccia Fm is divisible into a lower porphyry-bearing unit (the 153 Kenton Mbr), typical of most of the outcrop on the west of the Exe Estuary, and an overlying 154 quartzite- and mudstone-bearing breccia (the Langstone Mbr). This upper member is well 155 exposed at Langstone Rock (6 on Fig. 1) which in the upper part is dominated by poorly sorted 156 sandstones and sandy siltstones (Gallois 2014; Fig. 4). The thickness of the Exe Breccia is 157 uncertain, due to faulting along the Exe Estuary; 85 m was suggested by Selwood et al. (1984), 158 but up to ~50 m is more likely (Laming & Roche 2013). The uppermost part of the Langstone 159 Mbr at Lympstone and Sowden Lane (3 on Fig. 1) displays both well-developed shallow fluvial 160 channels and aeolian sandstone units, and is gradational into the mudstones and siltstones 161 forming the base of the Aylesbeare Mudstone Gp (Gallois 2014; Fig. 4). Around Exeter and in 162 the Crediton Trough the Aylesbeare Mudstone Gp is unconformable on the Dawlish Sandstone 163 Fm, also onlapping onto older units (Edwards et al. 1997; Edwards & Scrivener 1999).

#### 164 Aylesbeare Mudstone Group

165 The Exmouth Mudstone and Sandstone Fm is a lacustrine, red-brown mudstone-dominated unit,

166 with interbedded fine to medium-grained fluvial and lacustrine sandstone units (thicker beds

167 labelled as Beds A to J by Selwood et al. 1984). These are most prominent towards the upper 168 part of the formation, where the term Straight Point Sandstone Mbr is introduced for these 169 persistent sandstone beds (i.e. beds I and J of Selwood et al. 1984). This member is mapped 170 between the coast and Aylesbeare, north of which the Aylesbeare Mudstone Gp is not sub-171 divided (Edwards & Scrivener 1999). The upper few metres of the Straight Point Sandstone Mbr 172 at outcrop has patchily developed, immature, nodular and sheet-like groundwater calcretes, 173 locally with rhizoconcretions (Fig. 3B). The base of the overlying Littleham Mudstone Fm is 174 taken at the base of the porphyry and murchisonite bearing breccia unit (Ormerod-Wareing 175 1875), which locally erosively overlies this calcrete-bearing sandstone (Fig. 3C), and grades into 176 overlying interbedded sandstone, siltstone and mudstone beds in the basal parts of the Littleham 177 Mudstone Fm, west of the Littleham Cove fault (Fig. 5).

178

179 The Littleham Mudstone Fm is well-exposed in the cliffs between Littleham Cove and Budleigh 180 Salterton, but is locally disrupted by faulting in the lower part and landslips in the cliff. The 181 complete succession in the cliffs was determined by using a montage of photographs taken from 182 offshore, which shows the full succession is divided by a number of prominent green mudstone, 183 thin sandstone and siltstone beds (Fig. 5). The succession in the cliffs can be divided into three 184 units, a lower unit (Division A) east of the Littleham Cove fault with a few green mudstone beds, 185 a middle unit (Division B) with relatively common sandstone and siltstone beds, and an upper 186 unit (Division C) with more frequent green mudstone beds and some impersistant sandstones. 187 The true thickness of the Littleham Mudstone Fm, in these outcrops, cannot be determined 188 because of the uncertain displacement on the Littleham Cove fault. However, the measured 189 cumulative thickness east and west of the fault (216 m), is similar to the  $\sim$ 205 m and 230 m 190 measured in the Blackhill and Withycombe Rayleigh boreholes respectively (Bateson & Johnson 191 1992; Fig. 1), so the cliff outcrops probably represent most of the Littleham Mudstone Fm. In the 192 Venn Ottery borehole (Fig. 1) the Littleham Mudstone Fm contains pods and veins of gypsum, 193 and thin interbedded aeolian sandstones (Bateson & Johnson 1992; Edwards & Scrivener 1999; 194 N.S Jones pers comm to RAE). A substantial unconformity separates the Littleham Mudstone Fm 195 from the overlying Budleigh Salterton Pebble Beds Fm, shown by the dramatic lithology change, 196 and the sharp and irregular boundary (Fig. 3A) with some authors suggesting a small bedding dip 197 difference (Irving, 1888). Gallois (2014) has suggested this contact is conformable.

#### 198 Regional relationships

199 Broadly the Permian units in the study area can be divided into 5 genetic sequences (Pm1 to 200 Pm5), bounded by a hiatus or unconformity (Fig. 2). The upper three of these are all 201 characterised by basal breccias units (low stand deposits), with conformable transitions into 202 overlying finer-grained upper parts. The relationships of the successions in Torbay to those in the 203 Crediton Trough area is less certain. It is probable that the earliest parts of the Torbay Breccia 204 Fm is timing-related to the Cadbury Breccia Fm in the Crediton Trough (sequence Pm1), since 205 both units are very poor in igneous clasts (Edwards et al. 1997). These five sequences may relate 206 to the four sequences (A to D) seen in the Plymouth Bay Basin (Harvey et al. 1994). Their oldest 207 megasequence A likely relates to Pm1 and megasequence B to Pm2, since it is capped by an 208 inferred volcanic unit. Megasequence C likely relates to Pm3, and is marked by a change in 209 orientation of the Plymouth Bay Basin depocentres. Divergent bedding dips between units under 210 and overlying the Watcombe Fm (Pm2), suggest that the most important extensional event 211 (Leveridge et al. 2003; Laming 1982) is at the Pm2-Pm3 boundary, consistent with depocentre 212 orientation change in the Plymouth bay basin. Megasequence D is probably equivalent to Pm4 213 and Pm5, since the Pm4-Pm5 boundary is subtle to detect in the field.

214

215 The continuity of these units to the east in the central parts of the Wessex Basin is uncertain. 216 Henson (1972) suggested, based on geophysics, that the breccia units thin to the east, so 217 eastwards the breccias may pass into the mudstone dominated units, equated with the Aylesbeare 218 Mudstone Gp in the central parts of the Wessex Basin, which are up to  $\sim 1.5$  km thick (Butler 219 1998; Hamblin et al. 1992). However, Henson's data failed to detect the faults, along the Exe 220 Estuary, so the interpretation may be flawed. In the Western Approaches basins 1 km or more of 221 anhydritic mudstones and sandstones underlie the equivalent of the Sherwood Sandstone Gp 222 (Evans 1990). These locally rest on a Permian volcanic sequence, presumably of a similar age to 223 the early Permian Exeter Volcanic Rocks (Chapman 1989; Fig. 2).

#### 224 Palaeomagnetic sampling

225 Almost the entire succession of the Aylesbeare Mudstone Gp is exposed in the sea-cliffs between

- 226 Budleigh Salterton and Exmouth. Only the mid and upper parts of the Exe Breccia could be
- sampled at Lympstone (location 3 on Fig.1) and Langstone Rock (6 on Fig. 1; see Supplementary
- data for details). Outcrops in the lower parts of the Exe Breccia Fm (Kenton Mbr), were all too
- 229 coarse-grained for palaeomagnetic sampling. Most of the Dawlish Sandstone and Teignmouth

- 230 Breccia are well exposed between Langstone Rock and Teignmouth, adjacent to the main
- 231 London-Penzance railway-line (Ussher 1913; Selwood et al. 1984), but large parts are
- 232 inaccessible due to rail-safety restrictions. The Dawlish Sandstone Fm was sampled in quarries
- 233 near Exeter (4 and 5 on Fig. 1; Fig. 4). The uppermost part of the Teignmouth Breccia was
- available for sampling in the Coryton Cove and Dawlish Station sections (7 and 8 on Figs. 1, 4).
- 235 Reconnaissance sampling of the Oddicombe and Watcombe Breccias was undertaken. For the
- 236 most part, these units are fully exposed in sea-cliffs and foreshore exposes between Teignmouth
- and Oddicombe (Fig. 4). The Knowle Sandstone Fm was sampled at west Sandford (Edwards *et*
- 238

al. 1997).

239

240 Samples from these units were collected using mostly hand samples, oriented with a compass. In

total some 153 samples were collected from 13 sites (see Supplementary data), largely focussed

on reddened lithologies. Cubic specimens were cut from the hand samples using a circular saw.

243 Some samples from sandstone units in the Dawlish Sandstone and Exe and Teignmouth Breccias

were poorly consolidated, and impregnated in the laboratory with a 2:1 mix of sodium silicate

and water (Kostadinova *et al.* 2004) to consolidate them prior specimen preparation.

#### 246 Laboratory Methodology

247 Measurements of Natural Remanent Magnetisation (NRM) were made using a CCL 3-axis 248 cryogenic magnetometer (noise level ~0.002 mA/m), using multiple specimen positions, from 249 which the magnetisation variance was determined. The magnetometer is not housed in a 250 controlled space which cancels the earth's magnetic field. Instead specimens were housed in Mu-251 metal boxes with an ambient magnetic field <10 nT at all times, other than when being measured 252 or demagnetised. Generally, 1 to 3 specimens from each sample were treated to stepwise thermal 253 demagnetisation, using a Magnetic Measurements Ltd thermal demagnetiser, in 50-40°C steps up 254 to 700°C. Low frequency magnetic susceptibility ( $K_{\rm lf}$ ) was monitored after heating stages, 255 measured using a Bartington MS2B sensor. Specimens from the Bishops Court Quarry gave poor 256 quality results and sister specimens were partly treated with a combination of thermal and 257 alternating field (AF) demagnetisation, the latter conducted using a Molspin tumbling AF 258 demagnetiser. In total 166 and 78 paleomagnetic specimens were demagnetised from the Aylesbeare Mudstone and Exeter groups respectively. The bedding dips in the Aylesbeare 259 260 Mudstone Gp are  $5-10^{\circ}$  in an easterly direction, so a fold test was not possible. However, in the

261 Exeter Gp dips are more variable and up to  $40^{\circ}$ , so a fold-test was possible.

262

- 263 Characteristic remanent magnetisation (ChRM) directions were isolated using principal
- 264 component-based statistical procedures as implemented in LINEFIND, which uses the
- 265 measurement variance along with rigorous statistical procedures for identifying linear and planar
- structure in the demagnetisation data (Kent *et al.* 1983). Both linear trajectory fits and great
- 267 circle (remagnetisation circle) data were used in defining the paleomagnetic behaviour, guided by
- 268 objective and qualitative selection of the excess standard deviation parameter ( $\rho$ ), which governs
- 269 how closely the model variance, used for analysis, matches the data measurement variance (Kent
- 270 et al. 1983). The PMAGTOOL software (available at
- 271 https://www.lancs.ac.uk/staff/hounslow/default.htm) was used for the analysis of mean directions
- and virtual geomagnetic poles.
- 273

274 Progressive isothermal remanent magnetisation (IRM) up to 1T is most and 4 T is some was

- applied to a representative sub-set of specimens, to investigate the magnetic mineralogy. Back
- field demagnetisation was also used on some specimens. This used an ASC Scientific IM-30
- 277 impulse magnetiser and a Molspin pulse magnetiser. The IRM was measured using a Molspin
- 278 spinner magnetometer. Thermal demagnetisation of a three component IRM was used to
- 279 investigate the unblocking temperatures (Lowrie 1990). A small set of specimens were measured
- 280 for magnetic hysteresis (maximum field 0.9 T) and thermomagnetic curves (maximum field 300
- 281 mT, in air using a Magnetic measurements variable field translation balance, MMVFTB).
- 282 Selected thin sections were investigated to assess the petrography of the Fe-oxides. The
- anisotropy of magnetic susceptibility (AMS), of selected specimens, was measured using an
- Agico KLY3S Kappameter, to assess the preservation of the detrital sedimentary fabric (Løvlie
- 285 & Torsvik 1984; Tarling & Hrouda 1993), and to assess if fabrics had been modified by the weak
- tectonism.

### 287 Magnetic Mineralogy

- 288 Changes in the NRM intensity and K<sub>lf</sub> of specimens are broadly related to:
- a) The amount of silt and clay, with those samples having larger amounts of silt and clay
- 290 generally having larger NRM intensity and K<sub>lf</sub>. For example, aeolian sandstones such as
- those in the Dawlish Sandstone Fm, have significantly lower NRM intensity and K<sub>lf</sub> (Fig. 5,
- see Supplementary data). In the Aylesbeare Mudstone Gp red mudstones possess average

293		NRM intensity and $K_{\rm lf}$ of 5.0 mA/m and 20.0 x10 <sup>-6</sup> SI respectively, compared to means of 1.8
294		mA/m and $7.2 \times 10^{-6}$ SI in the red sandstone beds.
295	b)	Reddened and non-reddened samples of the same lithology often possesses dramatically
296		different NRM intensity and $K_{lf}$ ; with the non-reddened samples typically having lower
297		values. For example grey, green and white sandstones in the Aylesbeare Mudstone Gp have
298		mean NRM intensity and $K_{if}$ of 0.9 mA/m and 4.4 x10 <sup>-6</sup> SI respectively.
299	c)	The average NRM intensity and $K_{lf}$ shows progressively larger values into the Oddicombe
300		Breccia and Watcombe formations (see supplementary data). This may relate to a progressive
301		increase in volcanic-derived detritus (and iron oxide content) in the older units, which is
302		shown by the Cs content (Merefield et al. 1981).
303		
304	Spe	ecimens analysed do not saturate in IRM fields up to 4 T (Fig. 6A, C), indicating that canted
305	ant	iferrimagnetic minerals (haematite or goethite) are important magnetic minerals in these
306	spe	cimens. Durrance et al. (1978) also detected haematite as the main Fe-oxide in the Littleham
307	Mu	dstone Fm, with the addition of significant amounts of superparamagnetic haematite.
308	The	ermomagnetic curves were nearly reversible and exhibited Curie temperatures of 657-669°C,
309	and	I thermal demagnetisation of the IRM, shows that specimens display blocking temperatures up
310	to 6	650-700°C (Fig. 6). Coercivity of remanence (Bcr) ranged between 320 and 710 mT, all
311	sug	gesting predominant haematite (Frank & Nowaczyk 2008). Although the IRM does not
312	app	proach saturation by 4 T (Fig. 6B), there is no clear evidence for goethite, since we have high
313	SIF	RM/ K <sub>if</sub> values, and no well-defined Neel temperature for goethite. Percent IRM acquisition
314	bel	ow 100 mT ( $\%$ IRM <sub>100mT</sub> ) is mostly 5%-10% of the 1T IRM, for reddened lithologies. Only in
315	aec	lian sandstone units in the Dawlish Sandstone Fm, and grey or red mottled green/grey
316	lith	ologies does the $\%$ IRM <sub>100mT</sub> rise above 10% to ca. 50% at maximum (Fig. 6A,C,E). Data
317	fro	m synthetic mixtures (Frank & Nowaczyk 2008) suggest such $\%$ IRM <sub>100mT</sub> values indicate a
318	ma	gnetite contribution to the iron oxide load of ca. 0.05% (for red lithologies) to a maximum of
319	ca.	1% for aeolian samples DS16 and DS100 (Fig. 6E). In specimens DS16, (from Dawlish
320	Sar	ndstone Fm aeolian sandstones; Fig. 6F) and L3 (grey sandstone, Littleham Mudstones Fm;
321	Fig	. 6B) the 100 mT IRM demagnetises by 450°C- 550°C, which could suggest an oxidized, or
322	Ti-	poor magnetite (Fig. 6F). The >300 mT coercivity component in specimen DS16 has a
323	blo	cking temperature of ~550°C, probably due to a pigment-dominated haematite remanence
324	(Tu	rrner 1979) in this sample.

326 Petrography indicates, like other red-beds, that the haematite is present as two phases, firstly sub-327 micron haematite (pigmentary haematite), which coats pore perimeters and is often internal to 328 some rock clasts; secondly as larger specular haematite particles, most obvious as detrital opaque 329 grains (Turner 1979; Fig. 3E, F). The pore-lining pigmentary haematite is multiphase in origin, 330 since it both coats feldspar overgrowths, and to a lesser extent, coats the grains prior to the 331 overgrowths (observed in Dawlish Sandstone Fm only). Compaction related pressure solution at 332 some grain contacts, shows greater amounts of pigment coating the pores, and lesser amounts 333 between the grain contacts, demonstrating both pre and post-compaction pigmentary haematite 334 formation, with probably the bulk of the pigment produced post compaction. Some of the 335 pigmentary haematite may have formed pre-deposition, since it is widely dispersed within a 336 variety of siltstone and phyllite clasts.

337

338 The specular haematite is dominated by detrital opaques, which are either present as haematite

339 dominated particles, or compound particles in-part composed of other silicate minerals. The

340 compound particles are occasional haematised clastic rock fragments (intraformational?) but

341 most are of uncertain origin (Fig. 3E). These two types of specularite grains vary in abundance

342 from about 1% to trace amounts. Larger amounts of detrital opaques tend to occur in samples that

343 are finer-grained or less well sorted, and lesser amounts in the well-sorted aeolian sandstones.

### 344 Magnetic Fabric

345 The anisotropy of magnetic susceptibility (AMS) overall shows a primary depositional magnetic

346 fabric, characterised by vertical-to-bedding K<sub>min</sub> directions (Figs. 7 a-d) and largely oblate (T >0)

347 fabrics (Figs. 7 e - h). The mudstones have the stronger AMS (greater P values) and are always

348 oblate. The sandstones within the Aylesbeare Mudstone Gp and the various breccia units show

349 more variable AMS fabrics ranging into the prolate fields (T <0), especially so for some

350 sandstones from the breccia units (Fig. 7e, h). This may relate to the more poorly sorted,

351 probably more chaotically deposited grains in the breccia units (possibly related to mudflow

deposition, cf. Park et al. 2013). K<sub>max</sub> axis trends (Figs. 7i) for specimens from the breccia units

353 (Fig. 7l) show both N-S trends and ENE-WSW trends similar to the clast imbrication directions

- 354 (typically between easterly and northerly directions) of Laming (1982) and Selwood *et al.*
- 355 (1984). This demonstrates the K<sub>max</sub> directions parallel the fluvial transport directions. The N-S
- 356 directed K<sub>max</sub> axes trends are common near the Babbacombe-Torquay promontory and in the
- 357 Teignmouth Breccia Fm. Similar easterly and northeasterly K<sub>max</sub> axes trends are present in the

- Exmouth Sandstones and Mudstones, whereas those in the Littleham Mudstone Fm are morevariable.
- 360

361 Specimens from aeolian sandstones (from the Dawlish Sandstone Fm and upper part of the Teignmouth Breccia Fm) show a larger proportion of prolate fabrics (T <0) with many more  $K_{min}$ 362 363 axes deviating from vertical (Figs. 7c, g). This may be partly caused by the AMS in these 364 specimens being closer to the sensitivity limits of the KLY3S. However, it is also a reflection of 365 the rolling grain transport on the leeward slip-faces of the aeolian dunes (Ellwood & Howard 1981), producing a grain long-axis orientation transverse to the average wind direction 366 (Schwarzacher 1951), which was to the NW to NNW (Laming 1982; Newell 2001). This is 367 368 clearly shown in the specimens from the Bishops Court Quarry in which the K<sub>max</sub> axes are 369 transverse to the aeolian foresets (Edwards & Scrivener 1999).

#### 370 Mineralogical origin of magnetic properties

371 In summary, the magnetisation in these units is carried dominantly by haematite, with a likely 372 large range of grain size from superparamagnetic (pigmentary) haematite to larger (specularite) 373 particles of remanence carrying haematite. Magnetite may make a trace contribution. A strong 374 control on the concentration of haematite is related to the clay and silt content, and perhaps also 375 the concentration of volcanic rock detritus. The pigmentary haematite appears to have a 376 multiphase origin, ranging from possible pre-deposition to late diagenetic, a typical feature of 377 European Permian red beds (Turner et al. 1995; 1999). Largely detrital, specular haematite, 378 varies in amounts relating to the degree of sediment sorting and the sediment supply. In the 379 breccia units the maximum susceptibility axes reflect palaeocurrent-parallel trends, shown by clast imbrication directions. In aeolian transported sediments, the transverse trends in K<sub>max</sub> axes 380 381 reflect lee-face transport on dune slip faces. In the lacustrine mudstones the  $K_{max}$  directions may 382 represent wave-produced (or perhaps wind-related?) grain orientations in the lake playa systems 383 hence, the AMS shows a primary depositional fabric (unaffected by tectonism), probably carried 384 mostly by haematite.

#### 385 Palaeomagnetic Results

- 386 The majority of the 250 specimens demagnetised show little change in K<sub>lf</sub> during
- 387 demagnetisation, although the mudstones (particularly from the Littleham Mudstone Fm) tend to
- 388 show alteration at  $>600^{\circ}$ C, with lower temperature alteration in some specimens (Fig. 6). In some

- 389 specimens, this alteration obscures the recovery of the remanence at higher demagnetisation390 temperatures.
- 391

392 Demagnetisation isolates two remanence components. Firstly, a positive, often northerly, steeply 393 inclined component (Component A), between room temperature and often up to 350°C, but 394 sometimes up to 500-600°C (Fig. 8). This component is more northerly in specimens from the 395 Aylesbeare Mudstone Gp (Fisher mean,  $005^{\circ}$ ,  $+59^{\circ}$ , k=7.7, Ns=135), but more southerly in 396 specimens from the Exeter Gp (Fisher mean, 010°, +82°, k=6.7, Ns=44; see supplementary data). 397 This component is more prevalent in the Aylesbeare Mudstone Gp (79% of specimens) compared 398 to the Exeter Gp (56% of specimens), in which it is most prevalent in specimens from the 399 Dawlish Sandstone Fm. It does not correspond in direction particularly well to the expected 400 modern dipole field (i.e. inclination of 68°) and probably represents a composite component comprising mostly a Brunhes (viscous?) magnetisation plus the characteristic remanence. In 10% 401 402 of samples from the Aylesbeare Mudstone Gp, this was the only component present. In the 403 Exeter Gp 15% of specimens are dominated by this component, the bulk of these being from the 404 Dawlish Sandstone Fm.

405

406 A second component is recognised between about 400 and 650-700 °C that is a northerly, 407 positively inclined or southerly, negatively inclined direction (Fig. 8), interpreted as the 408 characteristic remanence (ChRM). In the Littleham Mudstone Fm the unblocking temperature 409 range of this component is mostly above  $500^{\circ}$ C-  $600^{\circ}$ C, whereas in specimens from the Exeter 410 Gp, the unblocking of the ChRM often starts at temperatures of  $\sim 400^{\circ}$ C. Some 52% of specimens 411 (49% in Aylesbeare Mudstone Gp and 57% in Exeter Gp) had suitable linear trajectory ChRM 412 line fits (here termed 'S-type' data; Fig. 8). This S-type demagnetisation behaviour was visually 413 classified into three quality classes, S1, S2 and S3 (Figs. 8, 9). The mean  $\alpha_{95}$  linear fits and  $\rho$  for 414 these classes indicate the generally larger model variance required to accommodate the less 415 quality line-fits (see supplementary data). Average confidence cone angles for these line-fit 416 classes vary from 3.2 to 13.9°. The mean directions for the ChRM line-fits pass the reversal test 417 (McFadden & McElhinney 1990), for all except those from the Littleham Mudstone Fm (Table 1; 418 Fig. 9a).

419

420 Some 28% of specimens displayed great circle trends, of varying arc length, towards interpreted

421 Permian reverse and normal polarity directions (here referred to as T-type demagnetization

422 behaviour; Fig. 8). This T-type behaviour was visually classified into three quality classes, T1,

423 T2 and T3, based on the visual length and scatter of the demagnetisation points about the great

424 circle, with T1 being the best quality. The mean  $\alpha_{95}$  for the poles to the fitted planes, for these

425 three data classes range from 9 to  $20^{\circ}$  (see supplementary data). These great circle fits included

- 426 the origin in 67% of these cases.
- 427

428 Data from the Dawlish Sandstone Fm yield the least well-defined results, particularly those from

429 Bishops Court Quarry, which are dominated by component A. These samples also display mainly

430 low blocking temperatures (i.e. the NRM is largely demagnetised by ~500°C). Some specimens

431 from this locality could be AF demagnetised, indicating that either these sandstones originally

432 had no haematite, or more likely a substantial proportion of haematite had been removed,

433 possibly by Quaternary ground water flow (e.g. Johnson et al. 1997). Notably, those samples that

did not retain a ChRM generally lacked specular haematite particles in thin section, whereas

435 samples of aeolian sandstone which possessed a ChRM often possessed specularite in small

436 amounts. Hence, the poor palaeomagnetic behaviour in the Bishops Court Quarry samples is due

to a paucity of specularite, and the dominance of pigment-dominated magnetisations, like

438 Permian aeolian sandstones such as the Penrith Sandstone (Turner *et al.* 1995).

#### 439 Mean directions and paleopoles

440 As well as the conventional means using the ChRM directions (Fig. 9), mean directions were also 441 determined using 'specimen-based' means, by combining the great circle paths with the specimen 442 line-fit ChRM (Table 1), to produce combined means using the method of McFadden & 443 McElhinney (1988). This method determines a mean direction, by including the 'fixed-point' 444 ChRM directions, and those points on the projected great circles, which maximise the resultant 445 length (i.e. points on the great circle which are closest to the combined mean direction). These 446 means are broadly similar to the line-fit ChRM means, except that for the Dawlish Sandstone and 447 Exe Breccia, which have steeper inclination and greater dispersion (Table 1). The great-circle 448 combined means pass the reversal test for the Littleham Mudstone Fm, Exmouth Mudstone and 449 Sandstone Fm, and Dawlish Sandstone plus Exe Breccia formations (Table 1). Using the line-fit 450 ChRM directions alone, the combined mean directions for the Aylesbeare Mudstone and Exeter 451 groups pass the reversal test (Table 1).

453 The S-class ChRM directions for the Exeter Gp, pass the fold test, indicating the pre-folding

454 nature of the magnetisations (Fig. 9b). The fold test of McFadden (1990) produced an f-statistic

455 (F [6,82]) of 1.90. Likewise, these data pass the DC fold test of Enkin (2003), with best

unfolding at 93.5%, with a 95% confidence interval of ±25.2%. A progressive unfolding test

457 (Watson & Enkin 1993) indicated best unfolding at 78%, with 95% confidence intervals on the

- 458 percent unfolding of 34% to 114% (Fig. 9b).
- 459

460 The virtual geomagnetic pole (VGP) data is consistent with other Permian data from stable-

461 Europe, confirming the Permian age of these magnetisations. The mean direction for the Exeter

462 Gp produces a virtual geomagnetic pole (VGP) similar to stable-Europe sediments from the

463 youngest part of the KRPS (see Supplementary Data), although the mean is slightly to the east of

the European apparent polar wander path of Torsvik & Cocks (2005). The Exeter Volcanic

465 Rocks VGP of Zijderveld (1967) is similar to that from the Aylesbeare Mudstone Gp (Table 1),

466 whereas the VGP pole for the Exeter Gp sediments from this study, is displaced slightly more to

467 the east (see Supplementary Data).

#### 468 Magnetostratigraphic Interpretation

469 The line-fit ChRM directions from the Aylesbeare Mudstone Gp (and Exe Breccia Fm) were 470 converted to virtual geomagnetic pole (VGP) latitude using the line-fit ChRM mean in Table 1 471 (Figs. 10a,b (iii)). For those specimens that had no line-fit, the point on their great circle nearest 472 this mean, were used for calculating the VGP latitude (Fig. 10). All specimens were also 473 assigned a polarity quality (Fig. 10a,b (ii)) based on the quality of demagnetisation behaviour 474 and, if from T-class specimens, the length and end point position of the great circle trend (similar 475 to the procedures used by Ogg & Steiner 1991; Hounslow & McIntosh 2003). One specimen of good-quality polarity (i.e. S-Type) was sufficient to define the horizon polarity, whereas with 476 477 specimens of poorer quality at least two are required (Figs. 4, 10). Some 12% of specimens failed 478 to yield data which could be used to determine horizon polarity (10% in Aylesbeare Mudstone 479 Gp, 15% in Exeter Gp) and eight horizons failed to yield any specimens which could reliably be 480 used to determine magnetic polarity (Figs. 4, 5). Most of these are from sandstones, with most of 481 these in the Dawlish Sandstone Fm at Bishops Court Quarry (Fig. 4).

482

All the samples collected from below the Exe Breccia Fm are of reverse polarity, with those
sections situated stratigraphically above the Langstone Rock outcrop having both reverse and

- 485 normal polarity (Figs. 4, 5). The single sample from the Knowle Sandstone Fm (Fig. 2; Table 1)
- 486 likewise confirms the reverse polarity results from the age-equivalent Exeter Volcanic Rocks
- 487 found by Creer (1957), Zijderveld (1967) and Cornwall (1967). Significantly, two sites in the
- 488 Torbay Breccia Fm sampled in the reconnaissance study of Cornwall (1967) produced reverse
- 489 polarity, suggesting that reverse polarity probably dominates to the base of the Exeter Gp.
- 490
- 491 Major magnetozone reverse and normal couplets have been numbered (Fig. 10) from the base of
- 492 the first normal polarity samples in the Exe Breccia Fm, using the prefix EA (for Exeter-
- 493 Aylesbeare). The magnetic polarities of six magnetozones are defined with multiple specimens
- 494 from a single sampling horizon (EA3n.1r, EA3n.2r, EA5n.1r), and EA3r.1n is defined with a
- 495 single specimen of S-class behaviour (Fig. 10b).

#### 496 **Discussion**

- 497 The major geomagnetic polarity marker in the Permian is the end of the Kiaman reverse polarity 498 Superchron, which has been comprehensively studied since the 1950's in Russian successions 499 (Molostovsky 1983; Burov et al. 1998). Studies on marine fossil-bearing rocks which 500 demonstrate the end of the Kiaman superchron are discontinuous studies in the SW USA (Steiner 501 2006), and Japan (Kirschvink et al. 2015), along with studies on successions in China (Steiner et 502 al. 1989; Embleton et al. 1996). The overlying reverse and normal polarity Illawarra Superchron, 503 has been investigated in marine successions in the Salt Range in Pakistan, China and Iran (Haag 504 & Heller 1991; Gallet et al. 2000; Jin et al. 2000; Steiner 2006), along with flood-basalts in 505 China (Ali et al. 2002; Zheng et al. 2010). Studies on non-marine rocks from the Illawarra 506 Superchron have been extensive in Russia, on outcrop and borehole material (Molostovsky 1983; 507 Burov et al. 1998) and core material from the Southern Permian Basin (Menning et al. 1988; 508 Nawrocki 1997; Turner et al. 1999; Lawton & Roberson 2003; Szurlies 2013). These studies 509 together allow the magnetic polarity stratigraphy (Fig. 11) to be defined through the Roadian to 510 Changhsingian (Steiner 2006; Hounslow submitted). The base of the Illawarra superchron is in 511 the lower to mid Wordian, based on magnetostratigraphic data from the Grayburg Fm in Texas 512 and New-Mexico (Steiner 2006) and limestones from Japan (Kirschvink et al. 2015). 513 514 Magnetostratigraphic studies in the southern Permian Basin well Mirow 1/1a/74 (Menning et al.
- 515 1988; Langereis et al. 2010), and wells in Poland (Nawrocki 1997) show a long-duration reverse
- 516 polarity interval (equivalent to MP3r –UP1r interval), with under and overlying mixed polarity-

517	inte	ervals (Fig. 11). The normal magnetozones in the Lower Drawa Fm and Havel Subgroup are						
518	probably equivalent with the MP1n to MP3n interval in the GPTS of Hounslow (submitted).							
519	Equivalent normal magnetozones in the Notec and Hannover formations, are more fully							
520	expressed by studies of the Lower Leman Sandstone, from the Johnston and Jupiter field in the							
521	sou	thern North Sea (Turner et al. 1999; Lawton & Roberson 2003; Fig. 11). These correlations						
522	are	constrained by the overlying Zechstein, and indicate that the Zechstein successions are						
523	ent	irely Changhsingian in age, rather than as old as early Wuchiapingian, as suggested by the						
524	cor	nodonts Merrillina divergens and Mesogondolella britannica (Korte et al. 2005; Legler et al.						
525	200	05; Słowakiewicz et al. 2009), and the synthesis of Szurlies (2013). Like the						
526	ma	gnetostratigraphic interpretation here (Fig. 11), Sr-isotope data indicates a short duration for						
527	the	Zechstein of ~ 2 Ma, with a likely age range of 255-251.5 Ma, placing it firmly in the						
528	Ch	anghsingian (Denison & Peryt 2009). Attempts at dating the Kupferschiefer at the base of the						
529	Zeo	chstein (Z1 cycle) have failed to yield consistent results, with Re-Os ages giving wide $95\%$						
530	cor	fidence intervals (Pašava et al. 2010).						
531								
532	Fou	ir pieces of information have allowed the Exeter Gp succession to be assigned to the Permian.						
533	1)	Volcanic units interbedded with the Knowle Sandstone of the Exeter area, and similar units						
534		equivalent to the Thorverton Sandstone and Bow Breccia in the Crediton Trough, have Ar-Ar						
535		ages of 291- 282 Ma (Edwards & Scrivener 1999). Volcanic clasts in the breccia units give						
536		Ar-Ar dates of 280 Ma. This suggests the volcanism and associated interbedded sediments						
537		are late Sakmarian through to late Artinskian in age, using the timescale of Henderson et al.						
538		(2012).						
539	2)	The Dartmoor Granite has Rb-Sr, U-Pb and Ar-Ar ages of 280 ±1 Ma (Scrivener 2006),						
540		placing its formation in the latest Artinskian (timescale of Henderson et al. 2012). Clasts of						
541		the granite begin to occur in the unroofing succession of the Teignmouth and Heavitree						
542		Breccias (Dangerfield & Hawkes 1969; Edwards et al. 1997), indicating that these breccias						
543		were deposited several millions years after the granite formation, in order to allow time for						
544		granite unroofing.						
545	3)	Miospore assemblages containing Lueckisporites virkkiae, occur from the Whipton Fm,						
546		around Exeter, but also in younger units in the Crediton Trough, equivalent to the Alphington						
547		and Heavitree Breccias (Edwards et al. 1997). Assemblages containing this miospore are						
548		widespread in European Zechstein deposits and similar 'Thuringian' and Russian Tatarian-						

549 age assemblages (Visscher 1973; Utting 1996). In the northern hemisphere, *Lueckisporites* 

- *virkkiae* has its first appearance in the latest Kungurian (Shu 1999; Mangerud 1994) to early
  Roadian (lower Kazanian in Russia; Utting 1996), but occurrences range into the latest
  Changhsingian.
- 4) The foot-print trace-fossil *Cheilichnus bucklandi*, found in the Dawlish Sandstone near
  Exeter (Edwards *et al.* 1997) suggests equivalence to the Germanic 'Rotliegend' (McKeever
  & Haubold 1996). However, this genus is restricted to aeolian dune units and is probably
- only vaguely indicative of the Permian (Lucas & Hunt 2006).
- 557

558 Constraints on the youngest possible age of the Aylesbeare Mudstone Gp are

559 magnetostratigraphy and vertebrate fossils from the overlying Otter Sandstone Formation

560 (Hounslow & McIntosh 2003; Benton 1997), which indicate the Sherwood Sandstone Gp is as

old as early Anisian (Middle Triassic), and probably ranges down into the Olenekian of the

562 Lower Triassic (Hounslow & McIntosh 2003; Hounslow & Muttoni 2010). Based on regional

climate comparisons between the Budleigh Salterton Pebble Beds and the 'Conglomérate

564 principal' of the Vosges region in NE France, Durand (2006) suggested a probable early

- 565 Olenekian age for the Budleigh Salterton Pebble Beds Fm, generally consistent with the 566 magnetostratigraphy.
- 567

568 This work detects the oldest normal magnetozone in the mid-parts of the Exe Breccia (i.e. EA1n), 569 with a substantial thickness (perhaps up to  $\sim 1$  to 1.5 km) of reverse polarity in the underlying 570 parts of the Exeter Gp. Although we cannot locate the base of EA1n precisely due to lack of 571 suitable outcrop, this normal magnetozone is the earliest evidence of the Illawarra Superchron 572 (Fig. 11). EA1n is probably equivalent to early MP2n, or the normal polarity part of MP1. 573 However, ~55 m of unsampled strata occur in the interval between our outcrops at the Exe 574 Breccia- Dawlish Sandstone boundary (Figs. 4, 11). Therefore, it is possible the equivalent of the 575 MP1n chron is within this unsampled interval. The end of the KRPS provides an important 576 dating-point (267.1±0.8 Ma; Hounslow submitted) to the early-mid Wordian in the Middle 577 Permian (Guadalupian). The oldest occurrence of the Lueckisporites virkkiae assemblage is 578 found in the Whipton Fm, which suggests that this formation could be as old as latest Kungurian 579 to early Roadian (~272 Ma; Henderson et al. 2012). This would give a minimum of ~8 Ma of 580 exhumation time, between formation of the Dartmoor Granite, and the first granite detritus 581 appearing in the Teignmouth-Heavitree breccias. 582

- 583 The overlying normal polarity magnetozone EA3n, is therefore likely to be equivalent to the
- 584 MP3n normal magnetozone in the upper and mid parts of the Capitanian (Fig. 11). The EA3r
- 585 magnetozone is equivalent to the MP3r to UP1r interval (in the lower part of the Wuchiapingian),
- 586 with the overlying normal magnetozones (i.e. EA4n to EA5n) equivalent to those in the upper
- 587 parts of the Wuchiapingian to basal Changhsingian (Fig. 11). Reverse magnetozone EA2r, in the
- top of the Exe Breccia is probably the equivalent of MP2r in the basal Capitanian. Sub
- 589 magnetozone EA3r.1n in the Littleham Mudstone Fm is probably equivalent to UP1n in the
- 590 Wuchiapingian.
- 591 Alternative Lower Triassic age models?

592 The alternative Lower Triassic age of the Aylesbeare Mudstone Gp suggested by Warrington &

593 Scrivener (1990) and Edwards *et al.* (1997), is untenable using the magnetostratigraphy. To

594 evaluate their hypothesis using this data suggests the most likely early Triassic correlation model

595 would indicate EA3n is the age equivalent of the first Triassic magnetozone, LT1n (Fig. 11).

596 Therefore, the overlying EA3r to EA5n interval would extend into the earliest Olenekian, an

- interval of some 1.4 Ma (Hounslow & Muttoni 2010). However, this seems unlikely for thefollowing reasons:
- 599

The local clast lithologies (e.g. murchisonite) seen in the breccia at the base of the Littleham
Mudstone Fm, are similar to those in the Exeter Gp, and very different to those found in the
Budleigh Salterton Pebble Beds (and other Lower Triassic units further north in the UK).

- Ducleigh Sulerton record Deus (and other Dower Triassie ands rather north in the Ork),
- which contain Armorican-derived clasts and zircons from Cadomian basement (Cocks 1993;
  Morton *et al.* 2013).
- 605 2) It would require a minimum hiatus of ~ 13-15 Ma between the Exe Breccia and the
- Aylesbeare Mudstone Gp, which seems unlikely considering the apparently conformablenature of the boundary between these formations in the Exe Estuary area.

6083)If the hypothesis of Warrington & Scrivener (1990) was correct, it would predict numerous

- normal polarity intervals (from the Illawarra superchron) below the Aylesbeare Mudstone Gp
- but we have only found these in the Exe Breccia Fm with no evidence of normal polarity inthe underlying *c*.1 km of the Exeter Gp.
- 612 4) This Lower Triassic model would suggest a ~1.4 Ma duration for the EA3n to EA5n interval
  613 requiring very large accumulation rates, comparable to the deepest grabens in the Southern

614 Permian Basin, north of the Variscan front, which there contain substantial thicknesses of615 Zechstein.

#### 616 Wider regional implications

A consequence of these data is that it is now possible to assess the relationship of the SW 617 England successions to the much better studied Rotleigend-II group in the Southern Permian 618 619 Basin (Fig. 11). The magnetostratigraphy suggests a similarity in age of the Altmark 620 unconformities with the Devon Permian sequence boundaries. The magnetic polarity stratigraphy 621 from the Mirow, Czaplinek and Piła wells suggests that the Altmark III unconformity is roughly 622 equivalent to the base of the Littleham Mudstone Fm (base of Pm5 sequence; Fig. 2), Altmark II, 623 with the base of sequence Pm4 (Figs. 2, 11). Less certain is the correlation of the base of unit B 624 in the Littleham Mudstone Fm, with Altmark IV. The base of sequence Pm3 probably relates to 625 the Altmark I unconformity, which separates the Muritz Subgroup from the Havel Subgroup,

626 across the Saalian unconformity, since underlying successions both contain volcanic units.

627

628 The calcrete and rhizoconcretion bearing sandstone, in the uppermost part of the Straight Point 629 Sandstone Mbr, is unusual in that no other well developed palaeosols are seen in the remainder 630 of these Permian successions. It is not until the mid Triassic (Anisian) Otter Sandstone Fm, that 631 calcretes began to be widely developed in SW England. The Capitanian-Wuchiapingian 632 boundary was an interval with dramatic, but poorly understood shifts in the global carbon cycle 633 (Nishikane et al. 2014). A tentative reason for this palaeosol development is the rapid warming 634 associated with increased  $CO_2$  in the atmosphere (and associated increased evaporation rates to create calcretes; Alonso-Zarza 2003), that developed after the extinction at the Capitanian-635 Wuchiapingian boundary. The peak is associated with a negative  $\delta^{13}$ C excursion (Chen *et al.* 636 637 2011: Nishikane *et al.* 2014) in the early Wuchiapingian, which corresponds closely to the early 638 parts of MP3r (Zheng et al. 2010; Fig. 11).

639

640 The dramatic switch between breccia-dominated facies of the Exeter Gp to the mudstone-

dominated facies of the Aylesbeare Mudstone Gp, occurs within the early Capitanian (Fig. 11).

642 We tentatively relate this switch in facies to the Kamura cooling event (seen as a large positive

- 643 δ<sup>13</sup>C excursion during the Capitanian), which began in the early Capitanian (Isozaki *et al.* 2011).
- 644 This has been associated with lows in atmospheric CO<sub>2</sub>, and cooler oceanic surface waters in
- both the Panthalassa and Paleo-Tethys Oceans (Isozaki et al. 2011; Nishikane et al. 2014). This

- 646 cooling event may have allowed more moisture-bearing weather systems to penetrate further
- northwards into the heart of Pangaea, from the Paleo-Tethys, so allowing delivery of larger
- amounts of mud into the playa systems of the Aylesbeare Mudstone Gp.
- 649

650 The Southern Permian Basin, Parchim and Mirow formations shows a number of similarities to 651 the Devon successions. The Parchim Fm dominantly comprises thick conglomeratic braidplain-652 type deposits, extending to sandflat and locally playa mudstone deposits in the basin centre 653 (McCann 1998; Rieke et al. 2003). Tectonic control of facies was important during the Parchim 654 Fm. Like the Exeter Group in sequence Pm3 (Fig. 2) the Parchim Fm has an earlier wetter phase 655 and a later dryer phase (Rieke et al. 2003). This is overlain by the Mirow Fm which is 656 characterized by the progradation of sand-prone fluvial facies with frequent claystones, over a 657 much wider extent in the Southern Permian Basin than the Parchim Fm. The rarity of conglomerates (except at basin margins), with instead claystones (containing fossils indicative of 658 659 freshwater conditions) and the dominance of sand-prone facies, is very different to the 660 underlying Parchim Fm (McCann 1998). Hence, the start of the Mirow Fm sees a switch to 661 climatically wetter conditions (Rieke et al. 2003), like those seen in the Aylesbeare Mudstone Gp. The coincidence in timing and the switch to wetter environmental conditions, seen in the 662 663 Devon successions and German basins, suggests these major facies changes are climatically 664 controlled.

### 665 Conclusions

666 The palaeomagnetic signal in the Exeter and Aylesbeare Mudstone groups is dominantly carried 667 by haematite, whose mean directions pass the reversal test. The remanence in the Exeter Gp 668 passes a fold test. The AMS indicates the fabric carried by haematite is detrital in origin. Reverse polarity dominates in the lower part of the Exeter Gp, with the start of the Illawarra superchron, 669 in the Exe Breccia Fm dated to the early Wordian. Five normal-reverse couplets are found in the 670 671 overlying sediments starting in the upper part of the Exe Breccia Fm (Langstone Mbr) and into 672 the Aylesbeare Mudstone Gp. This magnetostratigraphic data allows the Exmouth Mudstone and 673 Sandstone Fm to be assigned to the Capitainian to the earliest Wuchiapingian, and the overlying 674 Littleham Mudstone Fm is earliest Wuchiapingian, to as young as the Wuchiapingian-Changhsingian boundary. With these data, the Permian successions in SW England are now the 675 676 most precisely dated Permian succession in the UK, and if similar methods were used elsewhere, 677 should provide the foundation for a much better understanding of other UK Permian basins. The

- similarity in the timing between facies changes and sequence boundaries, here and those of the
- 679 Rotliegend-II Group in the Southern Permian Basin, indicates that palaeoclimatic change is a
- 680 fundamental metric in their subdivision. The question of the position of the Permo-Triassic
- boundary in SW England has now been effectively resolved, and ironically corresponds to the
- 682 position taken by Victorian geologists such as Irving (1888).

#### 683 Acknowledgements

- 684 This work was part-funded by a British Geological Survey- University collaboration grant.
- 685 Richard Scrivener led fieldwork to the Crediton Trough, and Paulette Posen assisted during
- 686 fieldwork to the Exeter Gp. Robert Hawkins and Laurence Thistlewood measured some of the
- 687 samples. Sylvie Bourquin and Antoine Bercovici assisted in the fieldwork in 2007. Simon Chew
- drafted some of the figures. The MOD kindly allowed access to the Straight Point firing range. A
- 689 reviewer Tim Raub provided constructive suggestions for improvement.

#### 690 **References**

- Ali, J. R, Thompson, G. M., Song, X. & Wang, Y. 2002. Emeishan Basalts (SW China) and the
  'end-Guadalupian' crisis: magnetobiostratigraphic constraints. *Journal of the Geological Society*, 159, 21-29.
- Alonso-Zarza, A. M. 2003. Palaeoenvironmental significance of palustrine carbonates and
   calcretes in the geological record. *Earth-Science Reviews*, **60**, 261-298.
- 696 Barton, C. M., Woods, M. A., Bristow, C. R., Newell, A. J., Westhead, R. K., Evans, D. J.,
- 697 Kirby, G. A., & Warrington, G. 2011. The geology of south Dorset and south-east Devon
- 698 *and its World Heritage Coast.* Special Memoir of the British Geological Survey, Sheets 328,
- 699 341/342, 342/343 and parts of 326/340, 327, 329 and 339. HMSO, London
- Bateson, J. H. & Johnson, C. C. 1992. *Reduction and related phenomena in the New Red Sandstone of south-west England*, British Geological Survey, Technical report WP/92/1.
- Benton, M. J., Cook, E., & Turner, P. 2002. *Permian and Triassic red beds and the Penarth Group of Great Britain*. Geological Conservation Review Series. Joint Nature
- 704 Conservation Committee, Peterborough.
- Benton, M. J. 1997. The Triassic reptiles from Devon. *Proceedings of the Ussher Society*, 9, 141152.
- Burov, B. V., Zharkov, I. Y., Nurgaliev, D. K., Balabanov, Yu. P., Borisov, A. S. & Yasonov, P.
  G. 1998. Magnetostratigraphic characteristics of Upper Permian sections in the Volga

709	and the Kama areas. In: Esaulova, N. K., Lozonsky, V. R., Rozanov, A. Yu. (eds).
710	Stratotypes and reference sections of the Upper Permian in the regions of the Volga and
711	Kama Rivers. GEOS, Moscow, 236-270.
712	Butler, M. 1998. The geological history and the southern Wessex Basin- a review of new
713	information from oil exploration. In: Underhill, J. R. (ed.), Development, evolution and
714	petroleum geology of the Wessex Basin. Geological Society London Special Publication
715	<b>133</b> , 67-86.
716	Carter, A. Yelland, A. Bristow, C. & Hurford, A. J. 1995. Thermal histories of Permian and
717	Triassic basins in Britain derived from fission track analysis. In: Boldy, S.A. R. (ed.),
718	Permian and Triassic Rifting in Northwest Europe, Geological Society Special
719	Publication, <b>91</b> , 41-56.
720	Chapman, T. J. 1989. The Permian to Cretaceous structural evolution of the Western Approaches
721	Basin (Melville sub-basin), UK. In: M.A. Cooper (ed.), Inversion Tectonics, Geological
722	Society, London, Special Publications; 44, 177-200.
723	Chen, B., Joachimski, M. M., Sun, Y., Shen, S. & Lai, X. 2011. Carbon and conodont apatite
724	oxygen isotope records of Guadalupian-Lopingian boundary sections: Climatic or sea-
725	level signal? Palaeogeography, Palaeoclimatology, Palaeoecology, 311, 145-153.
726	Cocks, L. R. M. 1993. Triassic pebbles, derived fossils and the Ordovician to Devonian
727	palaeogeography of Europe. Journal Geological Society London, 150, 219-226.
728	Cornwall, J. D. 1967. Palaeomagnetism of the Exeter Lavas, Devonshire. Geophys. Journal
729	Royal Astro. Soc., 12, 181-196.
730	Creer, K. M. 1957. The Natural Remanent Magnetization of Certain Stable Rocks from Great
731	Britain. Philosophical Transactions of the Royal Society of London. Series A, Mathematical
732	and Physical Sciences, <b>250</b> , 111-129.
733	Dangerfield, J. & Hawkes, J. R. 1969. Unroofing of the Dartmoor Granite and possible
734	consequences with regard to mineralization. Proc. Ussher Soc., 2, 122-131.
735	Denison, R. E. & Peryt, T. M. 2009. Strontium isotopes in the Zechstein (Upper Permian)
736	anhydrites of Poland: evidence of varied meteoric contributions to marine brines. Geological
737	Quarterly, <b>53</b> , 15
738	Durand, M. 2006. The problem of the transition from the Permian to the Triassic Series in
739	southeastern France: comparison with other Peritethyan regions. In: Lucas, S. G., Cassinis,
740	G. & Schneider J.W. (eds) Non-Marine Permian Biostratigraphy and Biochronology,
741	Geological Society, London, Special Publications, 265, 281-296.

742	Durrance, E. M., Meads, R. E., Ballard, R. R. B. & Walsh, J. N. 1978. Oxidation state of iron in
743	the Littleham Mudstone Formation of the new red sandstone series (Permian-Triassic) of
744	southeast Devon, England. Geological Society of America Bulletin, 89, 1231-1240.
745	Edwards, R. A. & Scrivener, R. C., 1999. Geology of the country around Exeter. Memoir of the
746	British Geological Survey, Sheet 325 (England and Wales), HMSO, London.
747	Edwards, R. A., Warrington, G., Scrivener, R. C., Jones, N. S., Haslam, H. W. & Ault, L. 1997.
748	The Exeter Group, south Devon, England: a contribution to the early post-Variscan
749	stratigraphy of northwest Europe. Geological Magazine, 134, 177-197.
750	Ellwood, B. B. & Howard, J. H. 1981. Magnetic fabric development in an experimentally
751	produced barchan dune. Journal of Sedimentary Research, 51, 97-100.
752	Embleton, B. J. J., McElhinny, M. W., Zhang Z. & Li Z. X. 1996. Permo-Triassic
753	magnetostratigraphy in China: the type section near Taiyuan, Shanxi Province, North
754	China. Geophys. J. Int. 126: 382–388
755	Enkin, R. J. 2003. The direction- correction tilt test: an all purpose tilt/fold test for
756	palaeomagnetic studies. Earth Planet. Sci. Lett., 212, 151-166.
757	Evans, C. D. R. 1990. The geology of the western English Channel and its western Approaches.
758	UK Offshore regional report. British Geological Survey, HMSO, London.
759	Frank, U. and Nowaczyk, N. R. 2008. Mineral magnetic properties of artificial samples
760	systematically mixed from haematite and magnetite. Geophys. J. Int. 175, 449-461.
761	Gallet, Y., Krystyn, L., Besse, J., Saidi, A. & Ricou, L-E. 2000. New constraints on the upper
762	Permian and Lower Triassic geomagnetic polarity timescale from the Abadeh section
763	(central Iran). Journal of Geophysical Research, 105, 2805-2815.
764	Gallois, R.W. 2014. The position of the Permo-Triassic boundary in Devon, UK. Geoscience in
765	South-West England, <b>13</b> , 328-338.
766	Haag, M. & Heller, F. 1991. Late Permian to Early Triassic magnetostratigraphy. Earth Planet.
767	Sci. Letter, <b>107</b> , 42-54.
768	Hamblin, R. J. O., Crosby, A., Alson, R. S., Jones, S. M., Chadwick, R. A., Penn, I. E. & Arthur,
769	M. J. 1992. United Kingdom off-shore report: the geology of the English Channel.
770	British Geological Survey, HMSO, London.
771	Harvey, M. J. Stewart, S. A., Wilkinson, J.J., Ruffell, A. H. & Shall, R. K. 1994. Tectonic
772	evolution of the Plymouth Bay Basin. Proceedings of the Ussher Society, 8, 271-278.

773 Henderson, C. M., Davydov, V. I. & Wardlaw, B. R. 2012. The Permian Period. In: Gradstein, 774 F.M., Ogg, J.G, Schmitz, M.D. & Ogg, G.M (eds), The geologic time scale 2012, vol. 2, 775 Elsevier, Amsterdam, 653-679. 776 Henson, M. R. 1972. The form of the Permo-Triassic basin in south-east Devon. Proc. Ussher 777 Soc. 3, 447-457. 778 Hounslow, M.W. & McIntosh, G. 2003. Magnetostratigraphy of the Sherwood Sandstone Group 779 (Lower and Middle Triassic): South Devon, U.K.: Detailed correlation of the marine and 780 non-marine Anisian. Palaeogeogr. Palaeoclimat. Palaeoecol., 193, 325-348. 781 Hounslow M.W. & Muttoni G. 2010. The geomagnetic polarity timescale for the Triassic: 782 Linkage to stage boundary definitions. In: Lucas, S. G. (ed.) The Triassic timescale. 783 Special Publication of the Geological Society, **334**, 61-102. 784 Hounslow, M. W. (submitted). Palaeozoic geomagnetic reversal rates following superchrons 785 have a fast re-start mechanism. Nature Communications. 786 Irving, A. 1888. The red-rock series of the Devon Coast section. Quart. J. geol. Soc., 44, 149-787 163. 788 Isozaki, Y., Aljinovic, D. & Kawahata, H. 2011. The Guadalupian (Permian) Kamura event in 789 European Tethys. Palaeogeography, Palaeoclimatology, Palaeoecology 308, 12–21. 790 Jin, Y. Shang, Q. & Cao, C. 2000. Late Permian magnetostratigraphy and its global correlation. 791 Chinese Science Bulletin, 45, 698-705. 792 Johnson, S. A., Glover B. W. & Turner, P. 1997. Multiphase reddening and weathering events in 793 Upper Carboniferous red beds from the English West Midlands. Journal of the 794 Geological Society, 154, 735-745. 795 Kent, J. T., Briden, J. C. & Mardia, K. V. 1983. Linear and planar structure in ordered 796 mulivariate data as applied to progressive demagnetisation of palaeomagnetic remanence. 797 Geophysical Journal Royal Astronomical Society, 81, 75-87. 798 Kirschvink, J. L., Isozaki, Y., Shibuya, H., Otofuji, Y. I., Raub, T. D., Hilburn, I. A., Kasuya, T., 799 Yokoyama, M. & Bonifacie, M. 2015. Challenging the sensitivity limits of 800 Paleomagnetism: Magnetostratigraphy of weakly magnetized Guadalupian-Lopingian 801 (Permian) Limestone from Kyushu, Japan. Palaeogeography, Palaeoclimatology, 802 Palaeoecology, 418, 75-89. Korte, C., Kozur, H. W. & Veizer, J. 2005.  $\delta^{13}$  C and  $\delta^{18}$  O values of Triassic brachiopods and 803 804 carbonate rocks as proxies for coeval seawater and palaeotemperature. Palaeogeography, 805 Palaeoclimatology, Palaeoecology, 226, 287-306.

806	Kostadinova M., Jordanova N., Jordanova D. & Kovacheva M. 2004. Preliminary study on the
807	effect of water glass impregnation on the rock-magnetic properties of baked clay. Studia
808	Geophysica et Geodaetica, <b>48</b> , 637-646.
809	Laming, D. J. C. 1966. Imbrications, palaeocurrents and other sedimentary features in the Lower
810	New Red Sandstone, Devonshire, England. Journal of Sedimentary Petrology, 36, 940-
811	959.
812	Laming, D. J. C. 1969. A guide to the New Red Sandstone of Tor Bay, Petitor and Shaldon.
813	Report Transaction of Devonshire Association of Science, Literature and Art, 101, 207-
814	218.
815	Laming, D. J. C. 1982. The New Red Sandstone. In: Durrance, E. M. & Laming, D. J. C. (eds)
816	The geology of Devon. University of Exeter Press, Exeter, 148-178,
817	Laming, D. J. C. & Roche, D. P. 2013. Faulting in Permo-Triassic strata and buried channels
818	revealed by excavation of the Lympstone-Powderham tunnel, Exe Estuary, Devon.
819	Geoscience in South-West England, 13, 244.
820	Langereis, C. G., Krijgsman, W., Muttoni, G. & Menning, M. 2010. Magnetostratigraphy -
821	concepts, definitions, and applications. Newsletters on Stratigraphy, 43, 3, 207-233
822	Lawton, D. E. & Roberson, P. P. 2003. The Johnston Gas Field, Blocks 43/26a, 43/27a, UK
823	Southern North Sea. In: Gluyas, J. & Hichens, H. M. (eds), United Kingdom Oil and Gas
824	Fields, Commemorative Millennium Volume. Geological Society London Memoir, 20,
825	749-759.
826	Legler, B., Gebhardt, U. & Schneider, J. W. 2005. Late Permian non marine to marine
827	transitional profiles in central southern Permian Basin, northern Germany. Int. Journal of
828	Earth Sciences, 94, 851-862.
829	Leveridge, B. E. Scrivener, R. C. Goode, A. J. J. & Merriman, R. J. 2003. Geology of the
830	Torquay district, a brief explanation of the geological map sheet 350 Torquay.
831	Keyworth: British Geological Survey, HMSO, London.
832	Løvlie, R. & Torsvik, T. 1984: Magnetic remanence and fabric properties of laboratory-deposited
833	hematite-bearing red sandstone. Geophysical Research Letters, 11, 229-232.
834	Lowrie, W. 1990. Identification of ferromagnetic minerals in a rock by coercivity and unblocking
835	temperature properties. Geophysical Research Letters, 17, 159-162.
836	Lucas, S. G. & Hunt, A. P. 2006. Permian tetrapod foorptints: biostratigraphy and biochronology.
837	In: Lucas, S.G. Cassinis, G. & Schneider, J.W. (eds), Non-marine Permian

838	biostratigraphy and biochronology, Geological Society, London Special Publications,
839	<b>265</b> , 179-200.
840	Mangerud, G. 1994. Palynostratigraphy of the Permian and lowermost Triassic succession,
841	Finnmark Platform, Barents sea. Review of Palaeobotany & Palynology, 82, 317-349.
842	McCann, T., Pascal, C., Timmerman, M. J., Krzywiec, P., López-Gómez, J., Wetzel, L. &
843	Lamarche, J. 2006. Post-Variscan (end Carboniferous-Early Permian) basin evolution in
844	western and central Europe. In: Gee, D. G. & Stephenson, R. A. (eds) European
845	Lithosphere Dynamics. Geological Society London Memoirs, 32, 355–388.
846	McCann, T. 1998. Sandstone composition and provenance of the Rotliegend of the NE German
847	Basin. Sedimentary Geology, 116, 177-198.
848	McFadden, P. L. 1990. A new fold test for palaeomagnetic studies. Geophysical Journal
849	International, <b>103</b> , 163-169.
850	McFadden, P. L. & McElhinney, M. W. 1988. The combined analysis of remagnetisation circles
851	and direct observations in palaeomagnetism. Earth Planetary Science Letters, 87, 161-
852	172.
853	McFadden, P. L. & McElhinny, M. W. 1990. Classification of the reversal test in
854	palaeomagnetism. Geophysical Journal International, 103, 725-729.
855	McKeever, P. M. & Haubold, H. 1996. Reclassification of vertebrate trackways from the Permian
856	of Scotland and related forms from Arizona and Germany. Journal of Paleontology, 70,
857	1011-1022.
858	McKie, T. & Williams, B. P. J. 2009. Triassic palaeogeography and fluvial dispersal across the
859	northwest European Basins. Geological Journal, 44, 711-741.
860	Menning, M., Katzung, G. & Lützner, H. 1988. Magnetostratigraphic investigations in the
861	Rotliegendes (300-252 Ma) of Central Europe. Z. geol. Wiss., Berlin, 16, 1045-1063.
862	Merefield, J. R., Brice, C. J. & Palmer, A. J. 1981. Caesium from former Dartmoor volcanism:
863	its incorporation in NewRed sediments of SW England. Journal of the Geological Society,
864	<b>138,</b> 145-152.
865	Molostovsky, E. A. 1983. Paleomagnetic stratigraphy of the eastern European part of the
866	USSR. University of Saratov, Saratov [In Russian].
867	Morton, A. C., Hounslow, M. W. & Frei, D. 2013. Heavy-mineral, mineral-chemical and zircon-
868	age constraints on the provenance of Triassic sandstones from the Devon coast, southern
869	Britain. Geologos 19, 67–85.

870 Nawrocki, J. 1997. Permian to early Triassic magnetostratigraphy from the central european 871 basin in poland: Implications on regional and worldwide correlations. Earth. Planet. Sci 872 Lett., 152, 37-58. 873 Newell, A. J. 2001. Bounding surfaces in a mixed aeolian-Fluvial system (Rotliegend, Wessex 874 Basin, SW UK). Marine & Petroleum Geology, 18, 339-347. 875 Nishikane, Y., Kaiho, K., Henderson, C. M., Takahashi, S. & Suzuki, N. 2014. Guadalupian-876 Lopingian conodont and carbon isotope stratigraphies of a deep chert sequence in Japan. 877 Palaeogeography, Palaeoclimatology, Palaeoecology, 403, 16-29. 878 Ogg, J. G. & Steiner, M. B. 1991. Early Triassic polarity time-scale: integration of 879 magnetostratigraphy, ammonite zonation and sequence stratigraphy from stratotype 880 sections (Canadian Arctic Archipelago). Earth and Planetary Science Letters, 107, 69-881 89. 882 Ormerod- Wareing, G. 1875. On the Murchisonite beds of the Estuary of the Exe and an attempt 883 to classify the beds of the Trias thereby. Quart, J. Geol. Soc. Lond., 31, 346-354. 884 Park, M. E., Cho, H., Son, M. & Sohn, Y. K. 2013. Depositional processes, paleoflow patterns, 885 and evolution of a Miocene gravelly fan-delta system in SE Korea constrained by anisotropy of magnetic susceptibility analysis of interbedded mudrocks. Marine and 886 887 Petroleum Geology, 48, 206-223. 888 Pašava, J., Oszczepalski, S. & Du, A. 2010. Re-Os age of non-mineralized black shale from the 889 Kupferschiefer, Poland, and implications for metal enrichment. Mineralium Deposita, 890 45, 189-199. 891 Rieke, H., McCann, T., Krawczyk, C. M. & Negendank, J. F.W. 2003. Evaluation of controlling 892 factors on facies distribution and evolution in an arid continental environment: an 893 example from the Rotliegend of the NE German Basin. In: McCann, T. & Saintot. A. 894 (eds) Tracing Tectonic Deformation Using the Sedimentary Record. Geological Society, London, Special Publications, 208, 71-94. 895 896 Roscher, M. & Schneider, J. W. 2006. Permo-Carboniferous climate: Early Pennsylvanian to late 897 Permian climate development of central Europe in a regional and global context. In: 898 Lucas, S. G. Cassinis, G. & Schneider, J. W. (eds) Non-marine Permian biostratigraphy 899 and biochronology, Geological Society, London Special Publication, 265, 15-38. 900 Schwarzacher, W. 1951. Grain orientation in sands and sandstones. Journal of Sedimentary 901 Research, 21, 162-172.

- Scrivener, R. C. 2006. Cornubian granites and mineralization in SW England. *In:* Brenchley, P. J.
  & Rawson, P. F. *Geology of England and Wales*. Geological Society of London
  Publication, Bath, 257-267.
- Selwood, E. B. Edwards, R. A., Simpson, S., Chesher, J. A. & Hamblin, R. A. 1984. *Geology of the country around Newton Abbot. Memoir for 1:50,000 geological sheet 339*, British
  Geological Survey, HMSO, London.
- 908 Shen, S-Z., Henderson, C. H., Bowring, S. A., Cao1, C-Q. Wang, Y., Wang, W., Zhang, H.,
- 209 Zhang, Y-C. & Mu, L. 2010. High-resolution Lopingian (Late Permian) timescale of
  210 South China. *Geological Journal*, 45, 122–134.
- Shu, O. 1999. A brief discussion on the occurrences of *Scutasporites unicus* and *Lueckisporites virkkiae* complexes in the northern hemisphere. *Permophiles*, **33**, 21-23.
- Słowakiewicz, M., Kiersnowski, H. & Wagner, R. 2009. Correlation of the Middle and Upper
  Permian marine and terrestrial sedimentary sequences in Polish, German and USA
  Western Interior Basins with reference to global time markers. *Palaeoworld*, 18, 193211.
- 917 Steiner, M. 2006. The magnetic polarity timescale across the Permian-Triassic boundary. In:
- 918 Lucas, S. G. Cassinis, G. and Schneider, J. W. (eds), *Non-marine Permian biostratigraphy*
- 919 *and biochronology*, Geological Society, London Special Publication, **265**, 15-38.
- 920 Steiner, M. B., Ogg, J., Zhang, Z. & Sun, S. 1989. The Late Permian/early Triassic magnetic
- 921 polarity time scale and plate motions of south China. *Journal Geophysical Research*, 94,
  922 7343-7363.
- Strachan, R. A., Linnemann, U., Jeffries, T., Drost, K. & Ulrich, J. 2014. Armorican provenance
  for the mélange deposits below the Lizard ophiolite (Cornwall, UK): evidence for Devonian
  obduction of Cadomian and Lower Palaeozoic crust onto the southern margin of Avalonia.
- 926 International Journal of Earth Sciences, **103**, 1359-1383.

927 Szurlies, M., Bachmann, G. H., Menning, M., Nowaczyk, N. R. & Käding, K-C. 2003.

- Magnetostratigraphy and high resolution lithostratigraphy of the Permian- Triassic boundary
   interval in Central Germany. *Earth and Planetary Science Letters*, 212, 263-278.
- 930 Szurlies, M. 2013. Late Permian (Zechstein) magnetostratigraphy in western and central Europe.
- 931 In: Gasiewicz, A. & Słowakiewicz, M. (eds), Palaeozoic climate cycles: their evolutionary
- 932 *and sedimentological impact.* Geological Society, London, Special Publications, **376**, 73-85.
- 933 Tarling, D. H. & Hrouda, F. 1993. The magnetic anisotropy of rocks. Chapman and Hall,
- 934 London.

935	Thomas, H. H. 1909. A contribution to the petrography of the New Red Sandstone in the West
936	of England. Quarterly Journal of the Geological Society, London, 65, 229-245.
937	Timmerman, M. J. 2004. Timing, geodynamic setting and character of Permo-Carboniferous
938	magmatism in the foreland of the Variscan Orogen, NW Europe. In: Wilson, M.,
939	Neumann, ER., Davies, G. R., Timmerman, M. J., Heeremans, M. & Larsen, B. T.
940	(eds) Permo-Carboniferous Rifting and Magmatism in Europe, Special Publication
941	Geological Society, London, 223, 41-74.
942	Torsvik T. H. & Cocks L. R. M. 2005. Norway in space and time: A Centennial cavalcade.
943	Norwegian Journal of Geology, <b>85</b> , 73-86.
944	Turner, P. 1979. The palaeomagnetic evolution of continental red beds. Geological Magazine,
945	<b>116</b> , 289-301.
946	Turner, P., Burley, S. D. Rey, D. & Prosser, J. 1995. Burial history of the Penrith Sandstone
947	(Lower Permian) deduced from the combined study of fluid inclusion and palaeomagnetic
948	data. In: Turner, P. & Turner, A. (eds) Paleomagnetic applications in hydrocarbon
949	exploration. Geological Society London Special Publications, 98, 43-78.
950	Turner, P., Chandler, P., Ellis, D., Leveille, G. P. & Heywood, M. L. 1999. Remanance
951	acquisition and magnetostratigraphy of the Leman Sandstone Formation: Jupiter Fields,
952	southern North Sea. In: Tarling, D. H. & Turner, P. (eds) Palaeomagnetism and diagenesis
953	in sediments, Geological Society of London special publications, 151, 109-124.
954	Ussher, W. A. E 1876. On the Triassic rocks of Somerset and Devon. Quart Jour. Geol. Soc.,
955	32, 367-394.
956	Ussher, W. A. E 1913. The geology of the country around Newton Abbott: explanation of sheet
957	339. Geological Survey of Great Britain. HMSO, London.
958	Visscher, H. 1973. The Upper Permian of western Europe—a palynological approach to
959	chronostratigraphy. In: Logan A. & Hills L.V. (eds) The Permian and Triassic systems and
960	their mutual boundary. Can. Soc. Pet. Geol. Mem., 2, 200-219.
961	Warrington, G. & Scrivener, R. C. 1990. The Permian of Devon, England. Review Palaeobotany
962	Palynology, <b>66</b> , 263-272.
963	Watson, G. S & Enkin, R. J. 1993. The fold test in palaeomagnetism as a parameter estimation
964	problem. Geophysical Research Letters, 20, 2135-3137.
965	Wills, L. J. 1970. The Triassic succession in the central Midlands in its regional setting. Jour.
966	Geol. Soc. Lond., <b>126</b> , 225-283.

- 267 Zheng, L., Yang, Z., Tong, Y., & Yuan, W. 2010. Magnetostratigraphic constraints on two-stage
  268 eruptions of the Emeishan continental flood basalts. *Geochemistry, Geophysics,*269 *Geosystems*, 11, doi:10.1029/2010GC003267.
- Zijderveld, J. D. A. 1967. The natural remanent magnetisation of the Exeter Volcanic Traps
  (Permian, Europe). *Tectonophysics*, 4, 121-153.
- Underhill, J. R. & Stoneley, R. 1998. Introduction to the development, evolution and petroleum
  geology of the Wessex Basin. *In:* Underhill, J. R. (ed.) Development, Evolution and
  Petroleum Geology of the Wessex Basin, Geological Society, London, Special
  Publications, 133, 1-18.
- Utting, J., 1996. Palynology of the Ufimian and Kazanian Stages of Russian stratotypes, and their
   comparison to the Word and Road of Canadian Arctic. *In:* Esaulova, N. K. & Lozovsky,
- 978 V. R. (eds) Stratotypes and reference sections of the Upper Permian of regions of the
- 979 Volga and Kama Rivers. Izd. "Ecocentr", Kazan, 486-506.
- 980

#### 981 Figure Captions

982 Fig. 1. Sketch map of the Permian-Triassic in SE Devon. Inset a) shows the study location within

983 the UK, where grey=Lower Palaeozoic basement highs, dotted=Permian basins. CAB=Channel

984 Approaches Basin, LRS=Lizard-Rhenish Suture, RHZ=Rheno-Hercynian Zone, STZ= Saxo-

985 Thuringian Zone . In main map, numbers correspond to the sampling locations indicated in the

986 Supplementary Data. Adapted from Selwood *et al.* (1984), Edwards *et al.* (1997). Sampling

- 987 locations on coast indicated by  $\}$  and in-land as  $\blacksquare$ .
- 988

Fig. 2. The stratigraphy of the Permian- Triassic around Exeter and the SE Devon coasts. Based

990 on this work and Laming (1982), Selwood et al. (1984), Edwards & Scrivener (1999), Leveridge

991 *et al.* (2003). Thicknesses of the coastal units is based on Selwood *et al.* (1984), Laming (1982)

and this work. The chronology is based on Edwards *et al.* (1997), Edwards & Scrivenor (1999)

993 and this work. The Torbay Breccia Formation occurs west of the Stickepath fault zone (SFZ,

dashed in grey), and is divisible into an upper unit (the Paignton breccias, PB) probably

995 equivalent to the Oddicombe Breccia Fm , and a lower unit composed of several separate

996 breccias units. PTM=Petit Tor Member. Arrows indicate overstepping units.

997

998 Fig. 3. A) The erosional boundary between the Littleham Mudstone Fm (below) and the Budleigh 999 Salterton Pebble Beds Fm (photo courtesy of Richard Porter), B) Immature calcrete and 1000 calcretised rootlets, top part of Straight Point Sandstone Member. C) Erosional boundary of 1001 breccia (base of arrow) at base of the Littleham Mudstone Fm, Littleham Cove (photo courtesy 1002 of Ian West). Scale arrow height=1.5 m. D) Unconformable boundary (marked in white) between 1003 the Watcombe Fm and the Oddicombe Breccia Fm, Whitsands Bay, hammer for scale. E) Detrital 1004 opaques (black) and pigmentary haematite grain coating (in red), fluvial sandstone, Dawlish 1005 Sandstone Fm, Dawlish Station section. The right hand side opaque (a haematised rock fragment) 1006 shows compactional deformation from surrounding framework grains. F) Detrital opaque 1007 showing indentation due to compaction into the surrounding quartz grains. Pigmentary haematite 1008 rims not present at opaque-quartz boundary. Fluvial sandstone in Dawlish Sandstone Fm. Pore 1009 spaces in blue. E) and F) scale bar is  $100 \,\mu\text{m}$ .

1010

Fig. 4. Section logs and summary palaeomagnetic data (horizon polarity, demagnetisation
behaviour and specimen polarity) from sections in the Exeter Group. See Fig. 1 for location

details. Symbols for specimen polarity and behaviour are larger for better quality behaviour.
Ticks adjacent to logs are sampling levels. Sample numbers indicated, for data shown on other
figures or in the supplementary data.

1016

Fig. 5. Section logs and summary horizon magnetic polarity data for the stratigraphic section
between Lympstone (site 3 on Fig. 1) to the top of the Littleham Mudstone Fm at Budleigh
Salterton (Fig. 1). Bed numbers on the log for the Exmouth Mudstones and Sandstones Fm are
those of Selwood *et al.* (1984). The divisions in the Littleham Mudstone Fm are from this work
(detailed in the supplementary data). Ticks adjacent to logs are sampling levels. Sample numbers
indicated, for data shown on other figures or in the supplementary data.

1023

1024 Fig. 6. Isothermal remanent magnetisation curves (A, C, E) and thermal demagnetisation of

1025 orthogonal isothermal remanent magnetisation (IRM) in B, D, F, for representative specimens.

1026 The Two numbers after lithology type are the magnetic susceptibility in  $x10^{-5}$  SI and NRM

1027 intensity in mA/m, respectively. Specimen numbers are those shown on Figs. 4 and 5.

1028 sst=sandstone; TBF= Teignmouth Breccia Fm, EBF= Exe Breccia Fm.

1029

1030 Fig. 7. Anisotropy of magnetic susceptibility data for the Littleham Mudstone Fm (A, E, I), The

1031 Exmouth Mudstone and Sandstone Fm (B, F, J), aeolian sandstones in the Dawlish Sandstone

1032 and Teignmouth Breccia formations (C, G, K), and the various breccia units (D, H, I). A), B), C),

1033 D) are steroegraphic projections of the specimen  $K_{max}$  and  $K_{min}$  directions. E), F), G), H) are the

1034 AMS ellipsoid shape  $(T = [2(L_{int}-L_{min})/(L_{max}-L_{min})]-1;$  where  $L = Ln(K_i)$  and strength (P =

1035 K<sub>max</sub>/K<sub>min</sub>; Tarling & Hrouda, 1993), I), J), K), L), rose diagrams showing the directions of the

1036  $K_{max}$  axes (mirrored about 0-180 axis), indicating the preferred grain long-axis directions in the

1037 bedding plane. Ns=number of specimens.

1038

1039 Fig. 8. Representative demagnetisation data from: (A, B) the Littleham Mudstone Fm, (C, D)

1040 Exmouth Mudstone and Sandstone Fm, (E) Exe Breccia Fm, (F) Dawlish Sandstone Fm, (G)

1041 Teignmouth Breccia and (H) Watcombe Fm. A) Specimen L35, normal polarity (behaviour S1,

1042 ChRM 500-660°C), B) EM30-4, reverse polarity (behaviour T1, component A, 0-500°C), C) E20,

1043 normal polarity (behaviour S2, ChRM 600°C to origin), D) EL63, reverse polarity (behaviour T1,

1044 Component A, 0-300°C), E) EB8-1A, normal polarity (behaviour S2, ChRM 300-500°C & 540°C

1045 to origin), F) DS21-1, reverse polarity (behaviour T1, steps 500°C and above noisy due to

1046 thermal alteration), G) DS4-2, reverse polarity (behaviour S2, ChRM 500-650°C), H) WB1-4, 1047 reverse polarity (behaviour S1, ChRM 100-620°C, 680°C step shows thermal alteration). See 1048 Figs. 4, 5 for specimen locations.

1049

1050 Fig. 9. a) Stereographic projection of all ChRM directions, with mean of these directions

1051 indicated for the units from the Aylesbeare Mudstone Group. B) The progressive unfolding fold

1052 test of Watson & Enkin (1993), using the data from the Exeter Group; showing the change in

1053 Fisher k with unfolding (left) and a pseudo-sampling bootstrap (right) to estimate the 95%

- 1054 confidence interval on the percentage unfolding.
- 1055

1056 Fig. 10. A) Detailed magnetostratigraphic data for the stratigraphic section between the 1057 Lympstone sections (3 on Fig. 1) and Littleham Cove. i) Demagnetisation behaviour showing 1058 categorisation into good (S1) and poor (S3) ChRM line-fits; great circle fit quality range from 1059 good (T1) to poor (T3), and specimens with no Triassic magnetisation are indicated in the P/X 1060 column . ii) Interpreted specimen polarity quality, with those in the greyed column not assigned a 1061 polarity. Poorest quality in column headed '??'. iii) VGP latitude, with filled symbols for those 1062 specimens possessing an S-class ChRM, and unfilled symbols for specimens with T-class, great-1063 circle behaviour. Polarity column: white= reversed polarity, black =normal polarity, grey= 1064 uncertain, gap=X; half bar-width indicates a single useful specimen from this horizon. B)

1065

Detailed magnetostratigraphic data for the stratigraphic section between Littleham Cove and

1066 Budleigh Salterton (1 on Fig. 1). Column details as in A).

1067

Fig. 11. Summary magnetostratigraphic data for European Permian sections, compared to the 1068

1069 composite geomagnetic magnetic polarity timescale (GPTS) of Hounslow (submitted). Southern

1070 North Sea data for the Leman Sandstone Fm from Turner et al. (1999) and Lawton & Robertson

1071 (2003). Czaplinek, Piła and Jaworzna IG-1 well data based on Nawrocki (1997) and

1072 Słowakiewicz et al. (2009). Mirow well data from Menning et al. (1988) and Langereis et al.

1073 (2010), Schlierbachswald-4 and Everdingen 1 wells from Szurlies et al. (2003) and Szurlies

1074 (2013). Related Russian stages tied to the magnetostratigraphy from Hounslow (submitted).

1075 Conodont zones labelled with Guadalupian (G) and Lopingian (L) zonal codes from Jin et al.

1076 (2000) and Shen et al. (2010). Early Wuchiapingian carbon isotope excursions (CIE) and

1077 Kamura event duration from Chen et al. (2011), Isozaki et al. (2011).

JGS, SW England Permian 36

Type/ Location/ Unit	Dec (°)	Inc (°)	K	<b>α95</b> (°)	NI/Np	Reversal Test	$G_0/G_C(^{\circ})$	Plat (°)	Plong (°)	Dp/Dm (°)
Li	ittleham M	udstone I	<sup>7</sup> m							
Line fits <sup>\$</sup>	12.4	29.2	26.0	5.4	28/0	R-	11.9/10.4*	53.6	156.3	3.3/6.0
GC means <sup>+</sup>	10.3	29.2	24.1	4.1	28/25	Rc	11.2/11.2	54.0	159.6	2.5/4.5
Exmouth	Mudstone	and Sand	lstone F	m						
Line fits <sup>\$</sup>	14.0	27.1	22.2	4.3	52/0	Rc	11.3/13.3*	52.0	154.2	2.6/4.7
GC means <sup>+</sup>	14.2	29.0	18.2	3.8	52/27	Rb	2.4/10.0*	53.1	153.5	2.3/4.2
Dawlish S	andstone a	nd Exe B	reccia fr	ns						
Line fits <sup>\$</sup>	5.0	26.6	40.4	7.3	11/0	Rc	7.5/20.0	53.2	168.5	4.3/7.9
GC means <sup>+</sup>	359.4	32.7	8.1	11.3	11/12	Rc	9.6/11.9*	56.9	185.7	7.2/12.8
T	eignmouth	Breccia H	m							
GC mean <sup>+</sup>	174.8	-25.1	25.1	8.5	10/3	-	-	52.4	184.8	4.9/9.1
0	ddicombe	Breccia F	'n							
Shaldon and Maidencombe <sup>\$</sup>	191.4	-24.4	116	3.4	16/0	-	-	51.1	158.6	2.0/3.6
Watcombe	e Fm, basal	Oddicon	nbe Brec	cia						
Watcombe <sup>\$</sup>	173.4	-20.0	28.0	10.7	8/0	-	-	49.5	186.5	5.9/11.2
	Other u	inits and	data							
Knowle Sandstone <sup>\$</sup>	195	-17	6842	3	2/0	-		-	-	-
Exeter Volc. Fm <sup>1</sup>	198	-25	23	6.5	23/0	-		49.5	148.5	-3.8/7.0
Exeter Volc. Fm <sup>2</sup>	189	-19	29	10	9/0	-		48	163	-5.4/10.4
Exeter Gp sediments <sup>2</sup>	188	-14	24	26	3/0	-		-	-	
	Group	means								
Aylesbeare Mudstone Gp	13.5	27.8	23.6	3.3	80/0	Rb	6.7/7.4*	52.5	154.9	2.0/3.6
Exeter Gp sediments	3.3	24.8	35.4	3.6	45/0	Rb	5.5/10.0*	52.4	171.2	2.1/3.9

Table 1. Directional means (with tectonic correction), reversal tests and VGP poles. <sup>+</sup>=great circle combined mean using method of McFadden & McElhinney (1988). <sup>\$</sup>=conventional Fisher mean. Nl=number of specimens using with fitted lines, and Np =number of specimens with great circle planes used in the determining the mean direction.  $\alpha_{95}$ , Fisher 95% cone of confidence. k, Fisher precision parameter. G<sub>0</sub> is the angular separation between the inverted reverse and normal directions, and Gc is the critical value for the reversal test. In the reversal test the Go/Gc values flagged with \* indicate common K values, others not flagged have statistically different K-values for reverse and normal populations, in which case a simulation reversal test was performed. Plat and Plong are the latitude and longitude of the mean virtual geomagnetic pole. <sup>1</sup>From Zijderveld (1967); <sup>2</sup> from Cornwall (1967)



#### Fig.2 **Budleigh Salterton to Exeter Torquay coastline** Anisian sic Sherwood Sandstone (~160m) Otter Sandstone Fm (~250m) ഗ Group σ <del>. . . . . . . . . . .</del> Olenek-(20-30m) Budleigh Salterton Pebble Beds Fm (30m) ian 7 Ma hiatus Wuchia-Aylesbeare (200-220m) Littleham Mudstone Fm (200-220m) Mudstone pingian Group Straight Point Member Capitan-Exmouth Sandstone and Mudstone Fm (~250m) ian (200m) Langstone Mb Exe Breccia Fm (85-450?m) Kenton Mb g d **Dawlish Sandstone Fm** (120-160 m) Wordian (100-120m) 7 0 Teignmouth Breccia Fm Ε (120-800 m) Heavitree Monkerton വ Breccia Fm Fm (135-300m) (0-40m) Oddicombe Breccia Fm <u>ـــ</u> Roadian (130-360m) Alphington Breccia Fm Φ Φ (160-240m) **(**10 - 60m) Whipton Fm ۵ PTM Watcombe Fm × 9 Ma hiatus Gzhelian(?) to (60-120m) Exeter (~20 m) Ш Artinskian Volcanics Knowle Sandstone Fm **Torbay Breccia Fm** (200 - 480 m) Carboniferous? Variscan basement



+Base Littleham Fm Breccia (Sylvie?) +Watcombe/Oddicombe boundary (Deryck)

+Micrograph 1 (mwh) +Micrograph 2 (mwh)











+~ 70 m of EBF







	Class	Ns	95	
e o	S1	23	5.7	2.1
ear Pn€	S2	28	10.6	2.4
stc ou	S3	29	13.9	3.2
Gr	T1	21	18.1	2.9
₹₹	T2	21	20.7	2.2
ا	T3	24	19.8	2.9
D)				
/	S1	31	3.2	2.9
гd	S2	5	5	2.9
ete ou	S3	8	9.6	3.8
Щ	T1	7	17.3	3.2
	T2	2	8.5	2.5
	T3	13	16.5	4.0
Exeter <b>(</b> Ayle Group G	T1 T2 T3 S1 S2 S3 T1 T2 T3	21 21 24 31 5 8 7 2 13	18.1 20.7 19.8 3.2 5 9.6 17.3 8.5 16.5	2.9 2.2 2.9 2.9 3.8 3.2 2.5 4.0













## End of the Kiaman Superchron in the Permian of SW England: Magnetostratigraphy of the Aylesbeare Mudstone and Exeter groups.

Mark W. Hounslow, Gregg McIntosh, Richard A. Edwards, Deryck Laming, Vassil Karloukovski

# Supplementary Data

The data here is composed of:

Table s1: Sampling site details, and mean magnetic properties.

Figure s1. The new bed-division of the Littleham Mudstone Fm, placed onto the photographs of the cliff outcrops.

Figure s2. Rose diagrams of the AMS Kmax axes, in the Aylesbeare Mudstone Group, against the sediment logs of the section.

Figure s3. The spatial variation in the AMS Kmax axes, of all Permian-Triassic west of Sidmouth, placed onto their sampling sites, and the palaeocurrent directions inferred from the sedimentology.

Figure s4. Component A data, and details of demagnetisation characteristics.

Figure s5. Additional rock magnetic data, pertaining to magnetic mineralogy

Figure s6. Petromagnetic data for all palaeomagnetic specimens.

Figure s7. Virtual geomagnetic pole data for Permian Europe, and a discussion of how the new data here fits with this data.

References.

Section	No. on Fig 1	Grid ref	Lat/long	Bedding strike/dip	N <sub>H</sub>	J <sub>0</sub> (x10 <sup>-3</sup> A/m)	κ <sub>lf</sub> (x10 <sup>-5</sup> SI)	Formation/unit
Budleigh Salterton to Littleham Cove	1	SY040802 to SY063817	50.622N: - 3.342W	302/5	38	3.2	2.1	Littleham Mudstone Fm (LMF)
Straight Point to Maer	2	SY040802 to SY011799	50.608N: - 3.376W	340/5 to 013/8	60	4.0	1.35	Exmouth Mudstone and Sandstone Fm (EMSF), base of LMF
Sowden Lane to Lympstone Harbour	3	SX990836 to SX988842	50.638N: - 3.441W	321/9 to 315/8	10	2.1	3.3	Base of EMSF, top of Exe Breccia Fm
Clyst St Mary Garage	4	SX976910	50.710N: - 3.454W	045/5	2	2.4	0.68	Dawlish Sandstone Fm
Bishops Court Quarry	5	SX965915	50.711N: - 3.465W	045/5	6	0.74	0.57	Dawlish Sandstone Fm
Langstone Rock	6	SX980779	50.598N: - 3.445W	293/8	6	2.8	7.6	Exe Breccia Fm
Dawlish Station	7	SX964767	50.580N: - 3.465W	293/10	5	1.4	2.7	Teignmouth Breccia Fm
Coryton Cliff and Cove	8	SX962762	50.577N: - 3.468W	319/10	8	1.5	1.9	Teignmouth Breccia Fm
Holcombe Beach	9	SX957746	50.562N:- 3.475W	342/12	2	2.4	5.1	Teignmouth Breccia Fm
Ness Point to Bundle Head	10	SX941720 to SX937712	50.533N: - 3.500W	355/0 to 295/5	6	7.2	9.8	Oddicombe Breccia Fm
Maidencombe Beach	11	SX928684	50.505: - 3.514	275/10 to 346/6	5	9.3	7.0	Oddicombe Breccia Fm
Whitsand to Watcombe Beachs	12	SX927674	50.496: - 3.515	315/41 to 288/25	5	5.8	6.4	Watcombe Fm and Oddicombe Breccia Fm
West Sandford	Not on Fig. 1	SS811027	50.812N: - 3.689W	105/14	1	10.8	12.1	Knowle Sandstone Fm

Table. s1. Section and site details sampled, and average magnetic properties of the samples. NH number of sampled horizons, Jo= Initial natural remanent magnetisation intensity, klf= low frequency magnetic susceptibility. Most samples were collected from the sea cliffs, with those from Bishops Court Quarry from a working quarry (Table s1). Those from Dawlish Station and West Sandford were from small cuttings. Samples from below bed B to the base of bed A (Fig. 5), in the Exmouth Mudstone and Sandstone Fm, were from foreshore exposures and sea-cliffs, east of Exmouth. In the breccia units, suitable units for palaeomagnetic sampling were thin, discontinuous sandstone and mudstone beds.



Fig.s1. Annotated photos of the cliff between Littleham Cove (panel i) and Budleigh Salterton (panel iv), indicating the bed and division sub-division of the Littleham Mudstones Formation. The three sub-divisions are a lower division A, mid division B, and an upper division C. The oldest part of the division A is exposed on the west side of the Littleham Cove fault. The W-E correlation across the Littleham Cove fault is not entirely clear, but the three sandstone beds (bed -1 in division A) may correlate to the upper-most units exposed west of the fault in the cliff adjacent to the path down the cliff. The succession is interrupted by a number of small landslips, which make the stratigraphy difficult to follow at beach level, without the photographs. The full succession can be examined by using the headwall scars behind the landslips.





Impersistant pale beds between A.3 and A.4

Fig.s1. Panels iii) and iv)







Fig.s2. K<sub>max</sub> axis directions for samples from the Aylesbeare Mudstone Group. The directions have been mirrored about the 0-180 axis. The  $K_{max}$  axes directions illustrate the similarity in ENE or easterly flow directions between the Exe Breccia and the lower part of the Exmouth Mudstone and Sandstone Fm. The directions in the upper part of the Group are more variable, but likely indicate a more NE direction of sediment transport. The SE-NW Kmax axes trends seen in the mudstones above bed G, and particularly in the Littleham Mudstones Fm may represent NW directed wind transport of the clay and silt in these units.



Fig.s3. Summary of transport directions and  $K_{max}$  axis directions for samples from the entire Permian, and Triassic successions west of Budleigh Salterton. Inset shows data for aeolian units, from the Dawlish Sandstone Fm (from Jones 1992, Selwood *et al.* 1984). The Kmax directions have been mirrored about the 0-180° axis (fluvial transport data from Laming 1966; Henson 1971; Selwood *et al.* 1984; Smith & Edwards 1991). The Triassic AMS data (Otter Sandstone, Budleigh Salterton Pebble Beds, BSPB) is from the samples described by Hounslow & McIntosh (2003). The Exeter Group below the Dawlish Sandstone Fm typically has bi-modal groups inferred to display ENE to easterly fluvial transport and northerly transport. Its possible in some of the sandstone and mudstones, the northerly trend may be a wind-transport direction, like seen in the aeolian units in the Dawlish Sandstone. Fluvial units in and above the Dawlish Sandstone Fm time to the lower Aylesbeare Mudstone Group, show strong ENE to easterly fluvial transport. In the Littleham Mudstone Fm directions may be transitional to the northerly transport clearly seen in the overlying BSPB and Otter Sandstone.



Fig.s4. Cumulative unblocking spectra for, a), b) the component A directions, from a) the Aylesbeare Mudstone Group, b) Exeter Group. C) The characteristic remanence ranges divided into stratigraphic groups. D) Statistics relating to the demagnetisation class, the number of specimens (Ns), the average  $\alpha_{95}$  of the line (S-class) or plane (T-class) principle component fits, and the average excess standard deviation ( $\rho$ ).



Fig.s5. A,b) Representative thermomagnetic data, measured using the vertical field translation balance (red=heating, blue= cooling). A subtle change in slope is seen at temperatures less than 200°C, which disappears when the estimated paramagnetic contribution is subtracted from the curves. C), d) log-Gaussian isothermal remanent magnetisation coercivity distributions (Kruiver *et al.* 2001) for selected samples, showing the fitted low field (component-1, c)) and high field components (component 2, d)). IRM data obtained with backfield data. Specimen WB1-6 and MB5-5 have higher coercivities (Bcr\* of 741-851 and 63 mT for the 2 components) than EM22-11 and EM30-14 (Bcr\* of 407-417 and 32 mT). Furthermore, these former 2 specimens had wider high coercivity distributions (dispersion parameter of 0.40-0.45 as compared to 0.32 for EM22-11 and EM30-14).





Fig. s7. Stable-Europe virtual geomagnetic poles (VGP) and their comparison to the VGP data from the latest Carboniferous, Permian and early-mid Triassic of the UK. Confidence cones for the units in SW England shown in grey, no confidence cones shown for other data. Each of the plots (A to E) shows the average European VGP path from Torsvik & Cocks (2005) labelled in Ma increments. (A) Lower and Middle Triassic sedimentary units, with Sherwood Sandstone Fm pole from Hounslow & McIntosh (2003), B) Permian sedimentary VGP data younger than the end of the KRPS (i.e. mid-late Wordian and younger). T=data from Taylor *et al.* (2009), G=from Gialanella *et al.* (1997). C) Sedimentary VGP data near the end of the KRPS. D) Lower Permian and latest Carboniferous sedimentary poles. E) Permian and latest Carboniferous volcanic and igneous-based VGP poles. MV=igneous units from the Scottish Midland Valley, WS= Whin sill data (from Liss *et al.* 2004). Data mostly from the Global Palaeomagnetic Database (http://www.ngu.no/dragon/), with newer data from Burov *et al.* (1998), Chen *et al.* (2006), Diego-Orozco *et al.* (2002), Nawrocki *et al.* (2008) and Bazhenov *et al.* (2008). In C) and D) rotated poles are filled circles.

## Discussion of VGP data in Figure s7

Creer (1957), Zijderveld (1967) and Cornwall (1967) presented palaeomagnetic data for the Exeter Volcanic Rocks, around Exeter and further north in the Crediton Trough. Cornwall (1967) also performed reconnaissance sampling of the successions in this study, and from the underlying Torbay Breccia Fm. Whilst Zijderveld and Cornwall did use AF demagnetisation, and Cornwall in addition used thermal demagnetisation, relatively few of the sites measured by Cornwall (1967) had structural corrections, whereas the mean direction determined by Zijderveld utilised tilt corrections (Table 1). Cornwall's (1967) mean direction for the Exeter Volcanic Rocks has overly shallows inclination due to inclusion of some southerly-directed magnetisations with positive inclinations, probably due to incomplete demagnetisation of the specimens. The mean VGP pole of Zijderveld (1967) falls towards the end of the late Carboniferous to Permian European APWP path (Fig. s7e), which corresponds well to the range of VGP's from other late Carboniferous to Lower Permian volcanic and igneous units (Fig. S7e).

The sites from the Teignmouth Breccia and from Watcombe Cove have a significant number of specimens having declinations east of south (Table 1; Fig. 9a). The result of this is a more southerly mean, which does however, reflect new data acquired from the Russian Platform by Taylor *et al.* (2009) and Gialanella *et al.* (1997), which are clearly separated (Fig. s7b) from the bulk of previous Russian data obtained from successions younger than the Kiaman Superchron (i.e. Molostovsky 1983; Burov *et al.* 1998). Bazhenov *et al.* (2008) have discussed the problems of this new data as either due to mis-orientation, or vertical axis rotations, hitherto undetected in the eastern part of the Russian platform. Vertical axis rotations of up to 30° have also been inferred for Lower Permian sediments (Fig. s7c) in basins in France and Germany (Diego-Orozco *et al.* 2002; Chen *et al.* 2006), where there are clearer, strike-slip-related tectonic mechanisms to produce this. There are insufficient data in this study to attempt an answer to this for the UK successions, but the similarity in age and tectonic setting of these European basins, south of the Variscan Front infers a common geological or geomagnetic origin for these outlier VGP directions, warranting further investigation, beyond the scope of this study.

### **Supplementary Information References**

- Bazhenov, M.L., Grishanov, A.N., Van der Voo, R. & Levashova, N.M. 2008. Late Permian palaeomagnetic data east and west of the Urals. *Geophys. J. Int.* **173**, 395-408.
- Burov, B. V., Zharkov, I. Y., Nurgaliev, D. K., Balabanov, Yu. P., Borisov, A. S. & Yasonov, P. G. 1998.
  Magnetostratigraphic characteristics of Upper Permian sections in the Volga and the Kama areas.
  In: Esaulova, N. K., Lozonsky, V. R., Rozanov, A. Yu. (eds). *Stratotypes and reference sections of the Upper Permian in the regions of the Volga and Kama Rivers*. GEOS, Moscow, 236-270.
- Cornwall, J. D. 1967. Palaeomagnetism of the Exeter Lavas, Devonshire. Geophys. *Journal Royal Astro. Soc.*, **12**, 181-196.
- Chen Y., Henry, B. Faure, M. Becq-Giraudon, J-F, Talbot, J-Y. Daly, L. & Le Goff, M. 2006. New Early Permian paleomagnetic results from the Brive basin (French Massif Central) and their implications for Late Variscan tectonics. *Int. J Earth Sci.*, DOI 10.1007/s00531-005-0010-5.
- Creer, K. M. 1957. The Natural Remanent Magnetization of Certain Stable Rocks from Great Britain. *Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences*, **250**, 111-129.
- Diego-Orozco, A., Chen,Y., Henry, B. & Becq- Giraudond, J-F. 2002. Paleomagnetic results from the Permian Rodez basin implications: the Late Variscan tectonics in the southern French Massif Central. *Geodynamica Acta*, **15**, 249260.
- Gialanella, P.R Heller, F., Haag, M. Nurgaliev, D. Borisov, A., Burov., B. Jasonov, P., Khasanov, D. Ibraginov, S. & Zharkov, I. 1997. Late Permian magnetostratigraphy on the eastern Russian Platform. *Geologie en Mijnbouw* **76**, 145-154.
- Henson, M.R. 1971. *The Permo-Triassic Rocks of South Devon*. Unpublished Ph.D. Thesis, University of Exeter.
- Hounslow, M.W. & McIntosh, G. 2003. Magnetostratigraphy of the Sherwood Sandstone Group (Lower and Middle Triassic): South Devon, U.K.: Detailed correlation of the marine and non-marine Anisian. *Palaeogeogr. Palaeoclimat. Palaeoecol.*, **193**, 325-348.
- Jones, N.S. 1992. Sedimentology of the Permo-Triassic of the Exeter area, S.W. England. British Geological Survey Technical Report, WH/92/122R.
- Kruiver, P. P., Dekkers, M. J. & Heslop, D. 2001. Quantification of magnetic coercivity components by the analysis of acquisition curves of isothermal remanent magnetisation. *Earth and Planetary Science Letters*, **189**, 269-276.
- Laming, D. J. C. 1966. Imbrications, palaeocurrents and other sedimentary features in the Lower New Red Sandstone, Devonshire, England. *Journal of Sedimentary Petrology*, **36**, 940-959.
- Liss, D., Owens, W. H. & Hutton, D. H. W. 2004. New palaeomagnetic results from the Whin Sill complex: evidence for a multiple intrusion event and revised virtual geomagnetic poles for the late Carboniferous for the British Isles. *Journal of the Geological Society, London*, **161**, 927-938.
- Molostovsky, E. A. 1983. *Paleomagnetic stratigraphy of the eastern European part of the USSR*. University of Saratov, Saratov [In Russian].
- Nawrocki, J. Fanning, M. Lewandowska, A., Polechoñska, O. & Werner, T. 2008. Palaeomagnetism and the age of the Cracow volcanic rocks (S Poland). *Geophys. J. Int.*, **174**, 475-488.
- Selwood, E. B. Edwards, R. A., Simpson, S., Chesher, J. A. & Hamblin, R. A. 1984. Geology of the country around Newton Abbot. Memoir for 1:50,000 geological sheet 339, British Geological Survey, HMSO, London.
- Smith, S.A. & Edwards, R.A. 1991. Regional sedimentological variations in Lower Triassic fluvial conglomerates (Budleigh Salterton Pebble Beds), southwest England: some implications for palaeogeography and basin evolution. *Geological Journal*, **26**, 65-83.
- Taylor, G. K., Tucker, C., Twitchett, R. J., Kearsey, T., Benton, M. J., Newell, A. J. & Tverdokhlebov, V. P. 2009. Magnetostratigraphy of Permian/Triassic boundary sequences in the Cis-Urals, Russia: No evidence for a major temporal hiatus. *Earth and Planetary Science Letters*, **281**, 36-47.
- Torsvik T. H. & Cocks L. R. M. 2005. Norway in space and time: A centennial cavalcade. Norwegian *Journal of Geology*, **85**, 73-86.
- Zijderveld, J. D. A. 1967. The natural remanent magnetisation of the Exeter Volcanic Traps (Permian, Europe). *Tectonophysics*, **4**, 121-153.