

Camouflage and interception: how pathogens evade detection by intracellular nucleic acid sensors

Leonie Unterholzner  and
Jessica F. Almine 

Division of Biomedical and Life Sciences,
Faculty of Health and Medicine, Lancaster
University, Lancaster, UK

Summary

Intracellular DNA and RNA sensors play a vital part in the innate immune response to viruses and other intracellular pathogens, causing the secretion of type I interferons, cytokines and chemokines from infected cells. Pathogen RNA can be detected by retinoic-acid inducible gene I-like receptors in the cytosol, whereas cytosolic DNA is recognized by DNA sensors such as cyclic GMP-AMP synthase (cGAS). The resulting local immune response, which is initiated within hours of infection, is able to eliminate many pathogens before they are able to establish an infection in the host. For this reason, all viruses, and some intracellular bacteria and protozoa, need to evade detection by nucleic acid sensors. Immune evasion strategies include the sequestration and modification of nucleic acids, and the inhibition or degradation of host factors involved in innate immune signalling. Large DNA viruses, such as herpesviruses, often use multiple viral proteins to inhibit signalling cascades at several different points; for instance herpes simplex virus 1 targets both DNA sensors cGAS and interferon- γ -inducible protein 16, as well as the adaptor protein STING (stimulator of interferon genes) and other signalling factors in the pathway. Viruses with a small genome encode only a few immunomodulatory proteins, but these are often multifunctional, such as the NS1 protein from influenza A virus, which inhibits RNA sensing in multiple ways. Intracellular bacteria and protozoa can also be detected by nucleic acid sensors. However, as the type I interferon response is not always beneficial for the host under these circumstances, some bacteria subvert, rather than evade, these signalling cascades for their own gain.

Keywords: DNA sensing; immune evasion; interferon; intracellular pathogens; retinoic-acid inducible gene I-like receptors; viruses.

doi:10.1111/imm.13030

Received 13 August 2018; revised 24

October 2018; accepted 26 November 2018.

Correspondence: Leonie Unterholzner, Division of Biomedical and Life Sciences, Faculty of Health and Medicine, Lancaster University, Lancaster LA1 4YQ, UK.

Email: l.unterholzner@lancaster.ac.uk

Senior author: Leonie Unterholzner

Introduction

Our innate immune system provides a rapid, if relatively non-specific, response to invading pathogens. The cornerstones of innate immunity are the pattern recognition receptors (PRRs), which detect pathogen-associated molecular patterns (PAMPs) during infection. PAMPs are key molecular features shared by classes of pathogens that are usually absent from healthy cells, and include, for instance, bacterial or fungal cell wall components, or viral nucleic acids that would not usually be found in the host cell. PAMPs can be recognized *in trans* when phagocytes detect engulfed pathogens or debris from infected cells via Toll-like receptors on their cell surface or inside

endosomes. In addition, many cell types can also recognize when they have been infected themselves by an intracellular pathogen using intracellular PRRs that constantly survey the cell's contents for signs of infection. Intracellular PRRs raise the alarm either after sensing the presence of PAMPs directly, or after detecting the indirect effects of infection or tissue damage as danger-associated molecular patterns.¹

Intracellular nucleic acid sensors are PRRs that can perform both those functions: they can recognize pathogen-derived DNA or RNA species inside the cell, and also detect nucleic acids as danger signals, for instance when infection leads to the leakage of mitochondrial DNA or RNA into the cytosol. Intracellular nucleic acid receptors

are particularly important in the detection of viruses, which lack many other conspicuous features such as the unique cell wall components of bacteria or protozoa. However, nucleic acids are starting to emerge as key mediators during the innate immune response to a range of pathogens, including intracellular bacteria and protozoa.

In this review, we will focus on the intracellular DNA and RNA sensors that induce a transcriptional response leading to the production of interferons, cytokines and chemokines. The rapidly inducible transcriptional programme is a key factor in the early elimination of pathogens and for the initiation of appropriate local immune responses that ultimately determine the outcome of an infection. For this reason, it can be assumed that all pathogens need to evade or subvert the host's innate immune responses to be able to establish an infection. Here, we will describe the strategies that intracellular pathogens use to hide from the cell's nucleic acid receptors, and to disable or subvert their function. Ultimately, the activation of signalling cascades by nucleic acid sensors and many other PRRs converges on the activation of transcription factors such as nuclear factor- κ B (NF- κ B) and interferon regulatory factor 3 (IRF3). Many pathogens also block innate immune responses at the level of these downstream signalling nodes, or by inhibiting the function of interferons and cytokines after they have been secreted, and we refer the reader to further reviews on this topic.²⁻⁴

Detection of pathogens by RIG-I-like receptors

The RIG-I-like receptors (RLRs) retinoic-acid inducible gene I (RIG-I) and melanoma differentiation-associated protein 5 (MDA5) recognize exogenous RNA in the cytosol of infected cells. RIG-I detects short stretches of dsRNA as well as RNA species containing a triphosphate or diphosphate moiety at their 5' end, and MDA5 recognizes long dsRNA molecules and higher-order aggregates of viral dsRNA.⁵ Although the cytosol contains abundant cellular RNA molecules, these are not detected by RLRs. Messenger RNAs contain an m7-guanosine cap at their 5' end, tRNAs are cleaved to generate a 5' monophosphate, and both tRNAs and rRNAs contain modifications that prevent recognition by RIG-I.⁵ Double-stranded RNA regions, e.g. in stem loops of endogenous RNA molecules, are usually short and contain frequent mismatches, further enhanced by the conversion of adenosine to inosine (A→I editing) mediated by adenosine deaminase acting on RNA 1 (ADAR1).⁶ In this way, RLRs are able to detect exogenous RNA molecules that may be present during infection in the presence of an abundance of cellular RNA in the cytosol.

Following detection of specific RNA ligands in the cytosol, RIG-I and MDA5 undergo post-translational

modifications such as dephosphorylation and polyubiquitylation⁷ and signal via their caspase activation and recruitment domains (CARD) to the CARD-containing adaptor mitochondrial anti-viral signalling protein (MAVS), nucleating the assembly of prion-like MAVS filaments.⁸ This results in the MAVS-mediated activation of the transcription factors IRF3 and NF- κ B, which co-operate in the activation of the interferon- β promoter. RLR signalling is further regulated by the addition of K48-linked and K11-linked ubiquitin chains, which mediate the degradation of RIG-I and MAVS.^{7,9} Termination of RLR signalling is also achieved by RIG-I phosphorylation preventing the ubiquitination of CARDs¹⁰ and MAVS dephosphorylation, which impedes interaction with downstream signalling proteins.¹¹ To prevent spurious activity, RIG-I and MAVS exist in an autoinhibitory state facilitated by the phosphorylation and sequestration of CARDs in RIG-I^{12,13} and the repression of signalling motifs by adjacent regions in MAVS.¹⁴ Negative regulation of signalling is important to ensure that an innate immune response is transient, and does not cause extensive tissue damage.

The RLRs have been implicated in the detection of many viruses, as well as other intracellular pathogens. Most viral RNA polymerases generate transcripts with 5' triphosphate ends and many viruses also generate dsRNA during their life cycle. Hence, both RIG-I and MDA5 are predicted to be able to detect the dsRNA genomes of reoviruses and the dsRNA replicative intermediates of the positive-strand ssRNA viruses of the *Flaviviridae* and *Coronaviridae* families.¹⁵ RIG-I can also detect the short dsRNA panhandle structures containing a 5' triphosphate from negative-strand RNA viruses, such as influenza A virus (IAV) and Sendai virus,^{16,17} and RNA with 5' diphosphate ends, such as the genome of reoviruses.¹⁸ MDA5, on the other hand, is specifically required for the detection of picornaviruses, that contain protein-capped positive-stranded RNA genomes, and generate long, structured dsRNA intermediates that can be detected by MDA5, but not RIG-I.¹⁹⁻²² MDA5 also specifically detects ssRNA viruses of the *Caliciviridae* family, such as norovirus.^{23,24} Several DNA viruses are also sensed by the RLRs, including herpes simplex virus 1 (HSV-1), adenovirus, modified vaccinia virus Ankara and Epstein-Barr virus (EBV).²⁵⁻²⁹ This is thought to be the result of the convergent transcription of adjacent open reading frames generating dsRNA, and the transcription of structured RNAs by viral polymerases. RNA sensing can also contribute to the detection of intracellular bacteria, including *Legionella pneumophila*, *Salmonella typhimurium* and *Shigella flexneri*.³⁰⁻³² Furthermore, RNA sensors can also be activated by cellular RNA as a positive feedback loop during infection: RNA fragments generated by the interferon-stimulated gene, RNaseL, unmasked ribosomal RNA transcripts or mitochondrial RNA that leaks into the cytosol can also be detected by RLRs.^{25,33,34}

Detection of exogenous DNA and cyclic dinucleotides

Many mammalian cell types can also detect exogenous DNA. Bacterial, viral or synthetic dsDNA, and even dsDNA isolated from mammalian cells, can be sensed, if it gains access to the cytosol and is over 40–50 bp in length.⁵ The key sensor of cytosolic DNA that induces an interferon response is the enzyme cyclic GMP-AMP synthase (cGAS), which upon dsDNA binding catalyses the production of the second messenger 2'3'-cyclic-GMP-AMP (cGAMP).³⁵ cGAMP then binds to the adaptor protein stimulator of interferon genes (STING), causing a conformational change in the STING dimers.³⁶ This is linked to the activation of STING, its translocation from the endoplasmic reticulum to perinuclear signalling foci, and the association with TANK-binding kinase 1 (TBK1) and IRF3.^{36–38} TBK1 phosphorylates STING at serine 366, which, in analogy to MAVS signalling, leads to the recruitment of IRF3.³⁷ STING function is further regulated by palmitoylation, and modification with K63-, K48-, K11- and K27-linked ubiquitin chains, and degradation by autophagy.⁷

Additional DNA binding proteins have been proposed to function as DNA sensors.³⁹ As cGAS is essential for the response to cytosolic DNA in most systems investigated so far, any other proposed DNA sensors may synergise with cGAS as co-receptors or co-factors, in order to further amplify the response or to provide an additional layer of specificity. This has been shown for the DNA binding protein interferon- γ -inducible protein 16 (IFI16), which co-operates with cGAS in the activation of STING in human cells.^{40,41} In human cells, IFI16 may also be able to detect viral DNA in the nucleus, possibly by recognizing unchromatinized ('naked') stretches of DNA to distinguish viral DNA from our own chromatin.^{42–44} Another member of the PYHIN family, Absent in Melanoma 2 (AIM2), has also been shown to detect DNA.^{45–47} However, its downstream signalling diverges from IFI16 as it triggers inflammasome formation and interleukin-1 β production. Other DNA sensors include DNA-dependent activator of interferon-regulatory factors (DAI),⁴⁸ DNA-dependent protein kinase (DNA-PK)⁴⁹ DExD/H-box helicases DHX9 and DHX36,⁵⁰ DDX41⁵¹ and the double strand break repair endonuclease MRE11.⁵² Several of these proteins have been observed to be important for the induction of an interferon response after detection of cytosolic DNA, but their molecular role in the context of the cGAS–STING signalling axis remains to be determined.

The cytosolic DNA sensing pathway has been shown to be crucial for the detection of many different DNA viruses and retroviruses.^{53,54} Surprisingly, some RNA viruses such as Dengue virus also engage the DNA sensing machinery by causing mitochondrial damage, resulting in the release of mitochondrial DNA into the

cytosol.⁵⁵ STING-dependent DNA sensing has also been shown to be involved in the detection of DNA from protozoa such as *Plasmodium falciparum* and intracellular bacteria including *Mycobacterium tuberculosis* and *Listeria monocytogenes*.^{56–59}

During bacterial infection, the DNA sensing adaptor STING has a second function and can act as a PRR in its own right. STING can sense bacterial signalling molecules such as cyclic di-GMP and cyclic di-AMP.^{60,61} Like cGAMP, these cyclic dinucleotides can bind and activate STING and induce the production of type I interferons, which has been observed, for example, during infection with *Mycobacterium tuberculosis* and *Listeria monocytogenes*.^{61–63} Hence bacteria can activate a STING-dependent innate immune response by two different means, even though the resulting production of type I interferons is not always beneficial for the host, but instead can serve as an immune diversion strategy for some bacteria.^{63,64}

Immune evasion strategies employed by intracellular pathogens

Pathogens and their hosts co-evolve, and there are clear signs of selective pressure on host factors such as DNA and RNA sensors on the one hand and pathogen-encoded immune evasion proteins on the other.^{65–67} Immunomodulatory proteins and virulence factors are finely tuned to adapt to the host range and life cycle of the pathogen. The precise interplay between host and pathogen factors is often unique, and can differ in different host organisms, cell types and among even closely related pathogens. However, mapping these host–pathogen interactions at a molecular level can provide clues about the importance of host signalling cascades, can help us to understand changes in species specificity and virulence of emerging pathogens, and aid in the design of vaccine vectors. As PRR signalling pathways are being characterized in more molecular detail, the intricate counterstrategies employed by pathogens to evade recognition are also becoming apparent. Some general principles by which intracellular pathogens hide from detection or intercept the cell's signalling cascades are described below.

Sequestration of PAMPs

Pathogens that reside inside host cells are already sheltered from the immune system to some extent, by avoiding exposure to antibodies or complement components. However, due to the existence of intracellular PRRs, additional camouflage strategies are essential for the establishment of infection. Intracellular bacteria and protozoan parasites often co-opt vacuoles as a niche for replication, which prevents pathogen-derived nucleic acids and other PAMPs from being exposed to the cell's PRRs. Hence, detection of intracellular nucleic acids during bacterial

infection often requires the presence of bacterial secretion systems. For instance, the type VII secretion system ESX from *Mycobacterium tuberculosis* is responsible for the exposure of bacterial DNA to cGAS in macrophages, but also secretes virulence factors that interfere with host functions.^{56,68} Viruses require greater access to host factors for replication than bacteria, but also try to shield their nucleic acids from detection. Some RNA viruses, including Dengue virus and hepatitis C virus, sequester their genome and replication machinery in membrane-bound compartments, which function to create locally high concentrations of replication factors, as well as hiding viral RNA genomes from recognition by RIG-I.^{69,70} Human immunodeficiency virus type 1 carries out reverse transcription and replication of its genome inside its capsid, which contains selective pores to allow entry for nucleotides from the host cell, while shielding viral DNA from recognition by cGAS.⁷¹

Modification of viral RNA and DNA

Many pathogens aim to keep PAMP levels at a minimum: coronaviruses for instance quickly degrade any excessive dsRNA formed during the viral life cycle,⁷² and Group B streptococcus degrades cyclic di-AMP using an ectonuclease.⁷³ However, the production of nucleic acid PAMPs cannot be avoided altogether; thus, more sophisticated strategies of camouflage have evolved, so that the pathogen can blend into the cellular environment.

As viral RNA polymerases generate RNA species with a 5' triphosphate moiety, many viruses employ additional strategies to modify their RNA, so that it is not recognized efficiently by RIG-I. For instance, *Picornaviridae*, *Caliciviridae* and *Astroviridae* covalently attach a protein, Vpg, to the 5' end of viral RNAs, which prevents RIG-I-mediated recognition.¹⁵ Processing of the negative-strand ssRNA genome of members of the *Bornaviridae* and *Bunyaviridae* families involves the cleavage of the first nucleotide by a viral endonuclease to generate a monophosphate group at the 5' end.⁷⁴ Viruses that transcribe their mRNAs in the nucleus, including the DNA viruses of the *Herpesviridae* and *Papillomaviridae* families and retroviruses such as human immunodeficiency virus, co-opt the cell's own capping machinery to protect the 5' end of their mRNAs with a 7-methylguanosine cap, while others, including IAV, use 'cap snatching' to remove the capped 5' ends from cellular mRNAs and incorporate them into viral RNA transcripts.⁷⁵ Many viruses, including poxviruses and flaviviruses, encode their own capping enzymes that synthesize RNA caps that are indistinguishable from the mammalian cap structure.⁷⁵ Yellow fever virus in addition recapitulates the 2' O-methylation of the first RNA nucleotide next to the cap, which is found in cellular mRNAs and further decreases recognition by RIG-I.⁷⁶ Bacteria and protozoa also cap their mRNAs,

but not always with 100% efficiency. RNA derived from intracellular bacteria can activate the RNA sensing pathway if it reaches the cytosol, but it remains to be determined whether this is important during infection.⁷⁷

So far, it is not known whether DNA modifications could prevent recognition by cGAS or other DNA sensors. It has been proposed that IFI16 may be able to distinguish viral DNA from the cell's own genome by recognizing longer (>40 nucleotides) stretches of free DNA,^{43,44} and the detection of DNA by cGAS has also been shown to be length-dependent.⁷⁸ It remains to be tested whether the assembly of nucleosomes on the genomes of DNA viruses such as herpesviruses and papillomaviruses could be a strategy to limit their detection by intracellular DNA sensors.

Degradation and inhibition of host signalling factors

All intracellular pathogens encode virulence factors that interfere with the host's signalling cascades. Viral immunomodulatory proteins are usually expressed as immediate early genes, or are even part of the viral particle, so that they are able to inhibit the host's signalling cascades as soon as PAMPs are detected early in infection. Intracellular bacteria employ secretion systems to transport effector proteins through the vacuolar and bacterial membranes into the host cell, whereas protozoans can secrete virulence factors via exosomes or other membrane-bound vesicles. A vast repertoire of virulence factors that influence the cell's innate immune signalling cascades have been described, particularly in viruses, which often dedicate a large portion of their genome to interfering with the host's innate immune response. Although many viruses encode only a few proteins, these are often multi-functional and target several key host factors. Large DNA viruses, such as poxviruses and herpesviruses, encode a larger repertoire of dozens of immunomodulatory proteins that interfere with anti-viral signalling cascades in the cell at multiple points.^{79,80} Bacteria and protozoa encode thousands of genes, and so have an even greater capacity to interfere with host functions, but comparatively few immune evasion proteins have been described so far, with many pathogen-encoded effector proteins still uncharacterized.^{3,4,64}

Many pathogens have evolved similar strategies to inhibit the innate immune signalling cascades in the host cell: virulence factors act to either eliminate host signalling factors by degradation, sequester them or block their function in the signalling cascade. As we learn more about the regulation of innate immune signalling cascades that detect pathogens, more examples of counterstrategies employed by pathogens also emerge.

Degradation and inhibition of RNA sensors

Since the discovery of the intracellular RNA sensing system, many immune evasion proteins that target RLRs or

MAVS have been identified (see Fig. 1), mostly in viruses with an RNA genome, for which RIG-I and/or MDA5 are the key PRRs during infection. Some viral proteases cleave RNA sensors, with several picornaviruses using their 3C protease to cleave RIG-I, and the 2A protease to cleave MDA5.^{81–83} The leader protease L^{Pro} from foot-and-mouth-disease virus also cleaves Lpg2, a co-factor for MDA5 activation.⁸⁴ Some viruses modify RLRs with K48-linked poly-ubiquitin chains, diverting the RNA sensors for degradation by the cell's proteasome. Toscana virus non-structural proteins and the rotavirus NSP1 protein cause the degradation of RIG-I,^{85,86} whereas West Nile virus NS1 targets both RIG-I and MDA5.⁸⁷ RIG-I mRNA expression and translation can also be inhibited during viral infection: for instance, EBV expresses the virus-

encoded microRNA miR-BART6-3p, which targets RIG-I mRNA,⁸⁸ and hepatitis B virus induces the cellular micro-RNA miR146a, which inhibits RIG-I expression.⁸⁹

Even under conditions where the expression of RIG-I and MDA5 is intact, the RNA sensors can be prevented from carrying out their signalling function by viral proteins that bind to them. The HSV-1 protein UL37 interacts with RIG-I and causes the de-amidation of its helicase domain, which renders RIG-I unable to sense RNA ligands.⁹⁰ Another HSV-1 protein, US11 inhibits RNA sensing by associating with the dsRNA binding protein PACT, which potentiates RNA-induced responses.^{91,92} PACT is also targeted by Influenza A virus NS1, the VP35 protein from Ebola virus and the 4a protein from Middle East respiratory syndrome coronavirus.^{93–95} Many of the PACT-




Figure 1. Schematic representation of the pathogen-derived molecules employed to evade the intracellular RNA sensing pathway. The RNA sensors RIG-I and MDA5 detect various RNA species, primarily those containing 5' di- or tri-phosphates or long dsRNA, respectively. Some viruses modify or mask their viral RNA, for instance by cleaving 5' phosphates, attaching a viral protein or 'cap snatching'. Pathogens also inhibit RIG-I activation via degradation or blocking accessory proteins (TRIM25 and Riplet), by direct binding with the CARD or C-terminal domains of RIG-I and by preventing RIG-I translocation to the mitochondria to block its interaction with MAVS. MDA5 is targeted in a similar fashion, often by degradation or inhibition by binding of virulence factors. Various viral proteins also interact with or degrade MAVS, which provides a further layer of evasion of the RNA sensing pathway. DENV, Dengue virus; EBV, Epstein-Barr virus; EBOV, Ebola virus; HBV, hepatitis B virus; hCMV, human cytomegalovirus; HCV, hepatitis C virus; HHV8, human herpesvirus 8; hMPV, human metapneumovirus; HSV-1, herpes simplex virus 1; IAV, influenza A virus; MERS, Middle Eastern respiratory virus; RSV, respiratory syncytial virus; SARS, severe acute respiratory syndrome; TOSV, Toscana virus; WNV, West Nile virus; ZIKV, Zika virus.

interacting proteins also bind dsRNA, so they probably shield the viral RNA from detection while at the same time inactivating PACT and RNA sensors.

The activity of RIG-I and MDA5 is tightly regulated in the cell: In the absence of infection, the CARD of the RNA sensors are kept in a phosphorylated state, and activation involves their dephosphorylation catalysed by protein phosphatase 1.⁹⁶ RIG-I activity is further enhanced by the assembly of K63-linked poly-ubiquitin chains catalysed by the E3 ubiquitin ligases tripartite motif containing 25 (TRIM25) and Riplet, which release RIG-I from its autoinhibited state.^{97,98} Given the importance of these post-translational modifications, it is not surprising that many viruses have evolved ways of inhibiting these regulatory mechanisms. For instance, measles virus uses an elaborate signalling strategy involving the activation of the C-type lectin DC-SIGN to inhibit protein phosphatase 1, thus keeping the RLRs phosphorylated and inactive.⁹⁹ The E3 ubiquitin ligase Riplet is inhibited by the multi-functional IAV NS1 protein¹⁰⁰ and is cleaved by the NS3/4A protease from hepatitis C virus.¹⁰¹ Multiple viruses also inhibit TRIM25, including again IAV NS1, the SARS coronavirus N protein, the V protein from various paramyxoviruses, the large tegument protein BPLF1 from EBV and the E6 protein from human papillomavirus 16.^{102–107} The subgenomic flavivirus RNA from Dengue virus can also bind and inactivate TRIM25 in an RNA sequence-specific manner.¹⁰⁸ RIG-I ubiquitylation by TRIM25 is also inhibited by the bacterial quorum sensing molecule cyclo(Phe-Pro) from the opportunistic pathogen *Vibrio vulnificus*. Cyclo(Phe-Pro) specifically binds to RIG-I and prevents it from being ubiquitylated by TRIM25,¹⁰⁹ possibly providing a link between bacterial infection and susceptibility to co-infection with viruses.

A multitude of viral immunomodulators prevent RIG-I and MDA5 from interacting with its adaptor protein MAVS, so blocking the nucleation event that allows MAVS to form higher-order signalling assemblies. Dengue virus NS3 binds 14-3-3 ϵ to prevent the translocation of RIG-I to MAVS.¹¹⁰ The interaction between RLRs and MAVS is also disrupted by the picornavirus 3C^{Pro} proteases, the Z and N proteins of some arenaviruses, NS2 from respiratory syncytial virus, NS1 from IAV, NSP1 from rotavirus and US11 from HSV-1.^{16,86,111–114}

MAVS is also targeted directly by many viruses. MAVS-interacting proteins include the metapneumovirus protein M2-2,¹¹⁵ the NS4A proteins of Zika and Dengue viruses,^{116,117} viral IRF1 from human herpesvirus 8¹¹⁸ and ORF9b from SARS coronavirus.¹¹⁹ MAVS is also diverted for proteasome-mediated degradation by the hepatitis B virus X protein,¹²⁰ and is cleaved by the hepatitis C virus protease NS3/4A, which releases it from the mitochondria and other intracellular membranes to disrupt signalling.^{121–124} The US9 protein from HSV-1 causes MAVS to leak from the mitochondria by disrupting the

mitochondrial membrane potential,¹²⁵ and the IAV protein PB1-F2 decreases mitochondrial membrane potential causing mitochondrial fragmentation.^{126,127} Disrupting mitochondrial physiology is a common mechanism to block MAVS signalling. Infection with Dengue and Zika viruses causes mitochondrial elongation, which also blocks MAVS function.¹²⁸ The sustained interferon production caused by MAVS signalling from the peroxisomes is inhibited by the MIA protein from human cytomegalovirus (hCMV) and the VP16 protein from HSV-1.^{118,129,130}

Although there are abundant examples of inhibition of RNA sensing by pathogens, and in particular viruses, there is one instance where a pathogen actually enhances signalling by RIG-I and MDA5. The food-borne pathogen *Salmonella typhimurium* uses its effector protein SopA, an E3 ubiquitin ligase, to ubiquitylate the host proteins TRIM56 and TRIM65, which in turn promote RIG-I and MDA5 signalling.^{131,132} This is an example of subversion, rather than evasion, of the interferon response, in line with the observation that the production of type I interferons can be advantageous to the pathogen, rather than the host, during some bacterial infections.⁶⁴

Degradation and inhibition of DNA sensors

Even though the cGAS/STING DNA sensing pathway was only discovered relatively recently, several viral and bacterial virulence factors that target this pathway have already been discovered, see Fig. 2. The nuclear DNA viruses hCMV, HSV-1 and Kaposi's sarcoma-associated herpesvirus all target several key DNA sensing factors to prevent the detection of their genomic DNA. HSV-1 uses its ICP27 protein to inhibit STING.¹³³ In addition, the E3 ubiquitin ligase ICP0 causes the degradation of IFI16 protein, and the virion host shutoff protein UL41 promotes the turnover of the mRNAs encoding IFI16 and cGAS.^{42,134,135} In addition, the HSV-1 virion protein VP22 interacts with cGAS and inhibits its enzymatic activity¹³⁶ and also interacts with AIM2 and blocks its oligomerization.¹³⁷ HSV-1 tegument protein UL37, which targets and deamidates RIG-I, also deamidates cGAS, impairing cGAMP synthesis.¹³⁸ The deployment of different virulence factors to target several components of the DNA sensing pathway may be necessary to block the pathway more efficiently, and/or to inhibit some non-overlapping functions of cGAS and IFI16 which have been observed during herpesvirus infection.¹³⁹ Analogously, several different hCMV proteins inhibit DNA sensing: pUL31 binds to cGAS and dissociates it from DNA,¹⁴⁰ pUL83 inhibits both cGAS and IFI16,^{141–143} pUL82 inhibits STING translocation,¹⁴⁴ IE2 causes STING degradation,¹⁴⁵ and US9 inactivates both MAVS and STING.¹²⁵ Furthermore, pUL83 binds AIM2 and facilitates the degradation of AIM2-driven inflammasomes.¹⁴⁶

A similar multi-pronged approach has also been observed for the gammaherpesvirus Kaposi's sarcoma-associated herpesvirus.^{147–150} STING is also inhibited by oncoproteins from the nuclear DNA viruses human adenovirus and human papillomavirus,¹⁵¹ and by vaccinia virus, a DNA virus that resides in the cytosol.¹⁵² Vaccinia virus also inhibits DNA-PK, using its virulence factor C16 to bind the Ku70/Ku80 subunits of the DNA-PK complex.¹⁵³ The viral inhibitor of RIP activation (vIRA) from mouse CMV targets another proposed DNA sensor, DAI, to prevent the induction of programmed necrosis upon viral infection.¹⁵⁴

Surprisingly, not only DNA viruses but also some RNA viruses have been found to antagonise STING – even though they do not produce DNA or cyclic dinucleotides during their replication cycle. It has been reported that

STING can function in a cGAS-independent manner during the detection of viral membrane fusion, for instance during infection with IAV.¹⁵⁵ This non-canonical STING signalling pathway is antagonized by the IAV fusion peptide.¹⁵⁶ Flaviviruses, enveloped viruses with a positive-stranded ssRNA genome, also inhibit STING-dependent DNA sensing. Viral NS2B proteases encoded by Dengue virus, Zika virus, West Nile virus and Japanese encephalitis virus cleave human STING protein, but not its mouse orthologue.^{157,158} The dengue virus NS2B co-factor also targets cGAS for degradation.⁵⁵ This prevents the activation of an innate immune response after the release of mitochondrial DNA, which occurs during Dengue virus infection,^{55,159} highlighting the breadth of pathogen classes that can be detected directly or indirectly by the DNA sensing pathway (Fig. 2).




Figure 2. Schematic representation of the pathogen-derived molecules employed to evade the intracellular DNA sensing pathway. cGAS is a sensor of cytosolic DNA and the main activator of the adaptor protein, STING. Pathogens target cGAS for degradation, impede cGAS from binding DNA or inhibit its catalytic activity. Pathogens also degrade cGAMP, the second messenger produced by cGAS, as well as bacterial cyclic dinucleotides. IFI16 contributes to the activation of the cGAS–STING pathway, and several viral proteins have been identified that inhibit its activation and promote its degradation. Other DNA sensors, such as DNA-PK, DAI and AIM2 are also targeted by viral factors that block DNA binding and activation of downstream pathways. Pathogens also hinder multiple aspects of STING function through proteolytic cleavage, degradation, blocking its translocation from the endoplasmic reticulum (ER) to ER-Golgi intermediate compartment (ERGIC) and preventing interaction with downstream signalling proteins, such as TANK-binding kinase 1 (TBK1). DENV, Dengue virus; hAd, human adenovirus; hCMV, human cytomegalovirus; HSV-1, herpes simplex virus 1; HPV18, human papillomavirus 18; JEV, Japanese encephalitis virus; mCMV, mouse cytomegalovirus; WNV, West Nile virus; ZIKV, Zika virus.

While the production of type I interferons is not always an effective strategy to limit bacterial infections, the activation of STING by bacterial DNA or cyclic dinucleotides can also promote autophagy to clear bacteria from the infected cells. For this reason, some intracellular bacteria also inhibit STING.⁶² For instance, the *Shigella* effector protein IpaJ inhibits STING translocation from the endoplasmic reticulum to endoplasmic reticulum–Golgi intermediate compartment,³⁸ and the *Yersinia* YopJ protein also blocks STING trafficking and causes its de-ubiquitylation.¹⁶⁰ *Mycobacterium tuberculosis* secretes the cyclic di-nucleotide phosphodiesterase CdnP (also known as Rv2837c), which degrades cGAMP and bacterial cyclic dinucleotides, thus inhibiting both DNA- and cyclic-dinucleotide-induced STING activation.¹⁶¹ Group A streptococcus, *Streptococcus pyogenes*, subverts, rather than evades, STING signalling: it uses its M protein to activate STING, resulting in the production of the anti-inflammatory cytokine interleukin-10 downstream of type I interferon signalling.¹⁶² In this way, the bacterium exploits the reciprocal antagonism between the interferon response and inflammation, and shapes the immune response to favour the pathogen, rather than the host.

Concluding remarks

All pathogens, having co-evolved with their hosts, employ elaborate strategies to strike a fine balance between the requirements of their own life cycle, the evasion of host defences, and effects on pathology for the host, which can influence further transmission and hence the evolutionary fitness of a pathogen. As in so many fields in biology, much can be learnt from insights into the host–pathogen interactions that interfere with our cells’ defences. Immunomodulatory proteins from pathogens can point us to their targets, which are often key signalling nodes that play a role during infection. Although immune evasion has long been studied in human cells and mouse models, the investigation of innate immune signalling in reservoir hosts such as bats or birds, is an emerging field of research. The variation of these interactions during pathogen evolution often underlies the ability of pathogens to cross species boundaries or to adopt new traits in transmission or virulence during emerging infections. Knowing more about the functions of individual immunomodulatory proteins is also crucial for the rational development of vaccine vectors, as for instance the removal of individual immune evasion factors can aid safety and immunogenicity of a live vaccine. It is becoming increasingly clear that nucleic acid PAMPs and their PRRs also perform key roles during sterile inflammation, autoimmunity and even cancer treatment, so the lessons we learn from our pathogens may ultimately have applications for the treatment and prevention of both infectious and non-infectious diseases that involve the immune system.

Acknowledgements

Our work is funded by the Medical Research Council (Career Development Award MR/K00655X/1), the European Commission (MC-CIG631718) and North West Cancer Research (CR-1140).

Disclosures

The authors declare no competing interests.

References

- Blander JM, Sander LE. Beyond pattern recognition: five immune checkpoints for scaling the microbial threat. *Nat Rev Immunol* 2012; **12**:215–25.
- Beachboard DC, Horner SM. Innate immune evasion strategies of DNA and RNA viruses. *Curr Opin Microbiol* 2016; **32**:113–9.
- Dos-Santos ALA, Carvalho-Kelly LF, Dick CF, Meyer-Fernandes JR. Innate immunomodulation to trypanosomatid parasite infections. *Exp Parasitol* 2016; **167**:67–75.
- Reddick LE, Alto NM. Bacteria fighting back: how pathogens target and subvert the host innate immune system. *Mol Cell* 2014; **54**:321–8.
- Schlee M, Hartmann G. Discriminating self from non-self in nucleic acid sensing. *Nat Rev Immunol* 2016; **16**:566–80.
- Lässig C, Hopfner K-P. Discrimination of cytosolic self and non-self RNA by RIG-I-like receptors. *J Biol Chem* 2017; **292**:9000–9.
- Chiang C, Gack MU. Post-translational control of intracellular pathogen sensing pathways. *Trends Immunol* 2017; **38**:39–52.
- Hou F, Sun L, Zheng H, Skaug B, Jiang Q-X, Chen ZJ. MAVS forms functional prion-like aggregates to activate and propagate antiviral innate immune response. *Cell* 2011; **146**:448–61.
- Xing J, Zhang A, Minze LJ, Li XC, Zhang Z. TRIM29 negatively regulates the Type I IFN production in response to RNA virus. *J Immunol* 2018; **201**:183–92.
- Nistal-Villán E, Gack MU, Martínez-Delgado G, Maharaj NP, Inn K-S, Yang H *et al.* Negative role of RIG-I serine 8 phosphorylation in the regulation of interferon- β production. *J Biol Chem* 2010; **285**:20252–61.
- Xiang W, Zhang Q, Lin X, Wu S, Zhou Y, Meng F *et al.* PPM1A silences cytosolic RNA sensing and antiviral defense through direct dephosphorylation of MAVS and TBK1. *Sci Adv* 2016; **2**:e1501889.
- Gack MU, Nistal-Villán E, Inn K-S, García-Sastre A, Jung JU. Phosphorylation-mediated negative regulation of RIG-I antiviral activity. *J Virol* 2010; **84**:3220–9.
- Saito T, Hirai R, Loo Y-M, Owen D, Johnson CL, Sinha SC *et al.* Regulation of innate antiviral defenses through a shared repressor domain in RIG-I and LGP2. *Proc Natl Acad Sci USA* 2007; **104**:582–7.
- Shi Y, Yuan B, Qi N, Zhu W, Su J, Li X *et al.* An autoinhibitory mechanism modulates MAVS activity in antiviral innate immune response. *Nat Commun* 2015; **6**:7811.
- Schlee M. Master sensors of pathogenic RNA – RIG-I like receptors. *Immunobiology* 2013; **218**:1322–35.
- Pichlmair A, Schulz O, Tan CP, Näslund TI, Liljestrom P, Weber F *et al.* RIG-I-mediated antiviral responses to single-stranded RNA bearing 5'-phosphates. *Science* 2006; **314**:997–1001.
- Rehwinkel J, Tan CP, Goubau D, Schulz O, Pichlmair A, Bier K *et al.* RIG-I detects viral genomic RNA during negative-strand RNA virus infection. *Cell* 2010; **140**:397–408.
- Goubau D, Schlee M, Deddouche S, Pruijssers AJ, Zillinger T, Goldeck M *et al.* Antiviral immunity via RIG-I-mediated recognition of RNA bearing 5'-diphosphates. *Nature* 2014; **514**:372–5.
- Kato H, Takeuchi O, Sato S, Yoneyama M, Yamamoto M, Matsui K *et al.* Differential roles of MDA5 and RIG-I helicases in the recognition of RNA viruses. *Nature* 2006; **441**:101–5.
- Pichlmair A, Schulz O, Tan CP, Rehwinkel J, Kato H, Takeuchi O *et al.* Activation of MDA5 requires higher-order RNA structures generated during virus infection. *J Virol* 2009; **83**:10761–9.
- Gitlin L, Benoit L, Song C, Cella M, Gilfillan S, Holtzman MJ *et al.* Melanoma differentiation-associated gene 5 (MDA5) is involved in the innate immune response to Paramyxoviridae infection *in vivo*. *PLoS Pathog* 2010; **6**:e1000734.
- Feng Q, Hato SV, Langereis MA, Zoll J, Virgen-Slane R, Peisley A *et al.* MDA5 detects the double-stranded RNA replicative form in picornavirus-infected cells. *Cell Rep* 2012; **2**:1187–96.

- 23 McCartney SA, Thackray LB, Gitlin L, Gilfillan S, Virgin HW, Virgin Iv HW *et al.* MDA-5 recognition of a murine norovirus. *PLoS Pathog* 2008; **4**:e1000108.
- 24 Chhabra P, Ranjan P, Cromeans T, Sambhara S, Vinjé J. Critical role of RIG-I and MDA5 in early and late stages of Tulane virus infection. *J Gen Virol* 2017; **98**:1016–26.
- 25 Chiang JJ, Sparrer KJM, van Gent M, Lässig C, Huang T, Osterrieder N *et al.* Viral unmasking of cellular 5S rRNA pseudogene transcripts induces RIG-I-mediated immunity. *Nat Immunol* 2018; **19**:53–62.
- 26 Rasmussen SB, Jensen SB, Nielsen C, Quartin E, Kato H, Chen ZJ *et al.* Herpes simplex virus infection is sensed by both Toll-like receptors and retinoic acid-inducible gene-like receptors, which synergize to induce type I interferon production. *J Gen Virol* 2009; **90**:74–8.
- 27 Minamitani T, Iwakiri D, Takada K. Adenovirus virus-associated RNAs induce type I interferon expression through a RIG-I-mediated pathway. *J Virol* 2011; **85**:4035–40.
- 28 Samanta M, Iwakiri D, Kanda T, Imazumi T, Takada K. EB virus-encoded RNAs are recognized by RIG-I and activate signaling to induce type I IFN. *EMBO J* 2006; **25**:4207–14.
- 29 Delaloye J, Roger T, Steiner-Tardivel Q-G, Le Roy D, Knaup Reymond M, Akira S *et al.* Innate immune sensing of modified vaccinia virus Ankara (MVA) is mediated by TLR2-TLR6, MDA-5 and the NALP3 inflammasome. *PLoS Pathog* 2009; **5**: e1000480.
- 30 Schmolke M, Patel JR, de Castro E, Sanchez-Aparicio MT, Uccellini MB, Miller JC *et al.* RIG-I detects mRNA of intracellular *Salmonella enterica* serovar Typhimurium during bacterial infection. *MBio* 2014; **5**:e01006–14.
- 31 Jehl SP, Nogueira CV, Zhang X, Starnbach MN. IFN- γ inhibits the cytosolic replication of *Shigella flexneri* via the cytoplasmic RNA sensor RIG-I. *PLoS Pathog* 2012; **8**: e1002809.
- 32 Hagmann CA, Herzner AM, Abdullah Z, Zillinger T, Jakobs C, Schubert C *et al.* RIG-I detects triphosphorylated RNA of *Listeria monocytogenes* during infection in non-immune cells. *PLoS ONE* 2013; **8**:e62872.
- 33 Dhir A, Dhir S, Borowski LS, Jimenez L, Teittel M, Rötig A *et al.* Mitochondrial double-stranded RNA triggers antiviral signalling in humans. *Nature* 2018; **560**:238–42.
- 34 Malathi K, Dong B, Gale M, Silverman RH. Small self-RNA generated by RNase L amplifies antiviral innate immunity. *Nature* 2007; **448**:816–9.
- 35 Sun L, Wu J, Du F, Chen X, Chen ZJ. Cyclic GMP-AMP synthase is a cytosolic DNA sensor that activates the type I interferon pathway. *Science* 2013; **339**:786–91.
- 36 Gao P, Ascano M, Zillinger T, Wang W, Dai P, Serganov AA *et al.* Structure–function analysis of STING activation by c[G(2',5')pA(3',5')p] and targeting by antiviral DMXAA. *Cell* 2013; **154**:748–62.
- 37 Liu S, Cai X, Wu J, Cong Q, Chen X, Li T *et al.* Phosphorylation of innate immune adaptor proteins MAVS, STING, and TRIF induces IRF3 activation. *Science* 2015; **347**: aaa2630.
- 38 Dobbs N, Burnaevskiy N, Chen D, Gonugunta VK, Alto NM, Yan N. STING activation by translocation from the ER is associated with infection and autoinflammatory disease. *Cell Host Microbe* 2015; **18**:157–68.
- 39 Unterholzner L. The interferon response to intracellular DNA: why so many receptors? *Immunobiology* 2013; **218**:1312–21.
- 40 Almine JF, O'Hare CAJ, Dunphy G, Haga IR, Naik RJ, Atrih A *et al.* IFI16 and cGAS cooperate in the activation of STING during DNA sensing in human keratinocytes. *Nat Commun* 2017; **8**:14392.
- 41 Jonsson KL, Laustsen A, Krapp C, Skipper KA, Thavachelvam K, Hotter D *et al.* IFI16 is required for DNA sensing in human macrophages by promoting production and function of cGAMP. *Nat Commun* 2017; **8**:14391.
- 42 Orzalli MH, DeLuca NA, Knipe DM. Nuclear IFI16 induction of IRF-3 signaling during herpesviral infection and degradation of IFI16 by the viral ICP0 protein. *Proc Natl Acad Sci USA* 2012; **109**:E3008–17.
- 43 Morrone SR, Wang T, Constantoulakis LM, Hooy RM, Delannoy MJ, Sohn J. Cooperative assembly of IFI16 filaments on dsDNA provides insights into host defense strategy. *Proc Natl Acad Sci USA* 2014; **111**:E62–71.
- 44 Stratmann SA, Morrone SR, van Oijen AM, Sohn J. The innate immune sensor IFI16 recognizes foreign DNA in the nucleus by scanning along the duplex. *Elife* 2015; **4**: e11721.
- 45 Hornung V, Ablasser A, Charrel-Dennis M, Bauernfeind F, Horvath G, Caffrey DR *et al.* AIM2 recognizes cytosolic dsDNA and forms a caspase-1-activating inflammasome with ASC. *Nature* 2009; **458**:514–8.
- 46 Bürckstümmer T, Baumann C, Blüml S, Dixit E, Dürnberger G, Jahn H *et al.* An orthogonal proteomic-genomic screen identifies AIM2 as a cytoplasmic DNA sensor for the inflammasome. *Nat Immunol* 2009; **10**:266–72.
- 47 Fernandes-Alnemri T, Yu J-W, Datta P, Wu J, Alnemri ES. AIM2 activates the inflammasome and cell death in response to cytoplasmic DNA. *Nature* 2009; **458**:509–13.
- 48 Takaoka A, Wang Z, Choi MK, Yanai H, Negishi H, Ban T *et al.* DAI (DLM-1/ZBP1) is a cytosolic DNA sensor and an activator of innate immune response. *Nature* 2007; **448**:501–5.
- 49 Ferguson BJ, Mansur DS, Peters NE, Ren H, Smith GL. DNA-PK is a DNA sensor for IRF-3-dependent innate immunity. *Elife* 2012; **1**:e00047.
- 50 Kim T, Pazhoor S, Bao M, Zhang Z, Hanabuchi S, Facchinetti V *et al.* Aspartate-glutamate-alanine-histidine box motif (DEAH)/RNA helicase A helicases sense microbial DNA in human plasmacytoid dendritic cells. *Proc Natl Acad Sci USA* 2010; **107**:15181–6.
- 51 Zhang Z, Yuan B, Bao M, Lu N, Kim T, Liu Y-J. The helicase DDX41 senses intracellular DNA mediated by the adaptor STING in dendritic cells. *Nat Immunol* 2011; **12**:959–65.
- 52 Kondo T, Kobayashi J, Saitoh T, Maruyama K, Ishii KJ, Barber GN *et al.* DNA damage sensor MRE11 recognizes cytosolic double-stranded DNA and induces type I interferon by regulating STING trafficking. *Proc Natl Acad Sci USA* 2013; **110**:2969–74.
- 53 Gao D, Wu J, Wu Y-T, Du F, Aroh C, Yan N *et al.* Cyclic GMP-AMP synthase is an innate immune sensor of HIV and other retroviruses – supplementary materials. *Science* 2013; **341**:903–6.
- 54 Li X-D, Wu J, Gao D, Wang H, Sun L, Chen ZJ. Pivotal roles of cGAS-cGAMP signaling in antiviral defense and immune adjuvant effects. *Science* 2013; **341**:1390–4.
- 55 Aguirre S, Luthra P, Sanchez-Aparicio MT, Maestre AM, Patel J, Lamothe F *et al.* Dengue virus NS2B protein targets cGAS for degradation and prevents mitochondrial DNA sensing during infection. *Nat Microbiol* 2017; **2**:17037.
- 56 Waßermann R, Gulen MF, Sala C, Perin SG, Lou Y, Rybniker J *et al.* *Mycobacterium tuberculosis* differentially activates cGAS- and inflammasome-dependent intracellular immune responses through ESX-1. *Cell Host Microbe* 2015; **17**:799–810.
- 57 Collins AC, Cai H, Li T, Franco LH, Li X-D, Nair VR *et al.* Cyclic GMP-AMP synthase is an innate immune DNA sensor for *Mycobacterium tuberculosis*. *Cell Host Microbe* 2015; **17**:820–8.
- 58 Sisquella X, Ofir-Birin Y, Pimentel MA, Cheng L, Abou Karam P, Sampaio NG *et al.* Malaria parasite DNA-harboring vesicles activate cytosolic immune sensors. *Nat Commun* 2017; **8**:1985.
- 59 Hansen K, Prabhakaran T, Laustsen A, Jorgensen SE, Rahbaek SH, Jensen SB *et al.* *Listeria monocytogenes* induces IFN expression through an IFI16-, cGAS- and STING-dependent pathway. *EMBO J* 2014; **33**:1654–66.
- 60 Burdette DL, Monroe KM, Sotelo-Troha K, Iwig JS, Eckert B, Hyodo M *et al.* STING is a direct innate immune sensor of cyclic di-GMP. *Nature* 2011; **478**:515–8.
- 61 Dey B, Dey RJ, Cheung LS, Pokkali S, Guo H, Lee J-H *et al.* A bacterial cyclic dinucleotide activates the cytosolic surveillance pathway and mediates innate resistance to tuberculosis. *Nat Med* 2015; **21**:401–6.
- 62 Marinho FV, Benmerzoug S, Oliveira SC, Ryffel B, Quesniaux VFJ. The emerging roles of STING in bacterial infections. *Trends Microbiol* 2017; **25**:906–18.
- 63 Archer KA, Durack J, Portnoy DA. STING-dependent type I IFN production inhibits cell-mediated immunity to *Listeria monocytogenes*. *PLoS Pathog* 2014; **10**:e1003861.
- 64 Snyder DT, Hedges JF, Jutila MA. Getting “inside” Type I IFNs: Type I IFNs in intracellular bacterial infections. *J Immunol Res* 2017; **2017**:9361802.
- 65 Hancks DC, Hartley MK, Hagan C, Clark NL, Elde NC. Overlapping patterns of rapid evolution in the nucleic acid sensors cGAS and OAS1 suggest a common mechanism of pathogen antagonism and escape. *PLoS Genet* 2015; **11**:e1005203.
- 66 Lemos de Matos A, McFadden G, Esteves PJ. Positive evolutionary selection on the RIG-I-like receptor genes in mammals. *PLoS ONE* 2013; **8**:e81864.
- 67 Sorci G, Cornet S, Faivre B. Immunity and the emergence of virulent pathogens. *Infect Genet Evol* 2013; **16**:441–6.
- 68 Gröschel MI, Sayes F, Simeone R, Majlessi L, Brosch R. ESX secretion systems: mycobacterial evolution to counter host immunity. *Nat Rev Microbiol* 2016; **14**:677–91.
- 69 Uchida L, Espada-Murao LA, Takamatsu Y, Okamoto K, Hayasaka D, Yu F *et al.* The dengue virus conceals double-stranded RNA in the intracellular membrane to escape from an interferon response. *Sci Rep* 2014; **4**:7395.
- 70 Neufeldt CJ, Joyce MA, Van Buuren N, Levin A, Kirkegaard K, Gale M *et al.* The Hepatitis C Virus-induced membranous web and associated nuclear transport machinery limit access of pattern recognition receptors to viral replication sites. *PLoS Pathog* 2016; **12**:e1005428.
- 71 Jacques DA, McEwan WA, Hilditch L, Price AJ, Towers GJ, James LC. HIV-1 uses dynamic capsid pores to import nucleotides and fuel encapsidated DNA synthesis. *Nature* 2016; **536**:349–53.
- 72 Kindler E, Gil-Cruz C, Spanier J, Li Y, Wilhelm J, Rabouw HH *et al.* Early endonuclease-mediated evasion of RNA sensing ensures efficient coronavirus replication. *PLoS Pathog* 2017; **13**:e1006195.
- 73 Andrade WA, Firon A, Schmidt T, Hornung V, Fitzgerald KA, Kurt-Jones EA *et al.* Group B streptococcus degrades cyclic-di-AMP to modulate STING-dependent Type I interferon production. *Cell Host Microbe* 2016; **20**:49–59.
- 74 Habjan M, Andersson I, Klingström J, Schümann M, Martin A, Zimmermann P *et al.* Processing of genome 5' termini as a strategy of negative-strand RNA viruses to avoid RIG-I-dependent interferon induction. *PLoS ONE* 2008; **3**:e2032.

- 75 Decroly E, Canard B. Biochemical principles and inhibitors to interfere with viral capping pathways. *Curr Opin Virol* 2017; **24**:87–96.
- 76 Schuberth-Wagner C, Ludwig J, Bruder AK, Herzner AM, Zillinger T, Goldeck M *et al.* A conserved Histidine in the RNA sensor RIG-I controls immune tolerance to N1-2'O-methylated Self RNA. *Immunity* 2015; **43**:41–51.
- 77 Abdullah Z, Schlee M, Roth S, Mraheil MA, Barchet W, Böttcher J *et al.* RIG-I detects infection with live *Listeria* by sensing secreted bacterial nucleic acids. *EMBO J* 2012; **31**:4153–64.
- 78 Luecke S, Holleufer A, Christensen MH, Jönsson KL, Boni GA, Sørensen LK *et al.* cGAS is activated by DNA in a length-dependent manner. *EMBO Rep* 2017; **18**:1707–15.
- 79 Smith GL, Benfield CTO, Maluquer de Motes C, Mazzon M, Ember SWJ, Ferguson BJ *et al.* Vaccinia virus immune evasion: mechanisms, virulence and immunogenicity. *J Gen Virol* 2013; **94**:2367–92.
- 80 Su C, Zhan G, Zheng C. Evasion of host antiviral innate immunity by HSV-1, an update. *Virol J BioMed Central* 2016; **13**:38.
- 81 Feng Q, Langereis MA, Lork M, Nguyen M, Hato SV, Lanke K *et al.* Enterovirus 2Apro targets MDA5 and MAVS in infected cells. *J Virol* 2014; **88**:3369–78.
- 82 Barral PM, Sarkar D, Fisher PB, Racaniello VR. RIG-I is cleaved during picornavirus infection. *Virology* 2009; **391**:171–6.
- 83 Papon L, Oteiza A, Imaizumi T, Kato H, Brocchi E, Lawson TG *et al.* The viral RNA recognition sensor RIG-I is degraded during encephalomyocarditis virus (EMCV) infection. *Virology* 2009; **393**:311–8.
- 84 Rodríguez Pulido M, Sánchez-Aparicio MT, Martínez-Salas E, García-Sastre A, Sobrino F, Sáiz M. Innate immune sensor LGP2 is cleaved by the Leader protease of foot-and-mouth disease virus. *PLoS Pathog* 2018; **14**:e1007135.
- 85 Gori-Savellini G, Valentini M, Cusi MG. Toscana virus NSs protein inhibits the induction of type I interferon by interacting with RIG-I. *J Virol* 2013; **87**:6660–7.
- 86 Qin L, Ren L, Zhou Z, Lei X, Chen L, Xue Q *et al.* Rotavirus nonstructural protein 1 antagonizes innate immune response by interacting with retinoic acid inducible gene I. *Virol J BioMed Central* 2011; **8**:526.
- 87 Zhang H-L, Ye H-Q, Liu S-Q, Deng C-L, Li X-D, Shi P-Y *et al.* West Nile virus NS1 antagonizes interferon β production by targeting RIG-I and MDA5. *J Virol* 2017; **91**:e02396-16.
- 88 Lu Y, Qin Z, Wang J, Zheng X, Lu J, Zhang X *et al.* Epstein-Barr virus miR-BART6-3p inhibits the RIG-I pathway. *J Innate Immun* 2017; **9**:574–86.
- 89 Hou Z, Zhang J, Han Q, Su C, Qu J, Xu D *et al.* Hepatitis B virus inhibits intrinsic RIG-I and RIG-G immune signaling via inducing miR146a. *Sci Rep* 2016; **6**:26150.
- 90 Zhao J, Zeng Y, Xu S, Chen J, Shen G, Yu C *et al.* A viral deamidase targets the helicase domain of RIG-I to block RNA-induced activation. *Cell Host Microbe* 2016; **20**:770–84.
- 91 Kok K-H, Lui P-Y, Ng M-HJ, Siu K-L, Au SWN, Jin D-Y. The double-stranded RNA-binding protein PACT functions as a cellular activator of RIG-I to facilitate innate antiviral response. *Cell Host Microbe* 2011; **9**:299–309.
- 92 Kew C, Lui P-Y, Chan C-P, Liu X, Au SWN, Mohr I *et al.* Suppression of PACT-induced type I interferon production by herpes simplex virus 1 Us11 protein. *J Virol* 2013; **87**:13141–9.
- 93 Siu K-L, Yeung ML, Kok K-H, Yuen K-S, Kew C, Lui P-Y *et al.* Middle east respiratory syndrome coronavirus 4a protein is a double-stranded RNA-binding protein that suppresses PACT-induced activation of RIG-I and MDA5 in the innate antiviral response. *J Virol* 2014; **88**:4866–76.
- 94 Luthra P, Ramanan P, Mire CE, Weisend C, Tsuda Y, Yen B *et al.* Mutual antagonism between the Ebola virus VP35 protein and the RIG-I activator PACT determines infection outcome. *Cell Host Microbe* 2013; **14**:74–84.
- 95 Tawaratsumida K, Phan V, Hrinčius ER, High AA, Webby R, Redecke V *et al.* Quantitative proteomic analysis of the influenza A virus nonstructural proteins NS1 and NS2 during natural cell infection identifies PACT as an NS1 target protein and antiviral host factor. *J Virol* 2014; **88**:9038–48.
- 96 Wies E, Wang MK, Maharaj NP, Chen K, Zhou S, Finberg RW *et al.* Dephosphorylation of the RNA sensors RIG-I and MDA5 by the phosphatase PP1 is essential for innate immune signaling. *Immunity* 2013; **38**:437–49.
- 97 Gack MU, Shin YC, Joo C-H, Urano T, Liang C, Sun L *et al.* TRIM25 RING-finger E3 ubiquitin ligase is essential for RIG-I-mediated antiviral activity. *Nature* 2007; **446**:916–20.
- 98 Oshiumi H, Miyashita M, Inoue N, Okabe M, Matsumoto M, Seya T. The ubiquitin ligase Riplet is essential for RIG-I-dependent innate immune responses to RNA virus infection. *Cell Host Microbe* 2010; **8**:496–509.
- 99 Mesman AW, Zijlstra-Willems EM, Kaptein TM, de Swart RL, Davis ME, Ludlow M *et al.* Measles virus suppresses RIG-I-like receptor activation in dendritic cells via DC-SIGN-mediated inhibition of PP1 phosphatases. *Cell Host Microbe* 2014; **16**:31–42.
- 100 Rajsbbaum R, Albrecht RA, Wang MK, Maharaj NP, Versteeg GA, Nistal-Villán E *et al.* Species-specific inhibition of RIG-I ubiquitination and IFN induction by the influenza A virus NS1 protein. *PLoS Pathog* 2012; **8**:e1003059.
- 101 Oshiumi H, Miyashita M, Matsumoto M, Seya T. A distinct role of Riplet-mediated K63-Linked polyubiquitination of the RIG-I repressor domain in human antiviral innate immune responses. *PLoS Pathog* 2013; **9**:e1003533.
- 102 Gupta S, Ylä-Anttila P, Callegari S, Tsai M-H, Delecluse H-J, Masucci MG. Herpesvirus deconjugases inhibit the IFN response by promoting TRIM25 autoubiquitination and functional inactivation of the RIG-I signalosome. *PLoS Pathog* 2018; **14**:e1006852.
- 103 Gack MU, Albrecht RA, Urano T, Inn K-S, Huang I-C, Carnero E *et al.* Influenza A virus NS1 targets the ubiquitin ligase TRIM25 to evade recognition by the host viral RNA sensor RIG-I. *Cell Host Microbe* 2009; **5**:439–49.
- 104 Koliopoulos MG, Lethier M, Van der Veen AG, Haubrich K, Hennig J, Kowalinski E *et al.* Molecular mechanism of influenza A NS1-mediated TRIM25 recognition and inhibition. *Nat Commun* 2018; **9**:1820.
- 105 Sanchez-Aparicio MT, Feinman LJ, García-Sastre A, Shaw ML. Paramyxovirus V proteins interact with the RIG-I/TRIM25 regulatory complex and inhibit RIG-I signaling. *J Virol* 2018; **92**:396.
- 106 Chiang C, Pauli E-K, Biryukov J, Feister KF, Meng M, White EA *et al.* The human Papillomavirus E6 Oncoprotein targets USP15 and TRIM25 to suppress RIG-I-mediated innate immune signaling. *J Virol* 2018; **92**:1.
- 107 Hu Y, Li W, Gao T, Cui Y, Jin Y, Li P *et al.* The severe acute respiratory syndrome coronavirus nucleocapsid inhibits Type I interferon production by interfering with TRIM25-mediated RIG-I ubiquitination. *J Virol* 2017; **91**:e02143-16.
- 108 Manokaran G, Finol E, Wang C, Gunaratne J, Bahl J, Ong EZ *et al.* Dengue subgenomic RNA binds TRIM25 to inhibit interferon expression for epidemiological fitness. *Science* 2015; **350**:217–21.
- 109 Lee W, Lee S-H, Kim M, Moon J-S, Kim G-W, Jung H-G *et al.* *Vibrio vulnificus* quorum-sensing molecule cyclo(Phe-Pro) inhibits RIG-I-mediated antiviral innate immunity. *Nat Commun* 2018; **9**:1606.
- 110 Chan YK, Gack MU. A phosphomimetic-based mechanism of dengue virus to antagonize innate immunity. *Nat Immunol* 2016; **17**:523–30.
- 111 Xing J, Wang S, Lin R, Mossman KL, Zheng C. Herpes simplex virus 1 tegument protein US11 downmodulates the RLR signaling pathway via direct interaction with RIG-I and MDA-5. *J Virol* 2012; **86**:3528–40.
- 112 Zhou S, Cerny AM, Zacharia A, Fitzgerald KA, Kurt-Jones EA, Finberg RW. Induction and inhibition of type I interferon responses by distinct components of lymphocytic choriomeningitis virus. *J Virol* 2010; **84**:9452–62.
- 113 Ling Z, Tran KC, Teng MN. Human respiratory syncytial virus nonstructural protein NS2 antagonizes the activation of β interferon transcription by interacting with RIG-I. *J Virol* 2009; **83**:3734–42.
- 114 Fan L, Briese T, Lipkin WI. Z proteins of New World arenaviruses bind RIG-I and interfere with type I interferon induction. *J Virol* 2010; **84**:1785–91.
- 115 Ren J, Wang Q, Kolli D, Prusak DJ, Tseng C-TK, Chen ZJ *et al.* Human metapneumovirus M2-2 protein inhibits innate cellular signaling by targeting MAVS. *J Virol* 2012; **86**:13049–61.
- 116 Ma J, Ketkar H, Geng T, Lo E, Wang L, Xi J *et al.* Zika virus non-structural protein 4A blocks the RLR-MAVS signaling. *Front Microbiol* 2018; **9**:1350.
- 117 He Z, Zhu X, Wen W, Yuan J, Hu Y, Chen J *et al.* Dengue virus subverts host innate immunity by targeting adaptor protein MAVS. *J Virol* 2016; **90**:7219–30.
- 118 Hwang KY, Choi YB. Modulation of mitochondrial antiviral signaling by human herpesvirus 8 interferon regulatory factor 1. *J Virol* 2016; **90**:506–20.
- 119 Shi C-S, Qi H-Y, Boularan C, Huang N-N, Abu-Asab M, Shelhamer JH *et al.* SARS-coronavirus open reading frame-9b suppresses innate immunity by targeting mitochondria and the MAVS/TRAF3/TRAF6 signalosome. *J Immunol* 2014; **193**:3080–9.
- 120 Wei C, Ni C, Song T, Liu Y, Yang X, Zheng Z *et al.* The hepatitis B virus X protein disrupts innate immunity by downregulating mitochondrial antiviral signaling protein. *J Immunol* 2010; **185**:1158–68.
- 121 Meylan E, Curran J, Hofmann K, Moradpour D, Binder M, Bartschslager R *et al.* Cardif is an adaptor protein in the RIG-I antiviral pathway and is targeted by hepatitis C virus. *Nature* 2005; **437**:1167–72.
- 122 Bender S, Reuter A, Eberle F, Einhorn E, Binder M, Bartschslager R. Activation of Type I and III interferon response by mitochondrial and peroxisomal MAVS and inhibition by hepatitis C virus. *PLoS Pathog* 2015; **11**:e1005264.
- 123 Horner SM, Park HS, Gale M. Control of innate immune signaling and membrane targeting by the Hepatitis C virus NS3/4A protease are governed by the NS3 helix $\alpha 0$. *J Virol* 2012; **86**:3112–20.
- 124 Lin R, Lacoste J, Nakhaei P, Sun Q, Yang L, Paz S *et al.* Dissociation of a MAVS/IPS-1/VISA/Cardif-IKKEpsilon molecular complex from the mitochondrial outer membrane by hepatitis C virus NS3-4A proteolytic cleavage. *J Virol* 2006; **80**:6072–83.
- 125 Choi HJ, Park A, Kang S, Lee E, Lee TA, Ra EA *et al.* Human cytomegalovirus-encoded US9 targets MAVS and STING signaling to evade type I interferon immune responses. *Nat Commun* 2018; **9**:125.
- 126 Varga ZT, Grant A, Manicassamy B, Palese P. Influenza virus protein PB1-F2 inhibits the induction of type I interferon by binding to MAVS and decreasing mitochondrial membrane potential. *J Virol* 2012; **86**:8359–66.

- 127 Yoshizumi T, Ichinohe T, Sasaki O, Otera H, Kawabata S-I, Mihara K *et al.* Influenza A virus protein PB1-F2 translocates into mitochondria via Tom40 channels and impairs innate immunity. *Nat Commun* 2014; **5**:4713.
- 128 Chatel-Chaix L, Cortese M, Romero-Brey I, Bender S, Neufeldt CJ, Fischl W *et al.* Dengue virus perturbs mitochondrial morphodynamics to dampen innate immune responses. *Cell Host Microbe* 2016; **20**:342–56.
- 129 Magalhães AC, Ferreira AR, Gomes S, Vieira M, Gouveia A, Valença I *et al.* Peroxisomes are platforms for cytomegalovirus' evasion from the cellular immune response. *Sci Rep* 2016; **6**:26028.
- 130 Zheng C, Su C. Herpes simplex virus 1 infection dampens the immediate early antiviral innate immunity signaling from peroxisomes by tegument protein VP16. *Virology* 2017; **14**:35.
- 131 Kamanova J, Sun H, Lara-Tejero M, Galán JE. The *Salmonella* effector protein SopA modulates innate immune responses by targeting TRIM E3 ligase family members. *PLoS Pathog* 2016; **12**:e1005552.
- 132 Fiskin E, Bhogaraju S, Herhaus L, Kalayil S, Hahn M, Dikic I. Structural basis for the recognition and degradation of host TRIM proteins by *Salmonella* effector SopA. *Nat Commun* 2017; **8**:14004.
- 133 Christensen MH, Jensen SB, Miettinen JJ, Luecke S, Prabakaran T, Reinert LS *et al.* HSV-1 ICP27 targets the TBK1-activated STING signaling to inhibit virus-induced type I IFN expression. *EMBO J* 2016; **35**:1385–99.
- 134 Orzalli MH, Broekema NM, Knipe DM. Relative contributions of Herpes simplex virus 1 ICP0 and VHS to loss of cellular IFI16 vary in different human cell types. *J Virol* 2016; **90**:8351–9.
- 135 Su C, Zheng C. Herpes simplex virus 1 abrogates the cGAS/STING-mediated cytosolic DNA-sensing pathway via its virion host shutoff protein, UL41. *J Virol* 2017; **91**:e02414–16.
- 136 Huang J, You H, Su C, Li Y, Chen S, Zheng C. Herpes simplex virus 1 tegument protein VP22 abrogates cGAS/STING-mediated antiviral innate immunity. *J Virol* 2018; **92**:783.
- 137 Maruzuru Y, Ichinohe T, Sato R, Miyake K, Okano T, Suzuki T *et al.* Herpes simplex virus 1 VP22 inhibits AIM2-dependent inflammasome activation to enable efficient viral replication. *Cell Host Microbe* 2018; **23**:254–7.
- 138 Zhang J, Zhao J, Xu S, Li J, He S, Zeng Y *et al.* Species-specific deamidation of cGAS by Herpes simplex virus UL37 protein facilitates viral replication. *Cell Host Microbe* 2018; **24**:234–5.
- 139 Diner BA, Lum KK, Toettcher JE, Cristea IM. Viral DNA sensors IFI16 and cyclic GMP-amp synthase possess distinct functions in regulating viral gene expression, immune defenses, and apoptotic responses during Herpesvirus infection. *MBio* 2016; **7**:e01553-16.
- 140 Huang Z-F, Zou H-M, Liao B-W, Zhang H-Y, Yang Y, Fu Y-Z *et al.* Human cytomegalovirus protein UL31 inhibits DNA sensing of cGAS to mediate immune evasion. *Cell Host Microbe* 2018; **24**:69–80.e4.
- 141 Biolatti M, Dell'Oste V, Pautasso S, Gugliesi F, von Einem J, Krapp C *et al.* Human cytomegalovirus tegument protein pp65 (pUL83) Dampens Type I Interferon production by inactivating the DNA sensor cGAS without affecting STING. *J Virol* 2018; **92**:288.
- 142 Biolatti M, Dell'Oste V, Pautasso S, von Einem J, Marschall M, Plachter B *et al.* Regulatory interaction between the cellular restriction factor IFI16 and viral pp65 (pUL83) modulates viral gene expression and IFI16 protein stability. *J Virol* 2016; **90**:8238–50.
- 143 Li T, Chen J, Cristea IM. Human cytomegalovirus tegument protein pUL83 inhibits IFI16-mediated DNA sensing for immune evasion. *Cell Host Microbe* 2013; **14**:591–9.
- 144 Fu Y-Z, Su S, Gao Y-Q, Wang P-P, Huang Z-F, Hu M-M *et al.* Human cytomegalovirus tegument protein UL82 inhibits STING-mediated signaling to evade antiviral immunity. *Cell Host Microbe* 2017; **21**:231–43.
- 145 Kim J-E, Kim Y-E, Stinski MF, Ahn J-H, Song Y-J. Human cytomegalovirus IE2 86 kDa protein induces STING degradation and inhibits cGAMP-mediated IFN- β induction. *Front Microbiol* 2017; **8**:1854.
- 146 Huang Y, Ma D, Huang H, Lu Y, Liao Y, Liu L *et al.* Interaction between HCMV pUL83 and human AIM2 disrupts the activation of the AIM2 inflammasome. *Virology* 2017; **14**:34.
- 147 Ma Z, Jacobs SR, West JA, Stopford C, Zhang Z, Davis Z *et al.* Modulation of the cGAS-STING DNA sensing pathway by gammaherpesviruses. *Proc Natl Acad Sci USA* 2015; **112**:E4306–15.
- 148 Wu J-J, Li W, Shao Y, Avey D, Fu B, Gillen J *et al.* Inhibition of cGAS DNA Sensing by a Herpesvirus Virion Protein. *Cell Host Microbe* 2015; **18**:333–44.
- 149 Roy A, Dutta D, Iqbal J, Pisano G, Gjyshi O, Ansari MA *et al.* Nuclear innate immune DNA sensor IFI16 is degraded during lytic reactivation of Kaposi's Sarcoma-associated Herpesvirus (KSHV): role of IFI16 in maintenance of KSHV latency. *J Virol* 2016; **90**:8822–41.
- 150 Zhang G, Chan B, Samarina N, Abere B, Weidner-Glunde M, Buch A *et al.* Cytoplasmic isoforms of Kaposi sarcoma herpesvirus LANA recruit and antagonize the innate immune DNA sensor cGAS. *Proc Natl Acad Sci USA* 2016; **113**:E1034–43.
- 151 Lau L, Gray EE, Brunette RL, Stetson DB. DNA tumor virus oncogenes antagonize the cGAS-STING DNA-sensing pathway. *Science* 2015; **350**:568–71.
- 152 Georgana I, Sumner RP, Towers GJ, Maluquer de Motes C. Virulent poxviruses inhibit DNA sensing by preventing STING activation. *J Virol* 2018; **92**:e02145–17.
- 153 Peters NE, Ferguson BJ, Mazzon M, Fahy AS, Kryzstofinska E, Arribas-Bosacoma R *et al.* A mechanism for the inhibition of DNA-PK-mediated DNA sensing by a virus. *PLoS Pathog* 2013; **9**:e1003649.
- 154 Upton JW, Kaiser WJ, Mocarski ES. DAI/ZBP1/DLM-1 complexes with RIP3 to mediate virus-induced programmed necrosis that is targeted by murine cytomegalovirus vIRA. *Cell Host Microbe* 2012; **11**:290–7.
- 155 Holm CK, Jensen SB, Jakobsen MR, Cheshenko N, Horan KA, Moeller HB *et al.* Virus-cell fusion as a trigger of innate immunity dependent on the adaptor STING. *Nat Immunol* 2012; **13**:737–43.
- 156 Holm CK, Rahbek SH, Gad HH, Bak RO, Jakobsen MR, Jiang Z *et al.* Influenza A virus targets a cGAS-independent STING pathway that controls enveloped RNA viruses. *Nat Commun* 2016; **7**:10680.
- 157 Ding Q, Gaska JM, Douam F, Wei L, Kim D, Balev M *et al.* Species-specific disruption of STING-dependent antiviral cellular defenses by the Zika virus NS2B3 protease. *Proc Natl Acad Sci USA* 2018; **115**:E6310–8.
- 158 Aguirre S, Maestre AM, Pagni S, Patel JR, Savage T, Gutman D *et al.* DENV inhibits type I IFN production in infected cells by cleaving human STING. *PLoS Pathog* 2012; **8**:e1002934.
- 159 Sun B, Sundström KB, Chew JJ, Bist P, Gan ES, Tan HC *et al.* Dengue virus activates cGAS through the release of mitochondrial DNA. *Sci Rep* 2017; **7**:3594.
- 160 Cao Y, Guan K, He X, Wei C, Zheng Z, Zhang Y *et al.* Yersinia YopJ negatively regulates IRF3-mediated antibacterial response through disruption of STING-mediated cytosolic DNA signaling. *Biochim Biophys Acta* 2016; **1863**:3148–59.
- 161 Dey RJ, Dey B, Zheng Y, Cheung LS, Zhou J, Sayre D *et al.* Inhibition of innate immune cytosolic surveillance by an *M. tuberculosis* phosphodiesterase. *Nat Chem Biol* 2017; **13**:210–7.
- 162 Moverit E, Lienard J, Valfridsson C, Nordström T, Johansson-Lindbom B, Carlsson F. Streptococcal M protein promotes IL-10 production by cGAS-independent activation of the STING signaling pathway. *PLoS Pathog* 2018; **14**:e1006969.