

1 **A pilot study using wearable global position system (GPS) dataloggers to compare water**
2 **contact levels: *Schistosoma haematobium* infection in pre-school-aged children (PSAC)**
3 **and their mothers at Barombi Kotto, Cameroon**

4

5 Grace Macklin^{a,+}, Michelle C. Stanton^b, Louis Albert Tchuem-Tchuente^c, J. Russell Stothard^{a,*}

6

7 ^a Department of Parasitology, Liverpool School of Tropical Medicine, Liverpool, UK; ^b Faculty
8 of Health and Medicine, Lancaster University, Lancaster, UK; ^c Centre for Schistosomiasis &
9 Parasitology, Yaoundé, Cameroon

10

11 ⁺ present address: London School of Hygiene and Tropical Medicine, Keppel Street, London
12 WC1E 7HT, UK;

13 ^{*} Corresponding author: Tel: +44 151 7053724; E-mail: russell.stothard@lstmed.ac.uk

14

15 **Abstract**

16 Barombi Kotto, Cameroon serves as a reference location for assessing intervention strategies
17 against *Schistosoma haematobium*. As part of a pilot study, the whole community was treated
18 with praziquantel, inclusive of pre-school-aged children (PSAC) and their mothers. One year
19 later egg-patent infections were reassessed and water contact patterns of 12 pairs of PSAC
20 and their mother were measured with global position system (GPS) dataloggers. A substantial
21 reduction in general infection prevalence, from 44.8% to 11.7 %, was observed but certain
22 PSAC and mothers continued to have egg-patent infections. Analysis of GPS data
23 demonstrated similar water contact levels between child and mother groups, although
24 certain individuals were numerical outliers. This study shows the potential of GPS dataloggers
25 to clarify the at-risk status of PSAC and mothers.

26

27 **Keywords**

28 Urogenital schistosomiasis, i-gotU, paediatric schistosomiasis, female genital
29 schistosomiasis, praziquantel

30 **Introduction**

31 Urogenital schistosomiasis is an important waterborne disease, caused by infection with the
32 blood fluke *Schistosoma haematobium*, and common in many parts of sub-Saharan Africa (1).
33 In Cameroon, for example, there is a national control programme active in the distribution of
34 praziquantel (PZQ) to school-aged children (SAC) (2, 3). However, in the move towards local
35 interruption of schistosome transmission, the programme is developing new tactics of control
36 (4) and has benefited from recent bilateral support from China in snail control and
37 environmental surveillance (5), as well as from UK to expand access of interventions (6).

38 Overlooked for too long, expanding access of PZQ to pre-school-aged children (PSAC)
39 and their mothers is attracting increasing attention (7, 8). It has been shown elsewhere that
40 these groups can be patently infected (9-12) and alongside SAC, may contribute towards
41 schistosome transmission but their water contact(s) is rarely measured and hence the role of
42 PSAC in sustaining local transmission remains speculative (13, 14). As a pilot investigation of
43 expanded access to praziquantel treatment, in June 2016 Campbell *et al.* undertook a detailed
44 cross-sectional epidemiological and malacological survey at Barombi Kotto, Cameroon (15).
45 Barombi Kotto is well-known crater lake and is of significant international interest as a
46 longstanding focus of urogenital schistosomiasis (16-18). Before treating all community
47 members with PZQ, Campbell *et al.* noted that a quarter of PSAC had egg-patent infections.
48 Furthermore, adult women had raised signs and symptoms of female genital schistosomiasis
49 (FGS), the latter is of growing international concern (13, 19). Environmental water contact is
50 very common across the community, for example, bathing, washing and other domestic
51 chores are typically performed on the immediate shoreline of the island while potable water
52 is collected in plastic containers from a local stream which is only accessible by canoe (15).
53 The level of environmental water contact, however, on the immediate lake shoreline of both
54 PSAC and their mothers remained to be determined and compared.

55 To shed fresh light on the at-risk status of PSAC and their mothers, using wearable
56 global position system (GPS) dataloggers, we attempted to measure and compare the water
57 contact patterns of PSAC and their mothers (20, 21). Furthermore, we hoped to pinpoint
58 water contact sites, measuring putative immersion times, on the Barombi Kotto crater lake
59 shoreline as baseline information for future interventions.

60

61

62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93

Methods & Materials

Study location and parasitological examination

This parasitological resurvey and GPS study was conducted in June 2017 in the community on the central island of Barombi Kotto crater lake, where some 375 people are permanent residents. Study protocols were approved by the Liverpool School of Tropical Medicine Research Ethics Committee and the Cameroon National Ethical Committee of Research for Human Health. Participation involved obtaining written informed consent from mothers and their PSAC before deployment of the GPS datalogger. A total of 179 individuals (20 PSAC, 55 SAC and 104 adults) underwent a parasitological reinspection where each provided a 10ml urine sample which was filtered and stained with Lugol's iodine to visualise *S. haematobium* eggs by microscopy as described previously (15). All participants found infected with schistosomiasis were offered and observed to take praziquantel treatment (40 mg/kg)

Water exposure assessment

A subset of 12 mothers and PSAC pairs were randomly selected, then 6 pairs were assigned into two groups to wear the GPS dataloggers (i-gotU-120, Mobile Action, UK; dimension 44.5 x 28.5 x 13 mm, weight 20 g) over a 48-hour period on two occasions. The dataloggers were worn on the arm or wrist using a custom made elastic strap (20). The dataloggers were configured to record GPS location and velocity at 1 minute intervals during waking hours (05:00-21:00). Analysis of GPS data was conducted in QGIS (22) and filtered per the velocity filtering method (23). A zone was developed around the lakeshore of the island, 10m into the lake and 5m into the shore, a conservative assessment of the positional accuracy of the i-gotU-120 dataloggers based on previous observations (24). A water contact event was defined as a GPS location recording within a defined lakeshore geospatial buffer zone around the island circumference: a conservative assessment of the positional accuracy of the i-gotU dataloggers (24). As the GPS loggers recorded location at 1 minute intervals, each water contact event is analogous to 1 minute spent in the geospatial buffer zone and could be tallied and compared between individuals and groups.

94 **Statistical analysis**

95 Statistical analysis was performed using the R statistical software (25). Prevalence of
96 schistosomiasis was calculated with 95% binomial confidence intervals (95% CIs) with
97 correction for samples of $n < 30$. The track logs of each GPS unit were plotted and overlaid
98 against a base map of Barombi Kotto shoreline to identify travel patterns on and off the island.

99

100 **Results and Discussion**

101

102 *<please insert table 1 here>*

103

104 The characteristics of the study population and infection status is shown in Table 1. In June
105 2017, the overall prevalence of egg-patent infection was 11.7 % (95% CI 7.0 - 17.0) with only
106 1 infection of heavy intensity encountered albeit in a PSAC. The epidemiological survey
107 undertaken by Campbell *et al.* one year previously observed a much higher egg-patent
108 prevalence of 40.1% (24.6% in PSAC, 51.3% in SAC and 44.9% in adults). The overall reduction
109 across all demographic groups, see Table 1, is most likely due to the community-wide PZQ
110 treatment.

111

112 *<please insert Figure 1 here>*

113

114 Twelve PSAC and mother pairs were randomly selected and assigned into two groups
115 of 6 pairs, to wear the GPS dataloggers on two locations. In total, there were 3 individuals
116 with egg-patent infections (2 mothers and 1 PSAC) and their intensities of egg-patent
117 infections against the sampled population is shown in Figure 1A. The water contact levels of
118 the 12 mother and child pairs for the 48 hour period is shown in Figure 1B. This documents a
119 high level of water contact events (reaching up to 166 water contact events in 24 hours) in
120 both mothers and PSAC on the Barombi Kotto shoreline. As data points were recorded at 1
121 minute intervals, the number of water contact events can be interpreted as the time spent in
122 water contact which would likely positively correlate with actual bodily immersion or skin
123 contact with lake shore water.

124 The average number of water contact events in 24 hours were 27.4 [95% CIs: -1.3,
125 56.1] for mothers and 14.1 [95% CIs: 8.5 ,19.7] for PSAC, with no significant difference

126 between the two groups, Figure 1B. These findings have important implications in that water
127 contact levels of PSAC should not to be overlooked and follow similar levels to that observed
128 in Uganda (20). In Uganda PSAC were observed to spend on average half an hour on the
129 shoreline of Lake Albert and were clearly shown to be an at-risk vulnerable group not only to
130 first infection but also re-infection (10), as evidenced here in Barombi Kotto. Our study shows
131 the potential of GPS data logging technology to clarify their at-risk status which should assist
132 in better infection surveillance and control of urogenital schistosomiasis in general as well as
133 for regular access to treatment with the soon to be deployed paediatric PZQ formulation (12).

134 Upon more detailed inspection of individual water contact patterns, whilst our GPS
135 sample of two infected mothers (“M1”, “M2”) and one infected PSAC (“X4”) was too small to
136 determine a precise relationship between water exposure and infection status, two of these
137 individuals were clear numerical outliers in terms of their water contact(s) (4.6, 166.1, and
138 31.8 for “M1”, “M2” and “X4”, respectively). The latter two lying far outside the confidence
139 intervals for PSAC and mother averages. This demonstrates the importance of individual
140 variation in exposure and likely environmental contamination, Figure 1B. Indeed, it is very
141 plausible that the water contact behaviours of “M2” and “X4” could classify them as ‘raised-
142 spreaders’ who should be specifically targeted for increased frequency of treatment(s)
143 alongside behavioural change interventions. It remains to be seen if these individuals play
144 more pivotal roles than others in facilitating and sustaining local transmission of
145 schistosomes.

146 Another interesting facet revealed by the GPS dataloggers is the similar geospatial
147 pattern of water contact between the two groups which illustrate that PSAC frequently
148 accompany their mothers to the same locations, Figure C. This is also consistent with other
149 studies using questionnaires (13, 21). Furthermore, observed water contacts were largely co-
150 clustered on the South-West lakeshore of the island, notably an area where activities of the
151 3 infected cases were concentrated. Micro-spatial heterogeneity of schistosome transmission
152 has been described elsewhere (26, 27) and is further evidenced here, Figure 1D. In the future
153 context of interruption of schistosome transmission (14, 27), as a cost-effective measure, it
154 would be sensible to apply focal molluscicides at this location rather than elsewhere, to have
155 highest impact upon removal of infected snail hosts.

156 Our findings document that GPS dataloggers are an accepted method of measuring
157 water exposure in PSAC and their mothers and directly compare environmental risk of

158 schistosomiasis exposure. We suggest that in future the water contact levels of these two
159 demographic groups should further investigated. The wearable GPS technology is also of
160 value to identify putative transmission foci for spatial targeting of interventions.

161 **Authors Contributions**

162 JRS and MS conceived the study; GM, MS, LATT and JRS designed the study protocol; GM
163 carried out the field work; GM and MS carried out the analysis and interpretation of these
164 data. GM and JRS drafted the manuscript with LATT and MS critically revising. All authors read
165 and approved the final manuscript.

166

167 **Funding**

168 GM is in receipt of a Scholarship from the Medical Research Council, United Kingdom. This
169 study was additionally funded as part of the COUNTDOWN research programme consortium,
170 by the Research and Evidence Division of the Department for International Development, UK.
171 The funders had no role in study design, decision to publish, or preparation of the manuscript.

172 **Acknowledgments**

173 We are grateful to Drs James LaCourse and Martyn Stewart, LSTM who helped with local
174 fieldwork in Cameroon, as well as, the local community and field team volunteers at Barombi
175 Kotto.

176

177 **Competing interests**

178 None declared

179

180 **Ethical approval**

181 The study was approved by the Liverpool School of Tropical Medicine and the Cameroon
182 National Ethical Committee of Research for Human Health.

183

184 **References**

- 185 1. Colley DG, Bustinduy AL, Secor WE, et al.; Human schistosomiasis. *Lancet*
186 2014;**383**(9936):2253-64. doi: 10.1016/S0140-6736(13)61949-2.
- 187 2. Tchuente LAT, N'Goran EK; Schistosomiasis and soil-transmitted helminthiasis
188 control in Cameroon and Cote d'Ivoire: implementing control on a limited budget.
189 *Parasitology* 2009;**136**(13):1739-1745. doi: 10.1017/s0031182009005988.
- 190 3. Tchuente LAT, Noumedem CD, Ngassam P, et al.; Mapping of schistosomiasis and
191 soil-transmitted helminthiasis in the regions of Littoral, North-West, South and South-West

192 Cameroon and recommendations for treatment. *Bmc Infectious Diseases* 2013;**13**. doi:
193 10.1186/1471-2334-13-602.

194 4. Tchuente LAT, Rollinson D, Stothard JR, et al.; Moving from control to elimination of
195 schistosomiasis in sub-Saharan Africa: time to change and adapt strategies. *Infectious*
196 *Diseases of Poverty* 2017;**6**. doi: 10.1186/s40249-017-0256-8.

197 5. Xu J, Yu Q, Tchuente LAT, et al.; Enhancing collaboration between China and African
198 countries for schistosomiasis control. *Lancet Infectious Diseases* 2016;**16**(3):376-383. doi:
199 10.1016/s1473-3099(15)00360-6.

200 6. Stothard JR, Kabatereine NB, Archer J, et al.; A centenary of Robert T. Leiper's lasting
201 legacy on schistosomiasis and a COUNTDOWN on control of neglected tropical diseases.
202 *Parasitology* 2017;**144**(12):1602-1612. doi: 10.1017/s0031182016000998.

203 7. Bustinduy AL, Friedman JF, Kjetland EF, et al.; Expanding Praziquantel (PZQ) Access
204 beyond Mass Drug Administration Programs: Paving a Way Forward for a Pediatric PZQ
205 Formulation for Schistosomiasis. *Plos Neglected Tropical Diseases* 2016;**10**(9). doi:
206 10.1371/journal.pntd.0004946.

207 8. Lo NC, Addiss DG, Hotez PJ, et al.; A call to strengthen the global strategy against
208 schistosomiasis and soil-transmitted helminthiasis: the time is now. *Lancet Infectious*
209 *Diseases* 2017;**17**(2):E64-E69. doi: 10.1016/s1473-3099(16)30535-7.

210 9. Bustinduy AL, Stothard JR, Friedman JF; Paediatric and maternal schistosomiasis:
211 shifting the paradigms. *British Medical Bulletin* 2017;**123**(1):115-125. doi:
212 10.1093/bmb/ldx028.

213 10. Mduluzi T, Mutapi F; Putting the treatment of paediatric schistosomiasis into
214 context. *Infectious Diseases of Poverty* 2017;**6**. doi: 10.1186/s40249-017-0300-8.

215 11. Stothard JR, Sousa-Figueiredo JC, Betson M, et al.; Schistosomiasis in African infants
216 and preschool children: let them now be treated! *Trends in Parasitology* 2013;**29**(4):197-
217 205. doi: 10.1016/j.pt.2013.02.001.

218 12. Reinhard-Rupp J, Klohe K; Developing a comprehensive response for treatment of
219 children under 6 years of age with schistosomiasis: research and development of a pediatric
220 formulation of praziquantel. *Infectious Diseases of Poverty* 2017;**6**. doi: 10.1186/s40249-
221 017-0336-9.

222 13. Poole H, Terlouw DJ, Naunje A, et al.; Schistosomiasis in pre-school-age children and
223 their mothers in Chikhwawa district, Malawi with notes on characterization of schistosomes
224 and snails. *Parasites & Vectors* 2014;**7**. doi: 10.1186/1756-3305-7-153.

225 14. Stothard JR, Campbell SJ, Osei-Atweneboana MY, et al.; Towards interruption of
226 schistosomiasis transmission in sub-Saharan Africa: developing an appropriate
227 environmental surveillance framework to guide and to support 'end game' interventions.
228 *Infectious Diseases of Poverty* 2017;**6**. doi: 10.1186/s40249-016-0215-9.

229 15. Campbell SJ, Stothard JR, O'Halloran F, et al.; Urogenital schistosomiasis and
230 soiltransmitted helminthiasis (STH) in Cameroon: An epidemiological update at Barombi
231 Mbo and Barombi Kotto crater lakes assessing prospects for intensified control
232 interventions. *Infectious Diseases of Poverty* 2017;**6**. doi: 10.1186/s40249-017-0264-8.

233 16. Duke BOL, Moore PJ; Use of a molluscicide in conjunction with chemotherapy to
234 control *Schistosoma haematobium* at Barombi Lake Foci in Cameroon. 3. Conclusions and
235 costs. *Tropenmedizin Und Parasitologie* 1976;**27**(4):505-508.

236 17. Gonsufotsin J, Tagnizukam D, Moyousomo R, et al.; Ultrasonographic study of
237 urological lesions of urinary bilharziasis in children in Barombi Kotto (Cameroon). *Semaine*
238 *Des Hopitaux* 1990;**66**(13):680-684.

- 239 18. Somo RM, Zukam DT, Kouamouo J, et al.; An epidemiological and radiological study
 240 of vesical schistosomiasis foci of the Barombi Lakes, Meme Division, Cameroon. *Bulletin De*
 241 *La Societe De Pathologie Exotique* 1987;**80**(5):813-822.
- 242 19. Christinet V, Lazdins-Helds JK, Stothard JR, et al.; Female genital schistosomiasis
 243 (FGS): from case reports to a call for concerted action against this neglected gynaecological
 244 disease. *International Journal for Parasitology* 2016;**46**(7):395-404. doi:
 245 10.1016/j.ijpara.2016.02.006.
- 246 20. Seto EYW, Sousa-Figueiredo JC, Betson M, et al.; Patterns of intestinal
 247 schistosomiasis among mothers and young children from Lake Albert, Uganda: water
 248 contact and social networks inferred from wearable global positioning system dataloggers.
 249 *Geospatial Health* 2012;**7**(1):1-13.
- 250 21. Stothard JR, Sousa-Figueiredo JC, Betson M, et al.; Investigating the spatial micro-
 251 epidemiology of diseases within a point-prevalence sample: a field applicable method for
 252 rapid mapping of households using low-cost GPS-dataloggers. *Transactions of the Royal*
 253 *Society of Tropical Medicine and Hygiene* 2011;**105**(9):500-506. doi:
 254 10.1016/j.trstmh.2011.05.007.
- 255 22. QGIS Development Team; QGIS Geographic Information System. Open Source
 256 Geospatial Foundation, 2009.
- 257 23. Seto EYW, Knapp F, Zhong B, et al.; The use of a vest equipped with a global
 258 positioning system to assess water-contact patterns associated with schistosomiasis.
 259 *Geospatial Health* 2007;**1**(2):233-241. doi: 10.4081/gh.2007.271.
- 260 24. Morris G, Conner LM; Assessment of accuracy, fix success rate, and use of estimated
 261 horizontal position error (EHPE) to filter inaccurate data collected by a common
 262 commercially available GPS logger. *Plos One* 2017;**12**(11). doi:
 263 10.1371/journal.pone.0189020.
- 264 25. R Core Team; R: A Language and Environment for Statistical Computing. Vienna,
 265 Austria: R Foundation for Statistical Computing, 2016.
- 266 26. Levitz S, Standley CJ, Adriko M, et al.; Environmental epidemiology of intestinal
 267 schistosomiasis and genetic diversity of *Schistosoma mansoni* infections in snails at Bugoigo
 268 village, Lake Albert. *Acta Tropica* 2013;**128**(2):284-291. doi:
 269 10.1016/j.actatropica.2012.10.003.
- 270 27. Rollinson D, Knopp S, Levitz S, et al.; Time to set the agenda for schistosomiasis
 271 elimination. *Acta Tropica* 2013;**128**(2):423-440. doi: 10.1016/j.actatropica.2012.04.013.

272

273 **Figure Legend**

274 Figure 1. **A.** The *S. haematobium* egg count frequency for the 21 individuals found positive
 275 at the time of community resurvey; **B.** Plot of water contact events over 24 hour period for
 276 mothers (n=12) and PSAC (n=12), the black lines denote average with 95% CIs others for
 277 mothers [27.4 (-1.3, 56.1)] and PSAC [14.1 (8.6, 19.7)] since our sample size was < 30 instead
 278 of using the formula of 1.96*Standard error, 2.201*standard error (11 degrees of freedom)
 279 was used. No significant difference was found between water contact events for mother
 280 and children groups ($P = 0.34$, *paired t-test*) evidencing similar water contact levels; **C.** GPS
 281 co-ordinates of individuals over a 48-hour period stratified by *S. haematobium* infection
 282 status and age with different colours representing individuals: (i) not infected PSAC (n=11);
 283 (ii) infected PSAC (n=1); (iii) uninfected mothers (n=10); and (iv) infected mothers (n=2). The

284 *S. haematobium* infected mothers (M1 and M2) and child (X4) from the GPS study are
285 identified in plots A and B.
286

287

288

289